

UNIVERSIDAD DE CONCEPCIÓN



CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA (CI²MA)



Hitting times for the stochastic wave equation with
fractional-colored noise

JORGE CLARKE, CIPRIAN A. TUDOR

PREPRINT 2012-11

SERIE DE PRE-PUBLICACIONES

Hitting times for the stochastic wave equation with fractional-colored noise

Jorge Clarke De la Cerda ^{1,2 *} Ciprian A. Tudor ^{2, †}

¹ CI²MA, Departamento de Ingeniería Matemática, Universidad de Concepción,
Casilla 160-C, Concepción, Chile.
jclarke@udec.cl

² Laboratoire Paul Painlevé, Université de Lille 1
F-59655 Villeneuve d'Ascq, France.
tudor@math.univ-lille1.fr

March 10, 2012

Abstract

We give sharp regularity results for the solution to the stochastic wave equation with linear fractional-colored noise. We apply these results in order to establish upper and lower bound for the hitting probabilities of the solution in terms of the Hausdorff measure and of the Newtonian capacity.

MSC 2000 subject classification: Primary 60H15; secondary 60H05

Keywords and phrases: stochastic wave equation, potential theory, hitting probability, capacity, Hausdorff dimension, spatially homogeneous Gaussian noise, fractional Brownian motion, Hölder continuity.

1 Introduction

The recent development of the stochastic calculus with respect to the fractional Brownian motion (fBm) naturally led to the study of stochastic partial differential equations (SPDEs) driven by this Gaussian process. The motivation comes from the wide area of applications of the fBm. We refer, among others, to [10], [11], [14], [16] and [18]. The purpose of our paper is to study the stochastic wave equation driven by fractional-colored Gaussian noise. Our work is situated somehow in the continuation of the line of research

*Partially supported by the MECESUP proyect UCO-0713 and the CONICYT-ECOS program C10E03.

†Associate member of the team Samm, Université de Paris 1 Panthéon-Sorbonne. Partially supported by the ANR grant "Mastérie" BLAN 012103.

which concerns SPDEs driven by the fBm but in the same time it follows the research initiated by Dalang in [4] which treats equations with white noise in time and correlated in space. More precisely, we consider a system of k stochastic wave equations

$$\frac{\partial^2 u_i}{\partial t^2}(t, x) = \Delta u_i(t, x) + \dot{W}_i(t, x), \quad t \in [0, T], x \in \mathbb{R}^d \quad (1)$$

with initial condition $u_i(t, x) = 0$ and $\frac{\partial u_i}{\partial t}(0, x) = 0$ for every $x \in \mathbb{R}^d$ and for every $i = 1, \dots, k$. The driving Gaussian process behaves as a fractional Brownian motion in time and has spatial covariance given by the Riesz kernel. More precisely

$$\mathbf{E}(W_i(t, A)W_j(s, B)) = \delta_{i,j}R_H(t, s) \int_A \int_B f(x - y)dx dy$$

for every $t, s \in [0, T]$ and A, B Borel sets in \mathbb{R}^d where $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is the Fourier transform of a non-negative tempered measure μ on \mathbb{R}^d whose density with respect to the Lebesgue measure is $|\xi|^{-(d-\beta)}$, $0 < \beta < d$. Above $\delta_{i,j}$ denotes the Kronecker symbol.

The equation (1) has been recently studied in [2]. It has been proven that (1) admits a unique mild solution if and only if $\beta < 2H + 1$ which extends the result obtained in [4] in the case $H = \frac{1}{2}$. The purpose of this work is to analyze further the solution of (1). We will actually give sharp results for the regularity of it, in time and in space, and we apply these regularity results to study the hitting probabilities for the solution u to (1). More precisely, given a Borel set $A \subset \mathbb{R}^k$ we want to determine whether the process $(u(t, x), t \in [0, T], x \in \mathbb{R}^d)$ hits the set A with positive probability. Recently, there has been several papers on hitting probabilities, and more generally speaking, on potential theory for systems of SPDEs. We refer, among others, to [5], [6], [7], [8] or [13]. The study of hitting probabilities for stochastic partial differential equations with fractional noise in time is new. As far as we know, only the paper [15] treated this problem. Actually, in this reference the authors give upper and lower bounds for the hitting times of solution to a system of stochastic heat equations on the circle with fractional noise in time.

Our aim is to make a new step in this research direction. As we mentioned before we make a potential analysis of the solution to the stochastic wave equation with fractional-colored noise. That means, the noise behaves as the fractional Brownian motion with respect to the time variable and it is a "colored" non-white spatial covariance. In our work this spatial covariance will be described by the Riesz kernel. It is known classical the fact that in order to obtain results on the hitting times of a stochastic process, a detailed analysis of the behavior of the increments of the process is needed. We address this question in our paper and we find the following: the solution $u(t, x), t \in [0, T], x \in \mathbb{R}^d$ to (1) is Hölder continuous of order $2H + 1 - \beta$, $\beta \in (2H - 1, d \wedge 2H + 1)$ in time as well as with respect to the space variable. This generalizes the result obtained in [8] and [9] for the wave equation with white noise in time and Riesz covariance in space. Although the main lines of our work follows the approach of [9], we stress that, as usually, the fractional cases involves more complex calculation and the techniques used in the standard white noise case need to be substantially adapted. this is mainly due to the nature of the noise and to the structure

of the Gaussian space associated to the noise. We will point out later in our paper, how the fractional noise involves more complexity than in e.g. [6] or [9]. Moreover, the study of the solution to the wave equation is generally recognized to be more difficult than e.g. the solution to the heat equation, due to the appearance of the trigonometric functions and this is also the case in our work.

We mention that there are more or less general criteria to determine the hitting times for a stochastic process. Such criteria have been given in [3], [6], [7] or [9] among others. We will use the approach in [3] because it concerns Gaussian processes and fits well with our context (note that the solution to (1) is Gaussian).

Our paper is structured as follows. Section 2 contains some preliminaries, we briefly describe the basic properties of the Gaussian noise and its associated Hilbert space, we list the elements of the potential theory that we will use in our paper and we will recall some fact related to the solution to the stochastic wave equation with fractional-colored noise. In Section 3 we analyze the Hölder regularity of the solution with respect to its time and space variables. Section 4 is devoted to the study of the hitting probabilities for this solution, based on a criterium in [3].

2 Preliminaries

This section is devoted to introduce the basic notion that we will need throughout the paper. We first introduce the canonical Hilbert space associated to the fractional-colored Gaussian noise. In the second part we present the basic elements related to the potential theory that intervene in the last section.

2.1 The canonical Hilbert space

We denote by $C_0^\infty(\mathbb{R}^{d+1})$ the space of infinitely differentiable functions on \mathbb{R}^{d+1} with compact support, and $\mathcal{S}(\mathbb{R}^d)$ the Schwartz space of rapidly decreasing C^∞ functions in \mathbb{R}^d . For $\varphi \in L^1(\mathbb{R}^d)$, we let $\mathcal{F}\varphi$ be the Fourier transform of φ :

$$\mathcal{F}\varphi(\xi) = \int_{\mathbb{R}^d} e^{-i\xi \cdot x} \varphi(x) dx.$$

We begin by introducing the framework of [4]. Let μ be a non-negative tempered measure on \mathbb{R}^d , i.e. a non-negative measure which satisfies:

$$\int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^l \mu(d\xi) < \infty, \quad \text{for some } l > 0. \quad (2)$$

Since the integrand is non-increasing in l , we may assume that $l \geq 1$ is an integer. Note that $1 + |\xi|^2$ behaves as a constant around 0, and as $|\xi|^2$ at ∞ , and hence (2) is equivalent to:

$$\int_{|\xi| \leq 1} \mu(d\xi) < \infty, \quad \text{and} \quad \int_{|\xi| \geq 1} \mu(d\xi) \frac{1}{|\xi|^{2l}} < \infty, \quad \text{for some integer } l \geq 1. \quad (3)$$

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ be the Fourier transform of μ in $\mathcal{S}'(\mathbb{R}^d)$, i.e.

$$\int_{\mathbb{R}^d} f(x)\varphi(x)dx = \int_{\mathbb{R}^d} \mathcal{F}\varphi(\xi)\mu(d\xi), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^d).$$

Simple properties of the Fourier transform show that for any $\varphi, \psi \in \mathcal{S}(\mathbb{R}^d)$,

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \varphi(x)f(x-y)\psi(y)dxdy = \int_{\mathbb{R}^d} \mathcal{F}\varphi(\xi)\overline{\mathcal{F}\psi(\xi)}\mu(d\xi).$$

An approximation argument shows that the previous equality also holds for indicator functions $\varphi = 1_A, \psi = 1_B$, with $A, B \in \mathcal{B}_b(\mathbb{R}^d)$, where $\mathcal{B}_b(\mathbb{R}^d)$ is the class of bounded Borel sets of \mathbb{R}^d :

$$\int_A \int_B f(x-y)dxdy = \int_{\mathbb{R}^d} \mathcal{F}1_A(\xi)\overline{\mathcal{F}1_B(\xi)}\mu(d\xi). \quad (4)$$

Now we introduce the *fractional Brownian motion* (fBm) with Hurst index $H \in (0, 1)$. This is a zero-mean Gaussian process $(B_t^H)_{t \in [0, T]}$ with covariance

$$R_H(t, s) := \frac{1}{2} (t^{2H} + s^{2H} - |t - s|^{2H}), \quad t, s \in [0, T].$$

Let us denote by \mathcal{H} the canonical Hilbert space associated with this Gaussian process. This canonical Hilbert space is defined as the closure of the linear space generated by the indicator functions $1_{[0, t]}, t \in [0, T]$ with respect to the inner product

$$\langle 1_{[0, t]}, 1_{[0, s]} \rangle_{\mathcal{H}} = R_H(t, s).$$

It is well known that for $H > 1/2$ we have the expression

$$R_H(t, s) = \alpha_H \int_0^t \int_0^s |u - v|^{2H-2} dudv \quad (5)$$

for every $s, t \in [0, T]$ with $\alpha_H := H(2H - 1)$. More generally, for $H > 1/2$ and every $\psi, \phi \in \mathcal{H} = \mathcal{H}([0, T])$ we have

$$\langle \psi, \phi \rangle_{\mathcal{H}} = \alpha_H \int_0^T \int_0^T \psi(u)\phi(v)|u - v|^{2H-2} dudv \quad (6)$$

As in [1], on a complete probability space (Ω, \mathcal{F}, P) , we consider a zero-mean Gaussian process $W = \{W_t(A); t \geq 0, A \in \mathcal{B}_b(\mathbb{R}^d)\}$ with covariance:

$$\mathbf{E}(W_t(A)W_s(B)) = R_H(t, s) \int_A \int_B f(x-y)dxdy =: \langle 1_{[0, t] \times A}, 1_{[0, s] \times B} \rangle_{\mathcal{HP}}. \quad (7)$$

Let \mathcal{E} be the set of linear combinations of elementary functions $1_{[0, t] \times A}$, $t \geq 0, A \in \mathcal{B}_b(\mathbb{R}^d)$, and \mathcal{HP} be the Hilbert space defined as the closure of \mathcal{E} with respect to the inner

product $\langle \cdot, \cdot \rangle_{\mathcal{HP}}$. (Alternatively, \mathcal{HP} can be defined as the completion of $C_0^\infty(\mathbb{R}^{d+1})$, with respect to the inner product $\langle \cdot, \cdot \rangle_{\mathcal{HP}}$; see [1].)

The map $1_{[0,t] \times A} \mapsto W_t(A)$ is an isometry between \mathcal{E} and the Gaussian space H^W of W , which can be extended to \mathcal{HP} . We denote this extension by:

$$\varphi \mapsto W(\varphi) = \int_0^\infty \int_{\mathbb{R}^d} \varphi(t, x) W(dt, dx).$$

In the present work, we assume that $H > 1/2$. Hence, (5) holds. From (4) and (5), it follows that for any $\varphi, \psi \in \mathcal{E}$,

$$\begin{aligned} \langle \varphi, \psi \rangle_{\mathcal{HP}} &= \alpha_H \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \varphi(u, x) \psi(v, y) f(x - y) |u - v|^{2H-2} dx \, dy \, du \, dv \\ &= \alpha_H \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \mathcal{F}\varphi(u, \cdot)(\xi) \overline{\mathcal{F}\psi(v, \cdot)(\xi)} |u - v|^{2H-2} \mu(d\xi) \, du \, dv. \end{aligned}$$

Moreover, we can interchange the order of the integrals $dudv$ and $\mu(d\xi)$, since for indicator functions φ and ψ , the integrand is a product of a function of (u, v) and a function of ξ . Hence, for $\varphi, \psi \in \mathcal{E}$, we have:

$$\langle \varphi, \psi \rangle_{\mathcal{HP}} = \alpha_H \int_{\mathbb{R}^d} \int_0^\infty \int_0^\infty \mathcal{F}\varphi(u, \cdot)(\xi) \overline{\mathcal{F}\psi(v, \cdot)(\xi)} |u - v|^{2H-2} du \, dv \, \mu(d\xi). \quad (8)$$

The space \mathcal{HP} may contain distributions, but contains the space $|\mathcal{HP}|$ of measurable functions $\varphi : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$\|\varphi\|_{|\mathcal{HP}|}^2 := \alpha_H \int_0^\infty \int_0^\infty \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |\varphi(u, x)| |\varphi(v, y)| f(x - y) |u - v|^{2H-2} dx \, dy \, du \, dv < \infty.$$

2.2 Elements of the potential theory

Our aim is to analyze the probability

$$P(u(I) \cap A) \neq \emptyset$$

where u is the solution to (1), I is a Borel set included in $[0, T] \times \mathbb{R}^d$ and A is a Borel set in \mathbb{R}^k . Here $u(I)$ means the image of I under the random map $(t, x) \rightarrow u(t, x)$.

We will briefly present the notion of the potential theory that we will need in our paper. For all Borel sets $F \subset \mathbb{R}^d$ we define $\mathcal{P}(F)$ to be the set of all probability measures with compact support included in F . For all $\mu \in \mathcal{P}(\mathbb{R}^d)$, let us denote by $I_\beta(\mu)$ the so-called β -energy of the measure μ defined by

$$I_\beta(\mu) = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} K_\beta(\|x - y\|) \mu(dx) \mu(dy) \quad (9)$$

where

$$K_\beta(r) = \begin{cases} r^{-\beta} & \text{if } \beta > 0; \\ \log\left(\frac{N_0}{r}\right) & \text{if } \beta = 0; \\ 1 & \text{if } \beta < 0. \end{cases} \quad (10)$$

Here N_0 is a constant.

For all $\beta \in \mathbb{R}$ and $F \in \mathcal{B}(\mathbb{R}^d)$ we define the β -dimensional capacity of F by

$$\text{Cap}_\beta(F) = \left[\inf_{\mu \in \mathcal{P}(F)} I_\beta(\mu) \right]^{-1} \quad (11)$$

with the convention $1/\infty := 0$. The β -dimensional Hausdorff measure of the set $F \in \mathcal{B}(\mathbb{R}^d)$ is given by

$$\mathcal{H}_\beta(F) = \lim_{\varepsilon \rightarrow 0^+} \inf \left[\sum_{i=1}^{\infty} (2r_i)^\beta; F \subset \bigcup_{i=1}^{\infty} B(x_i, r_i), \sup_{i \geq 1} r_i \leq \varepsilon \right] \quad (12)$$

where $B(x, r)$ denotes the Euclidean ball of radius $r > 0$ centered at $x \in \mathbb{R}^d$. When $\beta < 0$, the β -dimensional Hausdorff measure of F is infinite by definition.

2.3 The stochastic wave equation with linear fractional-colored noise

Consider the linear stochastic wave equation driven by an infinite-dimensional fractional Brownian motion W with Hurst parameter $H \in (0, 1)$. That is

$$\begin{aligned} \frac{\partial^2 u}{\partial t^2}(t, x) &= \Delta u(t, x) + \dot{W}(t, x), \quad t > 0, x \in \mathbb{R}^d \\ u(0, x) &= 0, \quad x \in \mathbb{R}^d \\ \frac{\partial u}{\partial t}(0, x) &= 0, \quad x \in \mathbb{R}^d. \end{aligned} \quad (13)$$

Here Δ is the Laplacian on \mathbb{R}^d and $W = \{W_t(A); t \geq 0, A \in \mathcal{B}_b(\mathbb{R}^d)\}$ is a centered Gaussian field with covariance

$$\mathbf{E}(W_t(A)W_s(B)) = R_H(t, s) \int_A \int_B f(x - y) dx dy,$$

where R_H is the covariance of the fractional Brownian motion and f is the Riesz kernel.

Let G_1 be the fundamental solution of $u_{tt} - \Delta u = 0$. It is known that $G_1(t, \cdot)$ is a distribution in $\mathcal{S}'(\mathbb{R}^d)$ with rapid decrease, and

$$\mathcal{F}G_1(t, \cdot)(\xi) = \frac{\sin(t|\xi|)}{|\xi|}, \quad (14)$$

for any $\xi \in \mathbb{R}^d, t > 0, d \geq 1$ (see e.g. [19]). In particular,

$$\begin{aligned} G_1(t, x) &= \frac{1}{2} 1_{\{|x| < t\}}, \quad \text{if } d = 1 \\ G_1(t, x) &= \frac{1}{2\pi} \frac{1}{\sqrt{t^2 - |x|^2}} 1_{\{|x| < t\}}, \quad \text{if } d = 2 \\ G_1(t, x) &= c_d \frac{1}{t} \sigma_t, \quad \text{if } d = 3, \end{aligned}$$

where σ_t denotes the surface measure on the 3-dimensional sphere of radius t .

The solution of (13) is a square-integrable process $u = \{u(t, x); t \geq 0, x \in \mathbb{R}^d\}$ defined by:

$$u(t, x) = \int_0^t \int_{\mathbb{R}^d} G_1(t - s, x - y) W(ds, dy). \quad (15)$$

By definition, $u(t, x)$ exists if and only if the stochastic integral above is well-defined, i.e. $g_{tx} := G_1(t - \cdot, x - \cdot) \in \mathcal{HP}$. In this case, $\mathbf{E}|u(t, x)|^2 = \|g_{tx}\|_{\mathcal{HP}}^2$.

The following result has been proved in [2].

Theorem 1 *The stochastic wave equation (13) admits an unique mild solution $(u(t, x))_{t \in [0, T], x \in \mathbb{R}^d}$ if and only if*

$$\int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^{H + \frac{1}{2}} \mu(d\xi) < \infty. \quad (16)$$

Remark 1 *Note that (16) is equivalent to:*

$$\int_{|\xi| \leq 1} \mu(d\xi) < \infty, \quad \text{and} \quad \int_{|\xi| \geq 1} \mu(d\xi) \frac{1}{|\xi|^{2H+1}} < \infty. \quad (17)$$

As mentioned in the Introduction, we will consider throughout the paper that the spatial covariance of the noise W is given by the Riezs kernel. That means the measure μ is

$$d\mu(\xi) = |\xi|^{-d+\beta} d\xi \text{ with } \beta \in (0, d).$$

In this case the kernel f is given by

$$f(\xi) = |\xi|^{-\beta} \text{ with } \beta \in (0, d).$$

Note that in the case of the Riesz kernel, condition (16) is equivalent to

$$\beta \in (0, d \wedge (2H + 1)). \quad (18)$$

Remark 2 *Since $H > \frac{1}{2}$ and so $2H + 1 \in (2, 3)$, for dimension $d = 1, 2$ we have $\beta \in (0, d)$ while for $d \geq 3$ we have $\beta \in (0, 2H + 1)$.*

3 Regularity of the solution

3.1 Time regularity

In this part we will focus our attention on the behavior of the increments of the solution $u(t, x)$ with respect to the variable t . We will give upper and lower bounds for the L^2 -norm of this increment. Usually, obtaining upper bounds is recognized to be easier than obtaining lower bounds, this is also the case in our work. Actually, to get the sharpness of the regularity of u with respect to the time variable, we need to impose a stronger assumption than (18) on the parameters β and H (condition (19) below). This is due to the characteristics of the scalar product in the \mathcal{HP} .

We will start with the following useful lemma that gives an explicit expression for the \mathcal{H} norm of the cosine and sine functions. These norms will widely appear further in our computations.

Lemma 1 *Let $f(x) = \cos(x)$ and $g(x) = \sin x$ for $x \in \mathbb{R}$. Then for every $a, b \in \mathbb{R}$, $a < b$*

$$\|f1_{(a,b)}\|_{\mathcal{H}}^2 = \alpha_H \int_0^{b-a} dv \cos(v) v^{2H-2} (b-a-v) + \alpha_H \cos(a+b) \int_0^{b-a} dv v^{2H-2} \sin(b-a-v).$$

and

$$\|g1_{(a,b)}\|_{\mathcal{H}}^2 = \alpha_H \int_0^{b-a} dv \cos(v) v^{2H-2} (b-a-v) - \alpha_H \cos(a+b) \int_0^{b-a} dv v^{2H-2} \sin(b-a-v).$$

Proof: Using the expression of the scalar product in the Hilbert space \mathcal{H} and the trigonometric identities

$$\begin{aligned} \cos(u \pm v) &= \cos u \cos v \mp \sin u \sin v \quad \text{and} \\ \sin(x) - \sin(y) &= 2\cos\left(\frac{x+y}{2}\right)\sin\left(\frac{x-y}{2}\right) \end{aligned}$$

we can write

$$\begin{aligned} \|f1_{(a,b)}\|_{\mathcal{H}}^2 + \|g1_{(a,b)}\|_{\mathcal{H}}^2 &= \alpha_H \int_a^b \int_a^b |u-v|^{2H-2} (\cos u \cos v + \sin u \sin v) du dv \\ &= \alpha_H \int_a^b du \int_a^b dv |u-v|^{2H-2} \cos(u-v) \\ &= 2\alpha_H \int_a^b du \int_0^{u-a} dv \cos(v) v^{2H-2} \\ &= 2\alpha_H \int_0^{b-a} dv \cos(v) v^{2H-2} (b-a-v) \end{aligned}$$

where we made the change of variables $\tilde{v} = u - v$ in the integral dv above and we computed the integral du . Similarly

$$\begin{aligned}\|f1_{(a,b)}\|_{\mathcal{H}}^2 - \|g1_{(a,b)}\|_{\mathcal{H}}^2 &= \alpha_H \int_a^b du \int_a^b dv |u - v|^{2H-2} (\cos u \cos v - \sin u \sin v) \\ &= \alpha_H \int_a^b \int_a^b |u - v|^{2H-2} \cos(u + v) dudv\end{aligned}$$

and by the change of variable $\tilde{v} = u - v$ in the integral dv ,

$$\begin{aligned}\|f1_{(a,b)}\|_{\mathcal{H}}^2 - \|g1_{(a,b)}\|_{\mathcal{H}}^2 &= 2\alpha_H \int_a^b du \int_0^{u-a} dv \cos(2u - v) v^{2H-2} \\ &= 2\alpha_H \int_0^{b-a} dv v^{2H-2} \int_{v+a}^b du \cos(2u - v) \\ &= \alpha_H \int_0^{b-a} dv v^{2H-2} (\sin(2b - v) - \sin(2a + v)) \\ &= 2\alpha_H \cos(a + b) \int_0^{b-a} dv v^{2H-2} \sin(b - a - v).\end{aligned}$$

■

Remark 3 As a consequence of the Lemma 1 we deduce the following

i. For any $x > 0$ the quantity $\int_0^x v^{2H-2} \cos(v)(x - v)dv$ is positive (it is the sum of two norms).

ii. For every $a, b \in \mathbb{R}$, $a < b$

$$\|f1_{(a,b)}\|_{\mathcal{H}}^2 \leq 2\alpha_H \int_0^{b-a} dv \cos(v) v^{2H-2} (b - a - v)$$

iii. For every $a, b \in \mathbb{R}$, $a < b$

$$\|f1_{(a,b)}\|_{\mathcal{H}}^2 \geq 2\alpha_H \cos(a + b) \int_0^{b-a} dv v^{2H-2} \sin(b - a - v).$$

Later, we use also the following lemma.

Lemma 2 For every $a, b \in \mathbb{R}$ with $a < b$,

$$\int_a^b \int_a^b dudv \sin(u - v) |u - v|^{2H-2} = 0.$$

Proof: This follows from the trivial equality

$$\int_a^b \int_a^b \sin(u) \cos(v) |u - v|^{2H-2} dudv = \int_a^b \int_a^b \sin(v) \cos(u) |u - v|^{2H-2} dudv.$$

■

Concretely, we will prove the following result concerning the time regularity of the solution to (13). We mention that, in the rest of our paper, c, C, \dots will denote generic positive constants that may change from line to line.

Proposition 1 *Assume that*

$$\beta \in (2H - 1, d \wedge (2H + 1)). \quad (19)$$

Let $t_0, M > 0$ and fix $x \in [-M, M]^d$. Then there exists a positive constants c_1, c_2 such that for every $s, t \in [t_0, T]$

$$c_1 |t - s|^{2H+1-\beta} \leq \mathbf{E} |u(t, x) - u(s, x)|^2 \leq c_2 |t - s|^{2H+1-\beta}.$$

Proof: Let $h > 0$ and let us estimate the $L^2(\Omega)$ -norm of the increment $u(t+h, x) - u(t, x)$. Splitting the interval $[0, t+h]$ into the intervals $[0, t]$ and $[t, t+h]$, and using the inequality $|a+b|^2 \leq 2(a^2 + b^2)$, we obtain:

$$\begin{aligned} \mathbf{E} |u(t+h, x) - u(t, x)|^2 &\leq 2\{\|(g_{t+h,x} - g_{t,x})1_{[0,t]}\|_{\mathcal{HP}}^2 + \|g_{t+h,x}1_{[t,t+h]}\|_{\mathcal{HP}}^2\} \\ &=: 2[E_{1,t}(h) + E_2(h)]. \end{aligned} \quad (20)$$

The first summand can be handled in the following way.

$$\begin{aligned} E_{1,t}(h) &= \alpha_H \int_{\mathbb{R}^d} \mu(d\xi) \int_0^t \int_0^t dv dv |u - v|^{2H-2} \mathcal{F}(g_{t+h,x} - g_{tx})(u, \cdot)(\xi) \\ &\quad \times \overline{\mathcal{F}(g_{t+h,x} - g_{tx})(v, \cdot)(\xi)} \\ &= \alpha_H \int_{\mathbb{R}^d} \mu(d\xi) \int_0^t \int_0^t du dv |u - v|^{2H-2} [\mathcal{F}G_1(u+h, \cdot)(\xi) - \mathcal{F}G_1(u, \cdot)(\xi)] \\ &\quad \times \overline{\mathcal{F}G_1(v+h, \cdot)(\xi) - \mathcal{F}G_1(v, \cdot)(\xi)} \\ &= \alpha_H \int_0^t \int_0^t du dv |u - v|^{2H-2} I_h, \end{aligned}$$

where

$$\begin{aligned} I_h &= \int_{\mathbb{R}^d} \mu(d\xi) [\mathcal{F}G_1(u+h, \cdot)(\xi) - \mathcal{F}G_1(u, \cdot)(\xi)] [\overline{\mathcal{F}G_1(v+h, \cdot)(\xi) - \mathcal{F}G_1(v, \cdot)(\xi)}] \\ &= \int_{\mathbb{R}^d} \mu(d\xi) \frac{(\sin((u+h)|\xi|) - \sin(u|\xi|))}{|\xi|} \frac{(\sin((v+h)|\xi|) - \sin(v|\xi|))}{|\xi|}. \end{aligned}$$

Using the last trigonometric identity presented before we obtain

$$\begin{aligned}
E_{1,t}(h) &= \alpha_H \int_0^t \int_0^t dudv |u-v|^{2H-2} \int_{\mathbb{R}^d} \mu(d\xi) \frac{\sin(\frac{h|\xi|}{2})^2}{|\xi|^2} \cos(\frac{(2u+h)|\xi|}{2}) \cos(\frac{(2v+h)|\xi|}{2}) \\
&= c\alpha_H \int_0^t \int_0^t dudv |u-v|^{2H-2} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2}} \sin(h|\xi|)^2 \cos((2u+h)|\xi|) \cos((2v+h)|\xi|),
\end{aligned}$$

and by making the change of variables $\tilde{u} = (2u+h)|\xi|$, $\tilde{v} = (2v+h)|\xi|$,

$$\begin{aligned}
E_{1,t}(h) &= c \cdot \alpha_H \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \int_{h|\xi|}^{(2t+h)|\xi|} \int_{h|\xi|}^{(2t+h)|\xi|} dudv |u-v|^{2H-2} \cos u \cos v \\
&= c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \|\cos(\cdot) 1_{(h|\xi|, (2t+h)|\xi|)}(\cdot)\|_{\mathcal{H}}^2,
\end{aligned} \tag{21}$$

and using Lemma 1,

$$\begin{aligned}
E_{1,t}(h) &= c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \times \left[\int_0^{2t|\xi|} \cos(v) v^{2H-2} (2t|\xi| - v) dv \right. \\
&\quad \left. + \cos(2t|\xi| + 2h|\xi|) \int_0^{2t|\xi|} v^{2H-2} (\sin(2t|\xi| - v)) \right] \\
&= c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \times \left[2t|\xi| \int_0^{2t|\xi|} \cos(v) v^{2H-2} dv \right. \\
&\quad \left. - \sin(2t|\xi|) (2t|\xi|)^{2H-1} + (2H-1) \int_0^{2t|\xi|} \sin(v) v^{2H-2} dv \right. \\
&\quad \left. + \cos(2t|\xi| + 2h|\xi|) \int_0^{2t|\xi|} v^{2H-2} (\sin(2t|\xi| - v)) \right]
\end{aligned} \tag{22}$$

where we use integration by parts. By Remark 3, point ii. we have the upper bound

$$\begin{aligned}
E_{1,t}(h) &\leq c \cdot \alpha_H \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \\
&\quad \times \left[2t|\xi| \int_0^{2t|\xi|} \cos(v) v^{2H-2} dv - \sin(2t|\xi|) (2t|\xi|)^{2H-1} \right. \\
&\quad \left. + (2H-1) \int_0^{2t|\xi|} \sin(v) v^{2H-2} dv \right].
\end{aligned}$$

We will treat separately the three summands above. Concerning the first one,

$$\begin{aligned}
& \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 2t|\xi| \int_0^{2t|\xi|} \cos(v) v^{2H-2} dv \\
&= c_{t,H} h^{2H+1-\beta} \left| \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+1}} \sin(|\xi|)^2 \int_0^{\frac{2t|\xi|}{h}} \cos(v) v^{2H-2} dv \right| \\
&\leq c_{t,H} h^{2H+1-\beta} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+1}} \sin(|\xi|)^2 \left| \int_0^{\frac{2t|\xi|}{h}} \cos(v) v^{2H-2} dv \right| \\
&\leq c_{t,H} h^{2H+1-\beta}
\end{aligned}$$

using condition (18) and the fact that the integral $\int_0^\infty \cos(v) v^{2H-2} dv$ is convergent (this implies that the function $x \in [0, \infty) \rightarrow \int_0^x \cos(v) v^{2H-2} dv$ admits a limit at infinity and it is therefore bounded). On the other hand

$$\begin{aligned}
& \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \sin(2t|\xi|) (2t|\xi|)^{2H-1} = c_t h^{3-\beta} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+3}} \sin(|\xi|)^2 \sin\left(\frac{2t|\xi|}{h}\right) \\
&= c_t h^{3-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+3}} \sin(|\xi|)^2 \sin\left(\frac{2t|\xi|}{h}\right) + c_t h^{3-\beta} \int_{|\xi| > 1} \frac{d\xi}{|\xi|^{d-\beta+3}} \sin(|\xi|)^2 \sin\left(\frac{2t|\xi|}{h}\right).
\end{aligned}$$

The second part over the region $|\xi| \geq 1$ is bounded by $ch^{3-\beta}$ simply by majorizing sinus by one. The second integral has a singularity for $|\xi|$ close to zero. Using that $\sin(x) \leq x$ for all $x \geq 0$, we will bound it above by

$$\begin{aligned}
& h^{3-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+3}} \sin(|\xi|)^2 \sin\left(\frac{2t|\xi|}{h}\right) \\
&\leq c_t h^{3-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+3}} |\xi|^2 \left| \sin\left(\frac{2t|\xi|}{h}\right) \right|^{2-2H} \left| \sin\left(\frac{2t|\xi|}{h}\right) \right|^{2H-1} \\
&\leq c_t h^{2H+1-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2H-1}}
\end{aligned}$$

where we bounded $\left| \sin\left(\frac{2t|\xi|}{h}\right) \right|^{2-2H}$ by $c_t (|\xi| h^{-1})^{2-2H}$ and $\left| \sin\left(\frac{2t|\xi|}{h}\right) \right|^{2H-1}$ by 1. The last

integral is finite since $\beta > 2H - 1$ (assumption (19)). Finally

$$\begin{aligned}
& \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \int_0^{2t|\xi|} \sin(v) v^{2H-2} dv \\
&= h^{2H+2-\beta} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(|\xi|)^2 \int_0^{\frac{2t|\xi|}{h}} \sin(v) v^{2H-2} dv \\
&= h^{2H+2-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(|\xi|)^2 \int_0^{\frac{2t|\xi|}{h}} \sin(v) v^{2H-2} dv \\
&\quad + h^{2H+2-\beta} \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(|\xi|)^2 \int_0^{\frac{2t|\xi|}{h}} \sin(v) v^{2H-2} dv \\
&\leq h^{2H+2-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} |\xi|^2 \int_0^{\frac{2t|\xi|}{h}} |\sin v| v^{2H-2} dv \\
&\quad + h^{2H+2-\beta} \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \int_0^{\frac{2t|\xi|}{h}} \sin(v) v^{2H-2} dv.
\end{aligned} \tag{23}$$

Using again the fact that $\int_0^\infty \sin(v) v^{2H-2} dv$ is convergent is easy to see that the integral over the region $|\xi| \geq 1$ is bounded by $c_t h^{2H+2-\beta}$. For the integral over $|\xi| \leq 1$ we make the change of variables $\tilde{v} = \frac{v|\xi|}{h}$ and we get

$$\begin{aligned}
& h^{3-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+1}} \int_0^{2t} \left| \sin\left(\frac{v|\xi|}{h}\right) \right| v^{2H-2} dv \\
&= h^{3-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+1}} \int_0^{2t} \left| \sin\left(\frac{v|\xi|}{h}\right) \right|^{2-2H} \left| \sin\left(\frac{v|\xi|}{h}\right) \right|^{2H-1} v^{2H-2} dv \\
&\leq c_t h^{2h+1-\beta} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2H-1}},
\end{aligned}$$

where we have made the same considerations as for the second summand in the decomposition of $E_{1,t}(h)$. In this way, we obtained the upper bound for the summand $E_{1,t}(h)$ in (20)

$$E_{1,t}(h) \leq C h^{2H+1-\beta}. \tag{24}$$

We study now the term $E_2(h)$ in (20) (its notation $E_2(h)$ instead of $E_{2,t}(h)$ is due to the fact that it does not depend on t , see below). Using successively the change of variables $\tilde{u} = \frac{u}{h}, \tilde{v} = \frac{v}{h}$ in the integral $du dv$ and $\tilde{\xi} = h\xi$ in the integral $d\xi$, the summand $E_2(h)$ can

be written as

$$\begin{aligned}
E_2(h) &= \alpha_H \int_{\mathbb{R}^d} \int_t^{t+h} \int_t^{t+h} \mathcal{F}G_1(t+h-u, \cdot)(\xi) \overline{\mathcal{F}G_1(t+h-v, \cdot)(\xi)} |u-v|^{2H-2} du \, dv \, \mu(d\xi) \\
&= \alpha_H \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{|\xi|^2} \int_0^h \int_0^h \sin(u|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv \\
&= \alpha_H h^{2H} \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{|\xi|^2} \int_0^1 \int_0^1 \sin(u|\xi|h) \sin(v|\xi|h) |u-v|^{2H-2} dudv \\
&= \alpha_H h^{2H+2-\beta} \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{|\xi|^2} \int_0^1 \int_0^1 \sin(u|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv.
\end{aligned}$$

Let us use the following notation:

$$N_t(\xi) = \frac{\alpha_H}{|\xi|^2} \int_0^t \int_0^t \sin(u|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv, \quad t \in [0, T], \xi \in \mathbb{R}^d. \quad (25)$$

By Proposition 3.7 in [2] the term

$$N_1(\xi) = \frac{\alpha_H}{|\xi|^2} \int_0^1 \int_0^1 \sin(u|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv$$

satisfies the inequality

$$N_1(\xi) \leq C_H \left(\frac{1}{1+|\xi|^2} \right)^{H+1/2},$$

with C_H a positive constant not depending on h . Consequently the term $E_2(h)$ is bounded by

$$E_2(h) \leq Ch^{2H+2-\beta} \int_{\mathbb{R}^d} \left(\frac{1}{1+|\xi|^2} \right)^{H+\frac{1}{2}} \mu(d\xi) \quad (26)$$

and this is clearly finite due to (18). Relations (24) and (26) give the first part of the conclusion.

Let us analyze now the lower bound of the increments of $u(t, x)$ with respect to the variable t . Let $h > 0, x \in [-M, M]^d$ and $t \in [t_0, T]$ such that $t+h \in [t_0, T]$. From the decomposition

$$\begin{aligned}
\mathbf{E} |u(t+h, x) - u(t, x)|^2 &= \| (g_{t+h, x} - g_{t, x}) 1_{[0, t]} \|_{\mathcal{HP}}^2 + \| g_{t+h, x} 1_{[t, t+h]} \|_{\mathcal{HP}}^2 \\
&\quad + 2 \langle (g_{t+h, x} - g_{t, x}) 1_{[0, t]}, g_{t+h, x} 1_{[t, t+h]} \rangle_{\mathcal{HP}}
\end{aligned}$$

we immediately obtain, since the second summand in the right-hand side is positive,

$$\begin{aligned}
\mathbf{E} |u(t+h, x) - u(t, x)|^2 &\geq \| (g_{t+h, x} - g_{t, x}) 1_{[0, t]} \|_{\mathcal{HP}}^2 + 2 \langle (g_{t+h, x} - g_{t, x}) 1_{[0, t]}, g_{t+h, x} 1_{[t, t+h]} \rangle_{\mathcal{HP}} \\
&:= E_{1,t}(h) + E_{3,t}(h).
\end{aligned}$$

We can assume, without any loss of the generality, that $t = \frac{1}{2}$. Denote $E_{1, \frac{1}{2}}(h) := E_1(h)$. We first prove that

$$E_1(h) \geq ch^{2H+1-\beta} - c'h^{2H+2-\beta}. \quad (27)$$

for h small enough. Recall that we have an exact expression for $E_1(h)$ (see (22)). Actually,

$$\begin{aligned} E_1(h) &= \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \|\cos(\cdot) 1_{(h|\xi|, h|\xi|+|\xi|)}\|_{\mathcal{H}}^2 \\ &= \alpha_H \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \int_{h|\xi|}^{(1+h)|\xi|} \int_{h|\xi|}^{(1+h)|\xi|} dudv |u-v|^{2H-2} \cos u \cos v \\ &= \alpha_H \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \int_0^{|\xi|} \int_0^{|\xi|} dudv \cos(u+h|\xi|) \cos(v+h|\xi|) |u-v|^{2H-2}. \end{aligned}$$

By the trigonometric formula $\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$ we can write

$$\begin{aligned} E_1(h) &= \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 \left[\cos(h|\xi|)^2 \int_0^{|\xi|} \int_0^{|\xi|} dudv \cos u \cos v |u-v|^{2H-2} \right. \\ &\quad \left. - 2 \sin(h|\xi|) \cos(h|\xi|) \int_0^{|\xi|} \int_0^{|\xi|} dudv \sin u \cos v |u-v|^{2H-2} \right. \\ &\quad \left. + \sin(h|\xi|)^2 \int_0^{|\xi|} \int_0^{|\xi|} dudv \sin u \sin v |u-v|^{2H-2} \right] \\ &:= A + B + C. \end{aligned}$$

We will neglect the first term since it is positive. We will bound the second one above by $ch^{2H+2-\beta}$. We have (we use Lemma 2 at the third line below)

$$\begin{aligned} &\sin(h|\xi|) \cos(h|\xi|) \int_0^{|\xi|} \int_0^{|\xi|} dudv \sin u \cos v |u-v|^{2H-2} \\ &\leq |\sin(2h|\xi|)| \left| \int_0^{|\xi|} \int_0^{|\xi|} dudv \sin u \cos v |u-v|^{2H-2} \right| \\ &= \frac{1}{2} |\sin(2h|\xi|)| \left| \int_0^{|\xi|} \int_0^{|\xi|} dudv \sin(u+v) |u-v|^{2H-2} \right| \\ &= c |\sin(2h|\xi|)| \left| \int_0^{|\xi|} v^{2H-2} (\cos(v) - \cos(2|\xi| - v)) dv \right| \end{aligned}$$

and thus

$$\begin{aligned}
B &\leq c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 |\sin(2h|\xi|)| \left| \int_0^{|\xi|} v^{2H-2} (\cos(v) - \cos(2|\xi| - v)) dv \right| \\
&\leq c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 |\sin(2h|\xi|)| \sin(|\xi|) \left| \int_0^{|\xi|} \sin(v + |\xi|) v^{2H-2} dv \right| \\
&= c \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 |\sin(2h|\xi|)| \sin(|\xi|) \\
&\quad \times \left| \left(\sin(|\xi|) \int_0^{|\xi|} \cos v v^{2H-2} dv + \cos(|\xi|) \int_0^{|\xi|} \sin v v^{2H-2} dv \right) \right| \\
&= c \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 |\sin(2h|\xi|)| \sin(|\xi|) \\
&\quad \times \left| \left(\sin(|\xi|) \int_0^{|\xi|} \cos v v^{2H-2} dv + \cos(|\xi|) \int_0^{|\xi|} \sin v v^{2H-2} dv \right) \right| \\
&+ c \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(h|\xi|)^2 |\sin(2h|\xi|)| \sin(|\xi|) \\
&\quad \times \left| \left(\sin(|\xi|) \int_0^{|\xi|} \cos v v^{2H-2} dv + \cos(|\xi|) \int_0^{|\xi|} \sin v v^{2H-2} dv \right) \right|.
\end{aligned}$$

The part over the set $|\xi| \leq 1$ is bounded by ch^3 by simply majorizing $\sin(x)$ by x and $\left| \left(\sin(|\xi|) \int_0^{|\xi|} \cos v v^{2H-2} dv + \cos(|\xi|) \int_0^{|\xi|} \sin v v^{2H-2} dv \right) \right|$ by a constant. Concerning the part over the region $|\xi| \geq 1$ we bound again $\left| \left(\sin(|\xi|) \int_0^{|\xi|} \cos v v^{2H-2} dv + \cos(|\xi|) \int_0^{|\xi|} \sin v v^{2H-2} dv \right) \right|$ by a constant and we use the change of variables $\tilde{\xi} = h\xi$. This part will be bounded by

$$\begin{aligned}
&h^{2H+2-\beta} \int_{|\xi| \geq h} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(|\xi|)^2 |\sin(2|\xi|)| \\
&\leq h^{2H+2-\beta} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2H+2}} \sin(|\xi|)^2 |\sin(2|\xi|)| \\
&\leq ch^{2H+2-\beta}
\end{aligned}$$

since the last integral is convergent at infinity by bounded sinus by one and at zero by bounding $\sin(x)$ by x and using the assumption $\beta > 2H - 1$. Therefore

$$B \leq ch^{2H+2-\beta}. \quad (28)$$

We bound now the summand C below. In this summand the \mathcal{H} norm of the sinus function appear and this has been analyzed in [2]. We have, after the change of variables

$$\tilde{u} = \frac{u}{|\xi|}, \tilde{v} = \frac{v}{|\xi|},$$

$$\begin{aligned} C &= \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2}} \sin(h|\xi|)^4 \int_0^1 \int_0^1 \sin(|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv \\ &\geq \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2}} \sin(h|\xi|)^4 \int_0^1 \int_0^1 \sin(|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv. \end{aligned}$$

We will use Proposition 3.8 in [2] (more precisely, we will use the inequality two lines before the formula (34) in that paper with $k = 0$; we notice that the term $\sin(h|\xi|)^4$ does not appear in this proof but by analyzing the step of the proof we can see that this term can be added without problems). We will have that, for h small,

$$\begin{aligned} C &\geq \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta}} \sin(h|\xi|)^4 \frac{1}{|\xi|^2} \int_0^1 \int_0^1 \sin(|\xi|) \sin(v|\xi|) |u-v|^{2H-2} dudv \\ &\geq \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta}} \sin(h|\xi|)^4 \frac{1}{|\xi|^{2H+1}} \\ &= h^{2H+1-\beta} \int_{|\xi| \geq h} \frac{d\xi}{|\xi|^{d-\beta+2H+1}} \sin(h|\xi|)^4 \\ &\geq h^{2H+1-\beta} \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+1}} \sin(h|\xi|)^4 \\ &= ch^{2H+1-\beta}. \end{aligned} \tag{29}$$

Relations (28) and (29) imply (27). Now, from relation (27), for every $t_0 \leq s < t < T$ with s, t close enough

$$E_1(t-s) \geq c(t-s)^{2H+1-\beta} - c'(t-s)^{2H+2-\beta} \geq \frac{c}{2}(t-s)^{2H+1-\beta}$$

if $|t-s| \leq \frac{c}{2c'}$. To extend the above inequality to arbitrary values of $|t-s|$, we proceed as in [9], proof of Proposition 4.1. Notice that the function $g(t, s, x, y) := \mathbf{E} |u(t, x) - u(s, x)|^2$ is positive and continuous with respect to all its arguments and therefore it is bounded below on the set $\{(t, s, x, y) \in [t_0, T]^2 \times [-M, M]^{2d}; |t-s| \geq \varepsilon\}$ by a constant depending on $\varepsilon > 0$. Hence for $|t-s| \geq \frac{c}{2c'}$ it also holds that

$$E_1(t-s) \geq c_1 |t-s|^{2H+1-\beta}.$$

On the other side, from (21) and (26) and Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} E_{3,t}(h) &= \langle (g_{t+h,x} - g_{t,x}) \mathbf{1}_{[0,t]}, g_{t+h,x} \mathbf{1}_{[t,t+h]} \rangle_{\mathcal{HP}} \\ &\leq \| (g_{t+h,x} - g_{t,x}) \mathbf{1}_{[0,t]} \|_{\mathcal{HP}} \| g_{t+h,x} \mathbf{1}_{[t,t+h]} \|_{\mathcal{HP}} \\ &\leq ch^{\frac{2H+1-\beta}{2} + \frac{2H+2-\beta}{2}}. \end{aligned}$$

Consequently,

$$\mathbf{E} |u(t+h, x) - u(t, x)|^2 \geq Ch^{2H+1-\beta} - C'h^{\frac{2H+1-\beta}{2} + \frac{2H+2-\beta}{2}}$$

and this implies that for every $s, t \in [t_0, T]$ and $x \in [-M, M]^d$

$$\mathbf{E} |u(t, x) - u(s, x)|^2 \geq \frac{C}{2} |t - s|^{2H+1-\beta} \quad \text{if} \quad |t - s| \leq \left(\frac{C}{2C'} \right)^{\frac{1}{2}}.$$

Similarly as above, the previous inequality can be extended to arbitrary values of $s, t \in [t_0, T]$. ■

Proposition (1) implies the following Hölder property for the solution to (13).

Corollary 1 *Assume (19). Then for every $x \in \mathbb{R}^d$ the application*

$$t \rightarrow u(t, x)$$

is almost surely Hölder continuous of order $\delta \in \left(0, \frac{2H+1-\beta}{2}\right)$.

Proof: This is consequence of the relations (21) and (26) in the proof of Proposition 1 and of the fact that u is Gaussian. ■

Let us make some comments on the result in Proposition 1.

Remark 4

- *Following the proof of Theorem 5.1 in [8] we can show that the mapping $t \rightarrow u(t, x)$ is not Hölder continuous of order $\frac{2H+1-\beta}{2}$.*
- *When $H = \frac{1}{2}$ the solution of the wave equation with fractional noise in time has the same regularity in time as the solution of the wave equation with white noise in time (see [9], [8]).*

3.2 Space regularity

Let us discuss the behavior of the solution u to the equation (13) with respect to the spatial variable. We have

Proposition 2 *Assume (19), fix $M > 0$ and $t \in [t_0, T]$. Then there exist positive constants c_3, c_4 such that for any $x, y \in [-M, M]^d$*

$$c_3 |x - y|^{2H+1-\beta} \leq \mathbf{E} |u(t, x) - u(t, y)|^2 \leq c_4 |x - y|^{2H+1-\beta}.$$

Proof: Let $z \in \mathbb{R}^d$. We compute

$$\begin{aligned}
\mathbf{E}|u(t, x+z) - u(t, x)|^2 &= \|g_{t, x+z} - g_{t, x}\|_{\mathcal{HP}}^2 \\
&= \alpha_H \int_{\mathbb{R}^d} \int_0^t \int_0^t \mathcal{F}(g_{t, x+z} - g_{t, x})(u, \cdot)(\xi) \overline{\mathcal{F}(g_{t, x+z} - g_{t, x})(v, \cdot)(\xi)} |u - v|^{2H-2} du \, dv \, \mu(d\xi) \\
&= \alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du \, dv \int_{\mathbb{R}^d} |e^{-i\xi \cdot (x+z)} - e^{-i\xi \cdot x}|^2 \mathcal{F}G_1(u, \cdot)(\xi) \overline{\mathcal{F}G_1(v, \cdot)(\xi)} \mu(d\xi) \\
&= \alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du \, dv \int_{\mathbb{R}^d} |e^{-i\xi \cdot z} - 1|^2 \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|} \mu(d\xi) \\
&=: E_{1,x}(z) + E_{2,x}(z),
\end{aligned}$$

where $E_{1,x}(z)$ and $E_{2,x}(z)$ are the integrals over the regions $|\xi| < 1$ and $|\xi| \geq 1$ respectively. For the first expression is easy to see that, using the inequality $|1 - e^{-i\xi z}|^2 \leq |\xi|^2 |z|^2$, we get the bound

$$E_{1,x}(z) \leq C|z|^2 \int_{|\xi| \leq 1} \mu(d\xi).$$

Developing the second expression we get

$$\begin{aligned}
E_{2,x}(z) &= \alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du \, dv \int_{|\xi| \geq 1} |e^{-i\xi \cdot z} - 1|^2 \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|} \mu(d\xi) \\
&= 2\alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du \, dv \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta}} (1 - \cos(z \cdot \xi)) \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|},
\end{aligned}$$

where $z \cdot \xi$ means the scalar product in \mathbb{R}^d . Again from Proposition 3.7 in [2] we have that

$$N_t(\xi) \leq c_{t,H} \left(\frac{1}{1 + |\xi|^2} \right)^{H+1/2}$$

for any $t > 0$, $|\xi| \geq 1$, where $N_t(\xi)$ is given by (25). Hence, denoting by $e = \frac{z}{|z|}$

$$\begin{aligned}
E_{2,x}(z) &\leq C \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta}} (1 - \cos(z \cdot \xi)) \left(\frac{1}{1 + |\xi|^2} \right)^{H+\frac{1}{2}} \\
&= C z^{2H+1-\beta} \int_{\mathbb{R}^d} \frac{dw}{|w|^{d-\beta}} (1 - \cos(w \cdot e)) \left(\frac{1}{|w|^2 + |z|^2} \right)^{H+\frac{1}{2}} \\
&\leq C |z|^{2H+1-\beta},
\end{aligned}$$

where we used the change of variables $w = \xi|z|$. This proves the upper bound.

Let us prove the sharpness of this bound (i.e. the lower bound). We can assume, without losing the generality, that $t = 1$. We note that

$$\begin{aligned}
&\mathbf{E}|u(1, x+z) - u(1, x)|^2 \geq F_2(z) \\
&:= 2\alpha_H \int_0^1 \int_0^1 du \, dv |u - v|^{2H-2} \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta}} (1 - \cos(\xi \cdot z)) \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|}.
\end{aligned}$$

Condition (17) implies that

$$\int_{|\xi| \geq 1} \frac{\mu(d\xi)}{|\xi|^3} \leq \int_{|\xi| \geq 1} \frac{\mu(d\xi)}{|\xi|^{2H+1}} < \infty.$$

We apply Proposition 3.8 in [2] (more precisely, the inequality two lines before (34) in [2] with $k = 0$) and we get (note that the result in [2] is stated without the factor $(1 - \cos(\xi \cdot z))$ but by analyzing the steps of the proof we can see that this factor may be added without problems)

$$F_2(z) \geq C \int_{|\xi| \geq 1} \frac{d\xi}{|\xi|^{d-\beta+2H+1}} (1 - \cos(\xi \cdot z)).$$

and by the change of variables $\xi|z| = w$ in the integral $d\xi$

$$F_2(z) \geq C z^{2H+1-\beta} \int_{|w| \geq |z|} \frac{dw}{|w|^{d-\beta+2H+1}} (1 - \cos(w \cdot e)).$$

As in the proof of Theorem 5.1 in [8], we obtain that the integral $\int_{|w| \geq |z|} \frac{dw}{|w|^{d-\beta+2H+1}} (1 - \cos(w \cdot e))$ is bounded below by a constant. (Notice that $\beta > 2H - 1$, implies that the first integral above is convergent when z is zero, because $1 - \cos(x) \approx x^2$ around zero). Thus, it is immediate that

$$\mathbf{E} |u(1, x + z) - u(1, x)|^2 \geq C z^{2H+1-\beta} - C' z^{2H+2-\beta}$$

and this implies

$$\mathbf{E} |u(1, x + z) - u(1, x)|^2 \geq \frac{1}{2} C z^{2H+1-\beta}$$

for $|z| \leq \frac{C}{2C'}$. It is a routine argument to extend the above inequality to arbitrary values of $|z|$ (see e.g. [9], page 22, see also the proof of Proposition 1 before). ■

We have the following result concerning the Hölder continuity in space. We mention that it is a little bit more than an extension of Proposition 2.

Proposition 3 *Assume (19). Then for any $t \in [t_0, T]$ the application*

$$x \rightarrow u(t, x)$$

is almost surely Hölder continuous of order $\delta \in \left(0, \left(\frac{2H+1-\beta}{2}\right) \wedge 1\right)$.

Proof: We claim that

$$\mathbf{E} |u(t, x) - u(t, y)|^2 \leq c |x - y|^{(2H+1-\beta) \wedge 2} \quad (30)$$

for $|x - y|$ small enough. From Proposition 2, (30) is true when $\beta > 2H - 1$. When $\beta \in (0, 2H - 1)$ then it suffices to regard the part of the quantity $\mathbf{E} |u(t, x + z) - u(t, x)|^2$ over the region $|\xi| \leq 1$ (the part over the region $|\xi| > 1$ is, as in the proof of Proposition 2, bounded by $cz^{2H+1-\beta}$ so by cz^2 for z small). It is immediate to see that, using the inequality $|1 - e^{-i\xi z}|^2 \leq |\xi|^2 |z|^2$ the considered part is less than $C|z|^2 \int_{|\xi| \leq 1} \mu(d\xi)$. This concludes the proof of (30).

The conclusion is a consequence of Proposition 2, the Gaussianity of u and the Kolmogorov continuity theorem. ■

Remark 5

- When $H = \frac{1}{2}$ the solution of the wave equation with fractional noise in time has the same regularity in space as the solution of the wave equation with white noise in time (see [9], [8]).
- We distinguish in Proposition 3 two cases: if $\beta \in (0, 2H - 1)$ then the solution to (13) has spatial Hölder continuity of order 1 (so, it is Lipschitz continuous in the space variable) while if $\beta \in (2H - 1, d \wedge (2H + 1))$ the Hölder exponent is $\delta \in (0, \frac{2H+1-\beta}{2}) < 1$.
- There is another way to see why the cases $\beta \in (0, 2H - 1]$ and $\beta \in (2H - 1, d \wedge (2H + 1))$ need to be separated. Denote by

$$\begin{aligned} g_t(z) &:= \mathbf{E} |u(t, x + z) - u(t, x)|^2 \\ &= 2\alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du dv \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta}} (1 - \cos(z \cdot \xi)) \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|} \end{aligned}$$

and let us study the behavior of g_t around $z = 0$. Let us also assume that $d = 1$. Notice first that $g_t(0) = 0$ and

$$g'_t(z) = 2\alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du dv \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta-1}} \sin(z \cdot \xi) \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|}$$

and thus $g'_t(0) = 0$ provided that $\beta < 2H$. Moreover

$$g''_t(z) = 2\alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du dv \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta-2}} \cos(z \cdot \xi) \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|}$$

and

$$\begin{aligned} g''_t(0) &= 2\alpha_H \int_0^t \int_0^t |u - v|^{2H-2} du dv \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta-2}} \frac{\sin(u|\xi|)}{|\xi|} \cdot \frac{\sin(v|\xi|)}{|\xi|} \\ &\leq C_t 2\alpha_H \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta-2}} \left(\frac{1}{1 + |\xi|^2} \right)^{H+\frac{1}{2}} \end{aligned}$$

which is a finite constant for $\beta < 2H - 1$. Therefore $g_t(z)$ behaves as Cz^2 for z close to zero.

3.3 Joint regularity

Let us denote by Δ the following metric on $[0, T] \times \mathbb{R}^d$

$$\Delta((t, x); (s, y)) = |t - s|^{2H+1-\beta} + |x - y|^{2H+1-\beta}. \quad (31)$$

From Propositions 1 and 2, we obtain the following result:

Theorem 2 *Fix $M > 0$ and assume (19). For every $t, s \in [t_0, T]$ and $x, y \in [-M, M]^d$ there exist positive constants C_1, C_2 such that*

$$C_1 \Delta((t, x); (s, y)) \leq \mathbf{E} |u(t, x) - u(s, y)|^2 \leq C_2 \Delta((t, x); (s, y)).$$

Proof: The upper bound can be easily obtained by using the upper bound in Propositions 1 and 2 since

$$\begin{aligned} \mathbf{E} |u(t, x) - u(s, y)|^2 &\leq 2\mathbf{E} |u(t, x) - u(s, x)|^2 + 2\mathbf{E} |u(s, x) - u(s, y)|^2 \\ &\leq C_2 \left(|t - s|^{2H+1-\beta} + |x - y|^{2H+1-\beta} \right). \end{aligned}$$

Concerning the lower bound, it suffices follow the lines of the proof of Lemma 2.1 in [15] (see also Steps 3 and 4 in the proof of Proposition 4.1 in [9]). We will briefly explain the main lines of the proof. The demonstration needs to be divided upon three cases: $|t - s|^{2H+1-\beta} \leq \frac{c_3}{4c_2} |x - y|^{2H+1-\beta}$, $|t - s|^{2H+1-\beta} \geq \frac{4c_4}{c_1} |x - y|^{2H+1-\beta}$ and $\frac{4c_4}{c_1} |x - y|^{2H+1-\beta} \geq |t - s|^{2H+1-\beta} \geq \frac{c_3}{4c_2} |x - y|^{2H+1-\beta}$ with the constants c_1, c_2, c_3, c_4 appearing in the statements of Propositions 1 and 2. The first case can be handled as follows

$$\begin{aligned} \mathbf{E} |u(t, x) - u(s, y)|^2 &\geq \frac{1}{2} \mathbf{E} |u(t, x) - u(t, y)|^2 - \mathbf{E} |u(t, y) - u(s, y)|^2 \\ &\geq \frac{1}{2} c_3 |x - y|^{2H+1-\beta} - c_2 |t - s|^{2H+1-\beta} \\ &\geq \frac{1}{2} c_3 |x - y|^{2H+1-\beta} - \frac{1}{4} c_3 |x - y|^{2H+1-\beta} \\ &= \frac{1}{4} c_3 |x - y|^{2H+1-\beta} \\ &\geq \frac{c_3}{8} |x - y|^{2H+1-\beta} + \frac{c_3}{8} \frac{4c_2}{c_3} |t - s|^{2H+1-\beta} \\ &\geq C_1 \Delta((t, x); (s, y)). \end{aligned}$$

The other cases follows similarly from Lemma 3.1 in [15], by replacing their exponents with our exponents. ■

Remark 6 *The result of Theorem 2 can be stated also in the following form: Fix $M > 0$ and assume (19). For every $t, s \in [t_0, T]$ and $x, y \in [-M, M]^d$ with (t, x) close enough to (s, y) , there exist positive constants C_1, C_2 such that*

$$C_1 (|t - s| + |x - y|)^{2H+1-\beta} \leq \mathbf{E} |u(t, x) - u(s, y)|^2 \leq C_2 (|t - s| + |x - y|)^{2H+1-\beta}.$$

4 Hitting times

Let us discuss the upper and lower bounds for the hitting probabilities of the solution u to equation (13). These bounds will be given in terms of the Newtonian capacity and the Hausdorff measure of the hit set (see Section 2 for the definition). Let us recall the notation: if $V = (V(x), x \in \mathbb{R}^m)$ is a \mathbb{R}^k valued stochastic process then $V(S)$ denote the range of the Borel set S under the random mapping $x \rightarrow V(x)$.

Our result is based on the following criteria for the hitting probabilities proven in [3], Theorem 2.1.

Theorem 3 *Let $X = X(t), t \in \mathbb{R}^N$ be a \mathbb{R}^k -valued centered Gaussian process and fix $I \subset \mathbb{R}^N$. Assume that there exist positive constants a_1, a_2, a_3, a_4 such that*

- i. For every $t \in I$, $\mathbf{E} [X(t)^2] \geq a_1 > 0$.*
- ii. There exists $\alpha_1, \dots, \alpha_N \in (0, 1)$ such that for every $t = (t_1, \dots, t_N), s = (s_1, \dots, s_N) \in I$ it holds that*

$$a_2 \sum_{j=1}^N |t_j - s_j|^{2\alpha_j} \leq \mathbf{E} |X(t) - X(s)|^2 \leq a_3 \sum_{j=1}^N |t_j - s_j|^{2\alpha_j}.$$

- iii. For every $t = (t_1, \dots, t_N), s = (s_1, \dots, s_N) \in I$*

$$\text{Var}(X(t)|X(s)) \geq a_4 \sum_{j=1}^N |t_j - s_j|^{2\alpha_j}.$$

Then there exist positive constants a_5, a_6 such that for every Borel set A in \mathbb{R}^k

$$a_5 \text{Cap}_{k-Q}(A) \leq P(X(I) \cap A \neq \emptyset) \leq a_6 \mathcal{H}_{k-Q}(A)$$

where $Q = \sum_{j=1}^N \frac{1}{\alpha_j}$.

Next, we will show that the solution to (13) satisfies the assumptions of the previous result. This will be done via several lemmas.

Lemma 3 *Assume (19) and let u be the solution to (13). Then for every $t \in [t_0, T]$ and $x \in \mathbb{R}^d$*

$$\mathbf{E} u(t, x)^2 \geq C.$$

Proof: Let $\sigma_{t,x}^2$ be the variance of $u(t, x)$. We need to give a lower bound for this variance. Assume for simplicity $t = 1$. Then

$$\begin{aligned}
\sigma_{1,x}^2 &= \mathbf{E} |u(1, x)|^2 \\
&= \alpha_H \int_0^1 \int_0^1 dudv |u - v|^{2H-2} \int_{\mathbb{R}^d} \frac{d\xi}{|\xi|^{d-\beta+2}} \sin(u|\xi|) \sin(v|\xi|) \\
&\geq \alpha_H \int_0^1 \int_0^1 dudv |u - v|^{2H-2} \int_{|\xi| \leq 1} \frac{d\xi}{|\xi|^{d-\beta+2}} \sin(u|\xi|) \sin(v|\xi|) \\
&\geq \alpha_H \sin^2 1 \int_0^1 \int_0^1 dudv |u - v|^{2H-2} uv = C > 0
\end{aligned}$$

where we used the bound $\sin x \geq x \sin 1$ for every $x \in [0, 1]$. The general case $t \in [t_0, T]$ follows in the same way by doing the change of variables $\tilde{u} = \frac{u}{t}, \tilde{v} = \frac{v}{t}$ and then working on the domain $D = \{\xi \in \mathbb{R}^d, |\xi| \leq \frac{1}{ut}\}$. \blacksquare

Now, we bound the conditional variance (condition iii. in Theorem 3).

Lemma 4 Assume (19) and fix $t_0, M > 0$. Then for every $s, t \in [t_0, T]$ and $x, y \in [-M, M]^d$

$$\text{Var}(u(t, x) | u(s, y)) \geq C \Delta((t, x); (s, y))$$

where Δ is the metric given by (31).

Proof: We will use the following formula: if (U, V) is a centered Gaussian vector, then

$$\text{Var}(U, V) = \frac{(\rho_{U,V}^2 - (\sigma_U - \sigma_V)^2)((\sigma_U + \sigma_V)^2 - \rho_{U,V}^2)}{4\sigma_V^2} \quad (32)$$

where $\rho_{U,V}^2 = \mathbf{E}(U - V)^2, \sigma_U^2 = \mathbf{E}U^2, \sigma_V^2 = \mathbf{E}V^2$. Denote by

$$\rho_{t,x,s,y}^2 = \mathbf{E} |u(t, x) - u(s, y)|^2, \quad \sigma^2(t, x) = \mathbf{E}u(t, x)^2, \quad \sigma_{s,y}^2 = \mathbf{E}u(s, y)^2.$$

It suffices to show that

$$(\rho_{t,x,s,y}^2 - (\sigma_{t,x} - \sigma_{s,y})^2)((\sigma_{t,x} + \sigma_{s,y})^2 - \rho_{t,x,s,y}^2) \geq c \Delta((t, x); (s, y))$$

for every $s, t \in [t_0, T]$ and $x, y \in [-M, M]^d$. By Theorem 2 the second factor in the left-hand side above is bounded below by a constant. So it remains to check that

$$(\rho_{t,x,s,y}^2 - (\sigma_{t,x} - \sigma_{s,y})^2) \geq c \Delta((t, x); (s, y))$$

but this has been done in [15], proof of Proposition 3.2. (see also [6], proof of Lemma 4.3). \blacksquare

Remark 7 Using the previous result we can give a bound on the joint density $p_{t,x,s,y}$ of the vector $(u(t,x), u(s,y))$. Actually, one can show that for every $t \in [t_0, T]$ and $x, y \in [-M, M]^d$ we have the inequality

$$p_{t,s,x,y}(z_1, z_2) \leq C_1 \Delta((t, x); (s, y))^{-\frac{1}{2}} \exp\left(-\frac{C_2 |z_1 - z_2|^2}{\Delta((t, x); (s, y))}\right)$$

for every $z_1, z_2 \in [-N, N]^k$, where Δ is the metric defined by (31). It suffices to follow the lines of Proposition 3.2 in [15].

We can state now the main result of this section.

Theorem 4 Let us consider I, J non-trivial compact sets in $[t_0, T]$ and $[-M, M]^d$ respectively. Fix $N > 0$ and let u be the solution to the system (1). Then for every Borel set A contained in $[-N, N]^k$ it holds that

$$C^{-1} \text{Cap}_{k-\gamma} \leq P(u(I \times J) \cap A \neq \emptyset) \leq C \mathcal{H}_{k-\gamma}(A)$$

with

$$\gamma = k - \frac{2(d+1)}{2H+1-\beta}.$$

Proof: The proof is a consequence of Theorem 3 and of the preceding two lemmas. ■

Remark 8

- Of course, for H close to $\frac{1}{2}$, our result recovers the findings in [9].
- it is also possible to give some results concerning the probability that, for fixed t, x , the sets $u(\{t\} \times J)$ and $u(I \times \{x\})$ (as before I, J non-trivial compact sets in $[t_0, T]$ and in $[-M, M]^d$ respectively) to hit a given Borel set A contained in $[-N, N]^k$. Actually, by routine arguments we will have

$$C^{-1} \text{Cap}_{k-\frac{2d}{2H+1-\beta}}(A) \leq P(u(\{t\} \times J) \cap A \neq \emptyset) \leq C \mathcal{H}_{k-\frac{2d}{2H+1-\beta}}(A)$$

and

$$C^{-1} \text{Cap}_{k-\frac{2}{2H+1-\beta}}(A) \leq P(u(I \times \{x\}) \cap A \neq \emptyset) \leq C \mathcal{H}_{k-\frac{2}{2H+1-\beta}}(A).$$

References

- [1] Balan R.M. and Tudor C. A. (2008): *The stochastic heat equation with fractional-colored noise: existence of the solution*. Latin Amer. J. Probab. Math. Stat. **4**, 57-87. Erratum in Latin Amer. J. Probab. Math. Stat. (2009) **6**, 343-347.

- [2] Balan R.M. and Tudor C. A. (2010): *The stochastic wave equation with fractional noise: A random field approach*. Stoch. Proc. Appl. **120**, 2468-2494.
- [3] Biermé H., Lacaux C. and Xiao Y. (2009): *Hitting probabilities and the Hausdorff dimension of the inverse images of anisotropic Gaussian random fields*. Bull. Lond. Math. Soc. **41**, 253-273.
- [4] Dalang R. C. (1999): *Extending the martingale measure stochastic integral with applications to spatially homogeneous SPDE's*. Electr. J. Probab. **4**, 1-29. pp. Erratum in Electr. J. Probab. **6** (2001), 5 pp.
- [5] Dalang R. C. and Nualart E. (2004): *Potential theory for hyperbolic SPDEs*. Ann. Probab. **32**, 2099-2148.
- [6] Dalang R. C., Khosnevisan D. and Nualart E. (2007): *Hitting probabilities for systems of non-linear heat equations with additive noise*. ALEA **3**, 231-271.
- [7] Dalang R. C., Khosnevisan D. and Nualart E. (2009): *Hitting probabilities for systems of non-linear stochastic heat equations with multiplicative noise*. Probab. Theory and Related Fields **144**, 371-427.
- [8] Dalang R. and Sanz-Solé M. (2009): *Hölder-Sobolev regularity of the solution to the stochastic wave equation in dimension three*. Memoirs of the American Mathematical Society **199**(931), 1-70.
- [9] Dalang R. and Sanz-Solé M. (2010): *Criteria for hitting probabilities with applications to systems of stochastic wave equations*. Bernoulli **16**(4), 1343-1368.
- [10] Gubinelli M., Lejay A. and Tindel S. (2006): *Young integrals and SPDE's*. Potential Anal. **25**, 307-326.
- [11] Maslovski B. and Nualart D. (2003): *Evolution equations driven by a fractional Brownian motion*. J. Funct. Anal. **202**, 277-305.
- [12] Mémin J., Mishura Y. and Valkeila E. (2001): *Inequalities for the moments of Wiener integrals with respect to fractional Brownian motions*. Stat. Probab. Letters **51**, 197-206.
- [13] Mueller C. and Tribe C. (2002): *Hitting probabilities of the random string*. Electronic Journal of probability **7**, 1-29.
- [14] Nualart D. and Vuillermont P.-A. (2006): *Variational solutions for partial differential equations driven by fractional a noise*. J. Funct. Anal. **232**, 390-454.
- [15] Nualart E. and Viens F. (2009): *The fractional stochastic heat equation on the circle: time regularity and potential theory*. Stochastic Processes and their Applications **119**, 1505-1540.

- [16] Quer-Sardanyons L. and Tindel S. (2007): *The 1-d stochastic wave equation driven by a fractional Brownian sheet*. Stoch. Proc. Appl. **117**, 1448-1472.
- [17] Sanz-Solé M. and Vuillermot P.-A. (2009). *Mild solutions for a class of fractional SPDE's and their sample paths*. J. Evolution Equations **9**, 235-265.
- [18] Tindel S., Tudor C. A. and Viens F. (2003). *Stochastic evolution equations with fractional Brownian motion*. Probab. Theor. Rel. Fields **127**, 186-204.
- [19] Treves F. (1975). *Basic Linear Partial Differential Equations*. Academic Press, New York.

Centro de Investigación en Ingeniería Matemática (CI²MA)


PRE-PUBLICACIONES 2011 - 2012

- 2011-30 RAIMUND BÜRGER, PEP MULET, LUIS M. VILLADA: *Spectral WENO schemes with adaptive mesh refinement for models of polydisperse sedimentation*
- 2012-01 GABRIEL N. GATICA, GEORGE C. HSIAO, SALIM MEDDAHI, FRANCISCO J. SAYAS: *On the dual-mixed formulation for an exterior Stokes problem*
- 2012-02 GABRIEL N. GATICA, LUIS F. GATICA, ANTONIO MARQUEZ: *Analysis of a pseudo-stress-based mixed finite element method for the Brinkman model of porous media flow*
- 2012-03 RAIMUND BÜRGER, STEFAN DIEHL, SEBASTIAN FARÅS, INGMAR NOPENS, ELENA TORFS: *A consistent modelling methodology for secondary settling tanks: A reliable numerical method*
- 2012-04 RAIMUND BÜRGER, STEFAN DIEHL, SEBASTIAN FARÅS, INGMAR NOPENS: *On reliable and unreliable numerical methods for the simulation of secondary settling tanks in wastewater treatment*
- 2012-05 MARGARETH ALVES, JAIME MUÑOZ-RIVERA, MAURICIO SEPÚLVEDA, OCTAVIO VERA, MARÍA ZEGARRA-GARAY: *The asymptotic behaviour of the linear transmission problem in viscoelasticity*
- 2012-06 SALIM MEDDAHI, DAVID MORA, RODOLFO RODRÍGUEZ: *Finite element spectral analysis for the mixed formulation of the elasticity equations*
- 2012-07 ANA I. GARRALDA GUILLEM, GABRIEL N. GATICA, ANTONIO MARQUEZ, MANUEL RUIZ GALAN: *A posteriori error analysis of twofold saddle point variational formulations for nonlinear boundary value problems*
- 2012-08 LEONARDO E. FIGUEROA, ENDRE SÜLI: *Greedy approximation of high-dimensional Ornstein–Uhlenbeck operators*
- 2012-09 CLAUDIO PADRA, RODOLFO RODRÍGUEZ, MARIO SCHEBLE: *An hp finite element method to solve a fluid-solid vibration problem*
- 2012-10 JULIO ARACENA, LAURENCE CALZONE, JEAN - PAUL COMET, JACQUES DEMON-GEOT, MARCELLE KAUFMAN, AURÉLIEN NALDI, ADRIEN RICHARD, EL HOUSSINE SNOUSSI, DENIS THIEFFRY: *On circuit functionality in Boolean networks*
- 2012-11 JORGE CLARKE, CIPRIAN A. TUDOR: *Hitting times for the stochastic wave equation with fractional-colored noise*

Para obtener copias de las Pre-Publicaciones, escribir o llamar a: DIRECTOR, CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE, TEL.: 41-2661324, o bien, visitar la página web del centro: <http://www.ci2ma.udec.cl>



**CENTRO DE INVESTIGACIÓN EN
INGENIERÍA MATEMÁTICA (CI²MA)
Universidad de Concepción**



Casilla 160-C, Concepción, Chile
Tel.: 56-41-2661324/2661554/2661316
<http://www.ci2ma.udec.cl>

