

UNIVERSIDAD DE CONCEPCIÓN



CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA (CI²MA)



**Advection-diffusion-reaction in poroelastic media. Part I:
Well-posedness and discrete analysis**

LUIS MIGUEL DE OLIVEIRA VILACA, BRYAN GOMEZ-VARGAS,
SARVESH KUMAR, RICARDO RUIZ-BAIER,
NITESH VERMA

PREPRINT 2019-35

SERIE DE PRE-PUBLICACIONES

Advection-diffusion-reaction in poroelastic media. Part I: Well-posedness and discrete analysis

Luis Miguel De Oliveira Vilaca^d, Bryan Gómez-Vargas^{b,c}, Sarvesh Kumar^a, Ricardo Ruiz-Baier^{e,*},
Nitesh Verma^a

^aDepartment of Mathematics, Indian Institute of Space Science and Technology, Trivandrum 695 547, India.

^bCI²MA and Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C, Concepción, Chile.

^cSección de Matemática, Sede de Occidente, Universidad de Costa Rica, San Ramón, Alajuela, Costa Rica.

^dLaboratory of Artificial & Natural Evolution (LANE), Department of Genetics and Evolution, University of Geneva, 4 Boulevard d'Yvoy,
1205 Geneva, Switzerland

^eMathematical Institute, University of Oxford, A. Wiles Building, Woodstock Road, Oxford OX2 6GG, UK.

Abstract

We analyse a PDE system modelling poromechanical processes (formulated in mixed form using the solid deformation, fluid pressure, and total pressure) interacting with diffusing and reacting solutes in the medium. We investigate the well-posedness of the nonlinear set of equations using fixed-point theory, Fredholm's alternative, a priori estimates, and compactness arguments. We also propose a mixed finite element method and rigorously demonstrate the stability of the scheme.

Keywords: Biot equations, reaction-diffusion, mixed finite elements, well-posedness and stability
2000 MSC: 65M60, 74F10, 35K57, 74L15

1. Introduction and problem statement

1.1. Scope of the paper

We aim at studying the spreading properties of a system of interacting species when the underlying medium is of a porous nature and it undergoes elastic deformations. The model we propose has the potential to deliver quantitative insight on the two-way coupling between the transport of solutes and poromechanical effects in the context of microscopic-macroscopic mechanobiology. Real biological tissues are conformed by living cells, and volume changes due to cell birth and death onset velocity fields and local deformation, eventually driving domain growth [22]. Interconnectivity of the porous microstructure is in this case sufficient to accommodate fluid flowing locally. The described problem can be encountered in numerous applications not only related to cell biomechanics, and these are explored in [29].

*Author for correspondence. Email: ruizbaier@maths.ox.ac.uk. Phone: +44 1865 615168.

Email addresses: LuisMiguel.DeOliveiraVilaca@unige.ch (Luis Miguel De Oliveira Vilaca),
bgomez@ci2ma.udec.cl (Bryan Gómez-Vargas), sarvesh@iist.ac.in (Sarvesh Kumar), ruizbaier@maths.ox.ac.uk
(Ricardo Ruiz-Baier), nitesh@iist.ac.in (Nitesh Verma)

Funding: This work has been partially supported by CONICYT through the Becas-Chile Programme for foreign students, by the London Mathematical Society through Scheme 5, Grant 51703, and by the CONICYT/PIA project AFB170001.

From the viewpoint of solvability analysis of partial differential equations and/or the theoretical aspects of finite element discretisations, the relevant literature contains a few works specifically targeting the coupling of diffusion in deformable porous media. We mention for instance the classical works of Showalter [27] and Showalter and Momken [28] which employ the theory of degenerate equations in Hilbert spaces, or the study of Hadamard well-posedness of parabolic-elliptic systems governing chemo-poroelasticity with thermal effects [21]. More recently, [20] introduces mixed finite element schemes and stability analysis for a system of multiple-network poroelasticity, that resembles the model problem we are interested in. Also, in [9] a six-field system including temperature dynamics has been rigorously analysed using linearisation tools, the Banach fixed-point theory and weak compactness, and piecewise continuation in time. As in [20], here we also employ the three-field formulation for the Biot consolidation equations introduced in [23] (see also [19]). However in the model we adopt here, we consider a two-way active transport: the poromechanical deformations affect the transport of the chemical species through advection and also by means of a volume-dependent modification of the reaction terms; and the solutes' concentration generate an active stress resulting in a distributed load depending linearly on the concentration gradients.

The coupled system is set up in mixed-primal structure, where the equations of poroelasticity have a mixed form using displacement, pressure, and a rescaled total pressure, and the advection-diffusion-reaction system is also set in primal form, solving for the concentrations. Then, we focus on the semidiscrete in-time formulation, rewriting the resulting scheme equivalently as a fixed-point equation [3, 5, 11], and then, Schauder fixed point theorem [3, 11], combined with Fredholm's alternative [6, 12, 23] and quasi-linear equations theory [5, 18], are applied to establish the solvability of the introduced formulation. Consequently, the well-known MINI-elements family and continuous piecewise polynomials are proposed to approximate the three-field formulation, whereas Lagrange elements are introduced to approximate the concentrations. Thus, making use of the discrete inf-sup condition together with classical inequalities, we obtain the corresponding stability result for our approximation. The advantage of using this approach is that the stability results are independent of the Lamé constants of the solid, and this is particularly important to prevent volumetric locking. We further stress that the main difficulties in the present analysis (which are not present in the literature cited above) are related to the advective coupling appearing in the advection-reaction-diffusion system. In contrast with e.g. [10, 9], the advecting velocity in our case is that of the solid (instead of the Darcy velocity), which is not a primary variable in our formulation. This implies that an extra $1/(\Delta t)$ appears from the backward Euler time discretisation of the solid velocity, complicating the analysis of the semidiscrete and fully discrete problems.

The remainder of this work is structured as follows. The governing equations as well as the main assumptions on the model coefficients will be stated in what is left of this Section. Then, in Section 2 we derive a weak formulation and include preliminary properties of the mathematical structure of the problem. Well-posedness of the coupled problem is then analysed also in Section 2, focusing in the semidiscrete case. We proceed in Section 3 with the introduction of a locking-free finite element scheme for the discretisation of the model equations, based on a stabilised formulation from [23] for the consolidation system, and a conforming method for the advection-diffusion-reaction subsystem. The convergence of the fully-discrete method is shown by means of a simple test presented in Section 4. We close with a discussion on model extensions in Section 5.

Let us point out that In a companion paper [29] we are addressing in more detail the modelling formalisms, we perform a linear stability analysis to identify suitable ranges for the key coupling parameters, and we give a full set of numerical tests in 2D and 3D.

1.2. Coupling poroelasticity and advection-diffusion-reaction

Let us consider a piece of soft material as a porous medium composed by a mixture of incompressible grains and interstitial fluid, whose description can be placed in the context of the classical Biot problem. As in [23, 19], we introduce an auxiliary unknown ψ representing the volumetric part of the total stress. In the absence of gravitational forces, and for a given body load $\mathbf{b}(t) : \Omega \rightarrow \mathbb{R}^d$ and a mass source $\ell(t) : \Omega \rightarrow \mathbb{R}$, one seeks for each time $t \in (0, t_{\text{final}}]$, the displacements of the porous skeleton, $\mathbf{u}^s(t) : \Omega \rightarrow \mathbb{R}^d$, and the pore pressure of the fluid, $p^f(t) : \Omega \rightarrow \mathbb{R}$, such that

$$\left(c_0 + \frac{\alpha^2}{\lambda}\right) \partial_t p^f - \frac{\alpha}{\lambda} \partial_t \psi - \frac{1}{\eta} \operatorname{div}(\kappa \nabla p^f) = \ell \quad \text{in } \Omega \times (0, t_{\text{final}}], \quad (1.1)$$

$$\boldsymbol{\sigma} = 2\mu \boldsymbol{\varepsilon}(\mathbf{u}^s) - \psi \mathbf{I}, \quad \text{in } \Omega \times (0, t_{\text{final}}], \quad (1.2)$$

$$\psi = \alpha p^f - \lambda \operatorname{div} \mathbf{u}^s, \quad \text{in } \Omega \times (0, t_{\text{final}}], \quad (1.3)$$

$$-\operatorname{div} \boldsymbol{\sigma} = \rho \mathbf{b} \quad \text{in } \Omega \times (0, t_{\text{final}}]. \quad (1.4)$$

Here $\kappa(\mathbf{x})$ is the hydraulic conductivity of the porous medium (possibly anisotropic), ρ is the density of the solid material, η is the constant viscosity of the interstitial fluid, c_0 is the constrained specific storage coefficient, α is the Biot-Willis consolidation parameter, and μ, λ are the shear and dilation moduli associated with the constitutive law of the solid structure.

We also consider the propagation of a generic species with concentration w_1 , reacting with an additional species with concentration w_2 . The problem can be written as follows

$$\partial_t w_1 + \partial_t \mathbf{u}^s \cdot \nabla w_1 - \operatorname{div}\{D_1(\mathbf{x}) \nabla w_1\} = f(w_1, w_2, \mathbf{u}^s) \quad \text{in } \Omega \times (0, t_{\text{final}}], \quad (1.5)$$

$$\partial_t w_2 + \partial_t \mathbf{u}^s \cdot \nabla w_2 - \operatorname{div}\{D_2(\mathbf{x}) \nabla w_2\} = g(w_1, w_2, \mathbf{u}^s) \quad \text{in } \Omega \times (0, t_{\text{final}}], \quad (1.6)$$

where D_1, D_2 are positive definite diffusion matrices (however we do not consider here cross-diffusion effects as in e.g. [5, 25]). In the well-posedness analysis the reaction kinetics are generic. Nevertheless, for sake of fixing ideas and in order to specify the coupling effects also through a stability analysis that will be conducted in [29], they will be chosen as a modification to the classical model from [26]

$$\begin{aligned} f(w_1, w_2, \mathbf{u}^s) &= \beta_1(\beta_2 - w_1 + w_1^2 w_2) + \gamma w_1 \partial_t \operatorname{div} \mathbf{u}^s, \\ g(w_1, w_2, \mathbf{u}^s) &= \beta_1(\beta_3 - w_1^2 w_2) + \gamma w_2 \partial_t \operatorname{div} \mathbf{u}^s, \end{aligned}$$

where $\beta_1, \beta_2, \beta_3, \gamma$ are positive model constants. Note that the mechano-chemical feedback (the process where mechanical deformation modifies the reaction-diffusion effects) is here assumed only through advection and an additional reaction term depending on local dilation. The latter term is here modulated by $\gamma > 0$, thus representing a source for both species if the solid volume increases, otherwise the additional contribution is a sink for both chemicals [22].

The poromechanical deformations are also actively influenced by microscopic tension generation. A very simple description is given in terms of active stresses: we assume that the total Cauchy stress contains a passive and an active component, where the passive part is as in (1.2) and

$$\boldsymbol{\sigma}_{\text{total}} = \boldsymbol{\sigma} + \boldsymbol{\sigma}_{\text{act}}, \quad (1.7)$$

where the active stress operates primarily on a given, constant direction $\mathbf{k}$, and its intensity depends on a scalar field $r = r(w_1, w_2)$ and on a positive constant τ , to be specified later on (see e.g. [16])

$$\boldsymbol{\sigma}_{\text{act}} = -\tau r \mathbf{k} \otimes \mathbf{k}. \quad (1.8)$$

In summary, the coupled system reads

$$-\operatorname{div}(2\mu \boldsymbol{\varepsilon}(\mathbf{u}^s) - \psi \mathbf{I} + \boldsymbol{\sigma}_{\text{act}}) = \rho \mathbf{b} \quad \text{in } \Omega \times (0, t_{\text{final}}],$$

$$\begin{aligned}
\left(c_0 + \frac{\alpha^2}{\lambda}\right) \partial_t p^f - \frac{\alpha}{\lambda} \partial_t \psi - \frac{1}{\eta} \operatorname{div}(\kappa \nabla p^f) &= \ell & \text{in } \Omega \times (0, t_{\text{final}}], \\
\psi - \alpha p^f + \lambda \operatorname{div} \mathbf{u}^s &= 0 & \text{in } \Omega \times (0, t_{\text{final}}], \\
\partial_t w_1 + \partial_t \mathbf{u}^s \cdot \nabla w_1 - \operatorname{div}(D_1(\mathbf{x}) \nabla w_1) &= f(w_1, w_2, \mathbf{u}^s) & \text{in } \Omega \times (0, t_{\text{final}}], \\
\partial_t w_2 + \partial_t \mathbf{u}^s \cdot \nabla w_2 - \operatorname{div}(D_2(\mathbf{x}) \nabla w_2) &= g(w_1, w_2, \mathbf{u}^s) & \text{in } \Omega \times (0, t_{\text{final}}],
\end{aligned} \tag{1.9}$$

which we endow with appropriate initial data at rest

$$w_1(0) = w_{1,0}, \quad w_2(0) = w_{2,0}, \quad \mathbf{u}^s(0) = \mathbf{0}, \quad p^f(0) = 0, \quad \psi(0) = 0 \quad \text{in } \Omega \times \{0\}, \tag{1.10}$$

and boundary conditions in the following manner

$$\mathbf{u}^s = \mathbf{0} \quad \text{and} \quad \frac{\kappa}{\eta} \nabla p^f \cdot \mathbf{n} = 0 \quad \text{on } \Gamma \times (0, t_{\text{final}}], \tag{1.11}$$

$$[2\mu \boldsymbol{\varepsilon}(\mathbf{u}^s) - \psi \mathbf{I} + \boldsymbol{\sigma}_{\text{act}}] \mathbf{n} = \mathbf{0} \quad \text{and} \quad p^f = 0 \quad \text{on } \Sigma \times (0, t_{\text{final}}], \tag{1.12}$$

$$D_1(\mathbf{x}) \nabla w_1 \cdot \mathbf{n} = 0 \quad \text{and} \quad D_2(\mathbf{x}) \nabla w_2 \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega \times (0, t_{\text{final}}], \tag{1.13}$$

where the boundary $\partial\Omega = \Gamma \cup \Sigma$ is disjointly split into Γ and Σ where we prescribe clamped boundaries and zero fluid normal fluxes; and zero (total) traction together with constant fluid pressure, respectively. Moreover, zero concentrations normal fluxes are prescribed on $\partial\Omega$. We point out that, if we would like to start with a model in terms of the divergence ($\operatorname{div}(w_i \partial_t \mathbf{u}^s)$ instead of $\partial_t \mathbf{u}^s \cdot \nabla w_i$ in (1.5)-(1.6), $i \in \{1, 2\}$), we need to assume zero total flux (including the advective term, see e.g. [5]). Homogeneity of the boundary conditions is only assumed to simplify the exposition of the subsequent analysis.

2. Well-posedness analysis

2.1. Weak formulation and a semi-discrete form

Let us multiply (1.9) by adequate test functions and integrate by parts (in space) whenever appropriate. Incorporating the boundary conditions (1.11)-(1.12) as well as the definition of the total stress (1.7), we end up with the following variational problem: For a given $t > 0$, find $\mathbf{u}^s(t) \in \mathbf{H}_\Gamma^1(\Omega)$, $p^f(t) \in H_\Sigma^1(\Omega)$, $\psi(t) \in L^2(\Omega)$, $w_1(t) \in H^1(\Omega)$, $w_2(t) \in H^1(\Omega)$ such that

$$\begin{aligned}
2\mu \int_\Omega \boldsymbol{\varepsilon}(\mathbf{u}^s) : \boldsymbol{\varepsilon}(\mathbf{v}^s) - \int_\Omega \psi \operatorname{div} \mathbf{v}^s &= \int_\Omega \rho \mathbf{b} \cdot \mathbf{v}^s + \int_\Omega \tau r \mathbf{k} \otimes \mathbf{k} : \boldsymbol{\varepsilon}(\mathbf{v}^s) \quad \forall \mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega), \\
\left(c_0 + \frac{\alpha^2}{\lambda}\right) \int_\Omega \partial_t p^f q^f + \frac{1}{\eta} \int_\Omega \kappa \nabla p^f \cdot \nabla q^f - \frac{\alpha}{\lambda} \int_\Omega \partial_t \psi q^f &= \int_\Omega \ell q^f \quad \forall q^f \in H_\Sigma^1(\Omega), \\
-\int_\Omega \phi \operatorname{div} \mathbf{u}^s + \frac{\alpha}{\lambda} \int_\Omega p^f \phi - \frac{1}{\lambda} \int_\Omega \psi \phi &= 0 \quad \forall \phi \in L^2(\Omega), \\
\int_\Omega \partial_t w_1 s_1 + \int_\Omega D_1 \nabla w_1 \cdot \nabla s_1 + \int_\Omega (\partial_t \mathbf{u}^s \cdot \nabla w_1) s_1 &= \int_\Omega f(w_1, w_2, \mathbf{u}^s) s_1 \quad \forall s_1 \in H^1(\Omega), \\
\int_\Omega \partial_t w_2 s_2 + \int_\Omega D_2 \nabla w_2 \cdot \nabla s_2 + \int_\Omega (\partial_t \mathbf{u}^s \cdot \nabla w_2) s_2 &= \int_\Omega g(w_1, w_2, \mathbf{u}^s) s_2 \quad \forall s_2 \in H^1(\Omega).
\end{aligned} \tag{2.1}$$

Next, let us discretise the time interval $(0, t_{\text{final}}]$ into equispaced points $t^n = n\Delta t$, and use the following general notation for the first order backward difference $\Delta t \delta_t X^{n+1} := X^{n+1} - X^n$. In this way, we can write a semidiscrete form of (2.1): From initial data $\mathbf{u}^{s,0}, p^{f,0}, \psi^0, w_1^0, w_2^0$ and for $n = 1, \dots$, find $\mathbf{u}^{s,n+1} \in \mathbf{H}_\Gamma^1(\Omega)$, $p^{f,n+1} \in H_\Sigma^1(\Omega)$, $\psi^{n+1} \in L^2(\Omega)$, $w_1^{n+1} \in H^1(\Omega)$, $w_2^{n+1} \in H^1(\Omega)$ such that

$$\begin{aligned}
a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) &+ b_1(\mathbf{v}^s, \psi^{n+1}) = F_{r^{n+1}}(\mathbf{v}^s) \quad \forall \mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega),
\end{aligned} \tag{2.2}$$

$$\tilde{a}_2(p^{f,n+1}, q^f) + a_2(p^{f,n+1}, q^f) - \tilde{b}_2(q^f, \psi^{n+1}) = G_{\ell^{n+1}}(q^f) \quad \forall q^f \in H_\Sigma^1(\Omega), \quad (2.3)$$

$$b_1(\mathbf{u}^{s,n+1}, \phi) + b_2(p^{f,n+1}, \phi) - a_3(\psi^{n+1}, \phi) = 0 \quad \forall \phi \in L^2(\Omega), \quad (2.4)$$

$$\tilde{a}_4(w_1^{n+1}, s_1) + a_4(w_1^{n+1}, s_1) + c(w_1^{n+1}, s_1, \mathbf{u}^{s,n+1}) = J_{f^{n+1}}(s_1) \quad \forall s_1 \in H^1(\Omega), \quad (2.5)$$

$$\tilde{a}_5(w_2^{n+1}, s_2) + a_5(w_2^{n+1}, s_2) + c(w_2^{n+1}, s_2, \mathbf{u}^{s,n+1}) = J_{g^{n+1}}(s_2) \quad \forall s_2 \in H^1(\Omega), \quad (2.6)$$

where the bilinear forms $a_1 : \mathbf{H}_\Gamma^1(\Omega) \times \mathbf{H}_\Gamma^1(\Omega) \rightarrow \mathbb{R}$, $a_2 : H_\Sigma^1(\Omega) \times H_\Sigma^1(\Omega) \rightarrow \mathbb{R}$, $a_3 : L^2(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$, $a_4, a_5 : H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{R}$, $b_1 : \mathbf{H}_\Gamma^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$, $b_2, \tilde{b}_2 : H_\Sigma^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$, the trilinear form $c : H^1(\Omega) \times H^1(\Omega) \times \mathbf{H}_\Gamma^1(\Omega) \rightarrow \mathbb{R}$, and linear functionals $F_r : \mathbf{H}_\Gamma^1(\Omega) \rightarrow \mathbb{R}$ (for r known), $G_\ell : H_\Sigma^1(\Omega) \rightarrow \mathbb{R}$, $J_f, J_g : H^1(\Omega) \rightarrow \mathbb{R}$ (for known f and known g), satisfy the following specifications

$$\begin{aligned} a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) &:= 2\mu \int_\Omega \boldsymbol{\varepsilon}(\mathbf{u}^{s,n+1}) : \boldsymbol{\varepsilon}(\mathbf{v}^s), \quad b_1(\mathbf{v}^s, \phi) := - \int_\Omega \phi \operatorname{div} \mathbf{v}^s, \quad b_2(p^{f,n+1}, \phi) := \frac{\alpha}{\lambda} \int_\Omega p^{f,n+1} \phi, \\ \tilde{a}_2(p^{f,n+1}, q^f) &:= \left(c_0 + \frac{\alpha^2}{\lambda}\right) \int_\Omega \delta_t p^{f,n+1} q^f, \quad a_2(p^{f,n+1}, q^f) := \frac{1}{\eta} \int_\Omega \kappa \nabla p^{f,n+1} \cdot \nabla q^f, \\ \tilde{b}_2(q^f, \psi^{n+1}) &:= \frac{\alpha}{\lambda} \int_\Omega \delta_t \psi^{n+1} q^f, \quad a_3(\psi^{n+1}, \phi) := \frac{1}{\lambda} \int_\Omega \psi^{n+1} \phi, \\ \tilde{a}_4(w_1^{n+1}, s_1) &:= \int_\Omega \delta_t w_1^{n+1} s_1, \quad a_4(w_1^{n+1}, s_1) := \int_\Omega D_1(\mathbf{x}) \nabla w_1^{n+1} \cdot \nabla s_1, \\ \tilde{a}_5(w_2^{n+1}, s_2) &:= \int_\Omega \delta_t w_2^{n+1} s_2, \quad a_5(w_2^{n+1}, s_2) := \int_\Omega D_2(\mathbf{x}) \nabla w_2^{n+1} \cdot \nabla s_2, \\ c(w, s, \mathbf{u}^{s,n+1}) &:= \int_\Omega (\delta_t \mathbf{u}^{s,n+1} \cdot \nabla w) s, \quad F_{r^{n+1}}(\mathbf{v}^s) := \rho \int_\Omega \mathbf{b}^{n+1} \cdot \mathbf{v}^s + \tau \int_\Omega r^{n+1} \mathbf{k} \otimes \mathbf{k} : \boldsymbol{\varepsilon}(\mathbf{v}^s), \\ G_{\ell^{n+1}}(q^f) &:= \int_\Omega \ell^{n+1} q^f, \quad J_{f^{n+1}}(s_1) := \int_\Omega f^{n+1} s_1, \quad J_{g^{n+1}}(s_2) := \int_\Omega g^{n+1} s_2. \end{aligned} \quad (2.7)$$

2.2. Preliminaries

We will consider that the initial data (1.10) are nonnegative and regular enough. Moreover, throughout the text we will assume that the anisotropic permeability $\kappa(\mathbf{x})$ and the diffusion matrices $D_1(\mathbf{x}), D_2(\mathbf{x})$ are uniformly bounded and positive definite in Ω . The latter means that, there exist positive constants κ_1, κ_2 , and $D_i^{\min}, D_i^{\max}$, $i \in \{1, 2\}$, such that

$$\kappa_1 |\mathbf{v}|^2 \leq \mathbf{v}^t \kappa(\mathbf{x}) \mathbf{v} \leq \kappa_2 |\mathbf{v}|^2, \quad \text{and} \quad D_i^{\min} |\mathbf{v}|^2 \leq \mathbf{v}^t D_i(\mathbf{x}) \mathbf{v} \leq D_i^{\max} |\mathbf{v}|^2 \quad \forall \mathbf{v} \in \mathbb{R}^d, \quad \forall \mathbf{x} \in \Omega.$$

Also, for a fixed $\mathbf{u}^s$, the reaction kinetics $f(w_1, w_2, \cdot), g(w_1, w_2, \cdot)$ satisfy the growth conditions

$$\begin{aligned} |f(w_1, w_2, \cdot)| &\leq C(1 + |w_1| + |w_2|), \quad |g(w_1, w_2, \cdot)| \leq C(1 + |w_1| + |w_2|) \quad \text{for } w_1, w_2 \geq 0, \\ |m(w_1, w_2, \cdot) - m(\tilde{w}_1, \tilde{w}_2, \cdot)| &\leq C(|w_1 - \tilde{w}_1| + |w_2 - \tilde{w}_2|) \quad \text{for } m = f, g, \\ f(w_1, w_2, \cdot) &= f_0 (\geq 0) \quad \text{and} \quad g(w_1, w_2, \cdot) = g_0 (\geq 0) \quad \text{if } w_1 \leq 0 \text{ or } w_2 \leq 0, \end{aligned} \quad (2.8)$$

and given $w_1, w_2 \in \mathbb{R}$, the scalar field $r(w_1, w_2)$ defined in (1.8) is such that

$$|r(w_1, w_2)| \leq |w_1| + |w_2|, \quad |r(w_1, w_2) - r(\tilde{w}_1, \tilde{w}_2)| \leq C(|w_1 - \tilde{w}_1| + |w_2 - \tilde{w}_2|). \quad (2.9)$$

In addition, according to [23], the terms in (2.2)-(2.6) fulfil the following continuity bounds

$$|a_1(\mathbf{u}^s, \mathbf{v}^s)| \leq 2\mu C_{k,2} \|\mathbf{u}^s\|_{1,\Omega} \|\mathbf{v}^s\|_{1,\Omega}, \quad |a_2(p^f, q^f)| \leq \frac{\kappa_2}{\eta} \|p^f\|_{1,\Omega} \|q^f\|_{1,\Omega},$$

$$\begin{aligned}
|a_3(\psi, \phi)| &\leq \lambda^{-1} \|\psi\|_{0,\Omega} \|\phi\|_{0,\Omega}, \quad |a_4(w_1, s_1)| \leq D_1^{\max} \|w_1\|_{1,\Omega} \|s_1\|_{1,\Omega}, \\
|a_5(w_2, s_2)| &\leq D_2^{\max} \|w_2\|_{1,\Omega} \|s_2\|_{1,\Omega}, \quad |b_1(\mathbf{v}^s, \phi)| \leq \sqrt{d} \|\mathbf{v}^s\|_{1,\Omega} \|\phi\|_{0,\Omega}, \\
|b_2(q^f, \phi)| &\leq \alpha \lambda^{-1} \|q^f\|_{1,\Omega} \|\phi\|_{0,\Omega}, \quad |F_r(\mathbf{v}^s)| \leq \rho \|\mathbf{b}\|_{0,\Omega} \|\mathbf{v}^s\|_{0,\Omega} + \tau \sqrt{C_{k,2}} \|r\|_{0,\Omega} \|\mathbf{v}^s\|_{1,\Omega}, \\
|G_\ell(q^f)| &\leq \|\ell\|_{0,\Omega} \|q^f\|_{0,\Omega}, \quad |J_f(s_1)| \leq \|f\|_{0,\Omega} \|s_1\|_{0,\Omega}, \quad |J_g(s_2)| \leq \|g\|_{0,\Omega} \|s_2\|_{0,\Omega},
\end{aligned} \tag{2.10}$$

for all $\mathbf{u}^s, \mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega)$, $p^f, q^f \in H_\Sigma^1(\Omega)$, $w_1, w_2, s_1, s_2 \in H^1(\Omega)$, $\psi, \phi \in L^2(\Omega)$. We also have the following coercivity and positivity bounds

$$\begin{aligned}
a_1(\mathbf{v}^s, \mathbf{v}^s) &\geq 2\mu C_{k,1} \|\mathbf{v}^s\|_{1,\Omega}^2, \quad a_2(q^f, q^f) \geq \frac{\kappa_1 c_p}{\eta} \|q^f\|_{1,\Omega}^2, \quad a_3(\phi, \phi) = \lambda^{-1} \|\phi\|_{0,\Omega}^2, \\
a_4(s_1, s_1) &\geq D_1^{\min} \|s_1\|_{1,\Omega}^2, \quad a_5(s_2, s_2) \geq D_2^{\min} \|s_2\|_{1,\Omega}^2,
\end{aligned} \tag{2.11}$$

for all $\mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega)$, $\phi \in L^2(\Omega)$, $s_1, s_2 \in H^1(\Omega)$, $q^f \in H_\Sigma^1(\Omega)$, where above $C_{k,1}$ and $C_{k,2}$ are the positive constants satisfying

$$C_{k,1} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 \leq \|\varepsilon(\mathbf{u}^{s,n+1})\|_{0,\Omega}^2 \leq C_{k,2} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2,$$

and c_p is the Poincaré constant. Moreover, the bilinear form b_1 satisfies the inf-sup condition (see e.g. [14]): For every $\phi \in L^2(\Omega)$, there exists $\beta > 0$ such that

$$\sup_{\mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega)} \frac{b_1(\mathbf{v}^s, \phi)}{\|\mathbf{v}^s\|_{1,\Omega}} \geq \beta \|\phi\|_{0,\Omega}. \tag{2.12}$$

Finally, we recall an important discrete identity and introduce the discrete-in-time norm

$$\int_\Omega X^{n+1} \delta_t X^{n+1} = \frac{1}{2} \delta_t \|X^{n+1}\|^2 + \frac{1}{2} \Delta t \|\delta_t X^{n+1}\|^2, \quad \|X\|_{\ell^2(V)}^2 := \Delta t \sum_{m=0}^n \|X^{m+1}\|_V^2, \tag{2.13}$$

respectively, which will be useful for the subsequent analysis.

2.3. Fredholm alternative and a fixed-point structure

We address the unique solvability and continuous dependence on data for the semi-discrete system (2.2)-(2.6). As in [5] we define adequate sets to be used in the subsequent analysis. For $i = 1, 2$ and $\forall t = t_n, n = 0, 1, \dots, N$ let

$$\mathcal{S} := \mathbf{D} \times \mathbf{D}, \quad \text{where } \mathbf{D} := \{w_i(\mathbf{x}, \cdot) \in L^2(\Omega) : 0 \leq w_i(\mathbf{x}, t_n) \leq e^{(\gamma-\nu)t_n} M \text{ for a.e. } \mathbf{x} \in \Omega\},$$

and where M is a constant that satisfies $M \geq \sup\{\|w_{1,0}\|_{\infty,\Omega}, \|w_{2,0}\|_{\infty,\Omega}\}$. In addition, the constants γ, ν will be defined as per the requirement of the analysis. It is evident that $\mathcal{S}$ is a closed, bounded and convex subset of the Banach space $[L^2(\Omega)]^2$. From system (2.2)-(2.6) we then define two uncoupled subproblems. For a given concentration pair $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in \mathcal{S}$, find a solution pair $(w_1^{n+1}, w_2^{n+1}) \in [H^1(\Omega)]^2$ of the following uncoupled advection-diffusion-reaction (ADR) system:

$$\begin{aligned}
\tilde{a}_4(w_1^{n+1}, s_1) + a_4(w_1^{n+1}, s_1) + c(w_1^{n+1}, s_1, \mathbf{u}^{s,n+1}) &= J_{f^{n+1}}(s_1) \quad \forall s_1 \in H^1(\Omega), \\
\tilde{a}_5(w_2^{n+1}, s_2) + a_5(w_2^{n+1}, s_2) + c(w_2^{n+1}, s_2, \mathbf{u}^{s,n+1}) &= J_{g^{n+1}}(s_2) \quad \forall s_2 \in H^1(\Omega).
\end{aligned} \tag{2.14}$$

In the above system, $\mathbf{u}^{s,n+1}$ is the solution of the following uncoupled poroelastic problem:

$$\begin{aligned}
a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) + b_1(\mathbf{v}^s, \psi^{n+1}) &= F_{\hat{r}^{n+1}}(\mathbf{v}^s) \quad \forall \mathbf{v}^s \in \mathbf{H}_\Gamma^1(\Omega), \\
\tilde{a}_2(p^{f,n+1}, q^f) + a_2(p^{f,n+1}, q^f) - \tilde{b}_2(q^f, \psi^{n+1}) &= G_{\ell^{n+1}}(q^f) \quad \forall q^f \in H_\Sigma^1(\Omega), \\
b_1(\mathbf{u}^{s,n+1}, \phi) + b_2(p^{f,n+1}, \phi) - a_3(\psi^{n+1}, \phi) &= 0 \quad \forall \phi \in L^2(\Omega),
\end{aligned} \tag{2.15}$$

for given $\hat{r}^{n+1} := r(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$. We first show that (2.14) and (2.15) are well-posed.

Lemma 2.1. Assume that $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in \mathcal{S}$. Then problem (2.15) has a unique solution

$$(\mathbf{u}^{s,n+1}, p^{f,n+1}, \psi^{n+1}) \in \mathbb{V} := \mathbf{H}_\Gamma^1(\Omega) \times H_\Sigma^1(\Omega) \times L^2(\Omega).$$

Moreover, there exists $C > 0$, independent of Δt and λ , such that for each n ,

$$\begin{aligned} & \|\mathbf{u}^{s,n+1}\|_{1,\Omega} + \sqrt{c_0} \|p^{f,n+1}\|_{0,\Omega} + \|\psi^{n+1}\|_{0,\Omega} + \|p^f\|_{L^2(H^1(\Omega))} \\ & \leq C \sqrt{\exp} \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega} + \|p^{f,0}\|_{0,\Omega} + \|\psi^0\|_{0,\Omega} + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega} + \|\ell\|_{\ell^2(L^2(\Omega))} + \sum_{m=0}^n \|\hat{r}^{m+1}\|_{0,\Omega} \right\}. \end{aligned} \quad (2.16)$$

Proof. The main ideas are borrowed from [23], which focuses on steady poromechanics, but possessing a similar structure to (2.15). In view of putting the formulation in operator form (amenable for analysis through the Fredholm alternative) we define, for $\vec{\mathbf{u}} = (\mathbf{u}^{s,n+1}, p^{f,n+1}, \psi^{n+1}) \in \mathbb{V}$, $\vec{\mathbf{v}} = (\mathbf{v}^s, q^f, \phi) \in \mathbb{V}$

$$\begin{aligned} \langle \mathcal{A}(\vec{\mathbf{u}}), \vec{\mathbf{v}} \rangle &:= a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) + b_1(\mathbf{v}^s, \psi^{n+1}) - b_1(\mathbf{u}^{s,n+1}, \phi) + \tilde{a}_2(p^{f,n+1}, q^f) \\ &\quad + a_2(p^{f,n+1}, q^f) + a_3(\psi^{n+1}, \phi), \\ \langle \mathcal{K}(\vec{\mathbf{u}}), \vec{\mathbf{v}} \rangle &:= -b_2(p^{f,n+1}, \phi) - \tilde{b}_2(q^f, \psi^{n+1}), \\ \langle \mathcal{F}, \vec{\mathbf{v}} \rangle &:= F_{\hat{r}^{n+1}}(\mathbf{v}^s) + G_{\ell^{n+1}}(q^f). \end{aligned}$$

As per the Fredholm alternative, the solvability of the operator problem $(\mathcal{A} + \mathcal{K})\vec{\mathbf{u}} = \mathcal{F}$ (which implies solvability of the uncoupled problem (2.15)), holds if $\mathcal{K}$ is compact, $\mathcal{A}$ is invertible and $\mathcal{A} + \mathcal{K}$ is injective.

Step 1. $\mathcal{K}$ is compact: Define an operator $\mathbb{B}_2 : H^1(\Omega) \rightarrow L^2(\Omega)$ such that $\langle \mathbb{B}_2(q^f), \phi \rangle := b_2(q^f, \phi)$, that is, $\mathbb{B}_2 q^f = (\frac{\alpha}{\lambda} I) \circ i_c$ where $i_c : H^1(\Omega) \rightarrow L^2(\Omega)$ is compact using Rellich-Kondrachov Theorem and $I : L^2(\Omega) \rightarrow L^2(\Omega)$ is the identity map. It implies that $\mathbb{B}_2$ is compact, so is $\mathbb{B}_2^*$. Note that $\mathcal{K}(\vec{\mathbf{u}}) = (0, \mathbb{B}_2(p^{f,n+1}), -\mathbb{B}_2^*(\delta_t \psi^{n+1}))$. Thus, $\mathcal{K}$ is compact.

Step 2. $\mathcal{A}$ is invertible and $(\mathcal{A} + \mathcal{K})$ is injective: Assume $\mathbf{V} := \mathbf{H}_\Gamma^1(\Omega)$, $Q := H_\Sigma^1(\Omega)$ and $Z := L^2(\Omega)$. The invertibility of $\mathcal{A}$ is equivalent to the existence of a unique solution to the operator problem: Given $\mathcal{L} := (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3) \in \mathbb{V}$, find $\vec{\mathbf{u}} \in \mathbb{V}$ such that $\mathcal{A}\vec{\mathbf{u}} = \mathcal{L}$, which is equivalent to the two uncoupled problems:

- Find $(\mathbf{u}^{s,n+1}, \psi^{n+1}) \in \mathbf{V} \times Z$ such that

$$\begin{aligned} a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) + b_1(\mathbf{v}^s, \psi^{n+1}) &= \mathcal{L}_1(\mathbf{v}^s) \quad \forall \mathbf{v}^s \in \mathbf{V}, \\ b_1(\mathbf{u}^{s,n+1}, \phi) - a_3(\psi^{n+1}, \phi) &= \mathcal{L}_3(\phi) \quad \forall \phi \in Z, \end{aligned} \quad (2.17)$$

- Find $p^{f,n+1} \in Q$ such that

$$\tilde{a}_2(p^{f,n+1}, q^f) + a_2(p^{f,n+1}, q^f) = \mathcal{L}_2(q^f) \quad \forall q^f \in Q. \quad (2.18)$$

The continuity and coercivity of the bilinear forms $a_1(\cdot, \cdot)$, $b_1(\cdot, \cdot)$ and $a_3(\cdot, \cdot)$, in combination with the inf-sup condition for $b_1(\cdot, \cdot)$ and the semi-positive definiteness of $a_3(\cdot, \cdot)$, ensure the unique solvability of (2.17) (see [13]). Moreover, in view of the coercivity of $a_2(\cdot, \cdot)$ and the classical result from [24], the existence of a unique solution to (2.18) can be easily shown. Therefore $\mathcal{A}$ is invertible. Furthermore, analogously to the proof of [23, Lemma 2.4], it is straightforward to show that $\mathcal{A} + \mathcal{K}$ is one-to-one.

Next we show the continuous dependence on data. We begin by taking $\mathbf{v}_h^s = \delta_t \mathbf{u}_h^{s,n+1}$ in the first row of (2.15), and then applying Cauchy-Schwarz and Young inequalities, to get

$$\begin{aligned} \mu \delta_t \|\varepsilon(\mathbf{u}^{s,n+1})\|_{0,\Omega}^2 + \mu C_{k,1} \Delta t \|\delta_t \mathbf{u}^{s,n+1}\|_{1,\Omega}^2 &\leq \frac{1}{2\delta_1} \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{\delta_1}{2} \|\delta_t \mathbf{u}^{s,n+1}\|_{1,\Omega}^2 \\ &\quad + \frac{\tau^2}{2\delta_2} \|\hat{r}^{n+1}\|_{0,\Omega}^2 + \frac{C_{k,2}\delta_2}{2} \|\delta_t \mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \frac{\rho^2}{2\delta_3} \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 + \frac{\delta_3}{2} \|\delta_t \mathbf{u}^{s,n+1}\|_{1,\Omega}^2. \end{aligned}$$

Next, defining $\delta_1 := \frac{\mu C_{k,1} \Delta t}{2}$, $\delta_2 := \frac{\mu C_{k,1} \Delta t}{2 C_{k,2}}$ and $\delta_3 := \frac{\mu C_{k,1} \Delta t}{2}$, and then, multiplying the resulting inequality by Δt and summing over n , we finally obtain

$$\begin{aligned} & \mu C_{k,1} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \frac{\mu C_{k,1} \Delta t^2}{4} \sum_{m=0}^n \|\delta_t \mathbf{u}^{s,m+1}\|_{1,\Omega}^2 \\ & \leq C_1 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\hat{r}^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 \right\}, \end{aligned} \quad (2.19)$$

where C_1 is a constant depending on μ , $C_{k,1}$, $C_{k,2}$, ρ , and τ . On the other hand, by taking $q = p^{f,n+1}$ and $\phi = \delta_t \psi^{n+1}$ in the second and third equation of (2.15), respectively, we get

$$\begin{aligned} & \frac{1}{2\lambda} \delta_t \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2\lambda} \|\delta_t \psi^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\delta_t \|p^{f,n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t p^{f,n+1}\|_{0,\Omega}^2 \right) + \frac{\kappa_1}{\eta} |p^{f,n+1}|_{1,\Omega}^2 \\ & \leq \frac{2\alpha}{\lambda} \|p^{f,n+1}\|_{0,\Omega} \|\delta_t \psi^{n+1}\|_{0,\Omega} + \|\ell^{n+1}\|_{0,\Omega} \|p^{f,n+1}\|_{0,\Omega} - \int_{\Omega} \delta_t \psi^{n+1} \operatorname{div} \mathbf{u}^{s,n+1}. \end{aligned} \quad (2.20)$$

Employing then Young's inequality in the first two terms on the right-hand side of (2.20), we obtain

$$\begin{aligned} & \frac{1}{2\lambda} \delta_t \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2\lambda} \|\delta_t \psi^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\delta_t \|p^{f,n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t p^{f,n+1}\|_{0,\Omega}^2 \right) + \frac{\kappa_1}{\eta} |p^{f,n+1}|_{1,\Omega}^2 \\ & \leq \frac{\delta_1}{\lambda} \|\delta_t \psi^{n+1}\|_{0,\Omega}^2 + \frac{\alpha^2}{\lambda \delta_1} \|p^{f,n+1}\|_{0,\Omega}^2 + \frac{1}{2\delta_2} \|\ell^{n+1}\|_{0,\Omega}^2 + \frac{\delta_2}{2} \|p^{f,n+1}\|_{0,\Omega}^2 - \int_{\Omega} \delta_t \psi^{n+1} \operatorname{div} \mathbf{u}^{s,n+1}. \end{aligned}$$

Now, choosing $\delta_1 := \frac{\Delta t}{2}$ and $\delta_2 := \frac{\kappa_1 c_p}{\eta}$, and then, multiplying the resulting inequality by Δt and summing over n , we deduce the following preliminar bound

$$\begin{aligned} & \frac{1}{2\lambda} \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\|p^{f,n+1}\|_{0,\Omega}^2 + \Delta t^2 \sum_{m=0}^n \|\delta_t p^{f,m+1}\|_{0,\Omega}^2 \right) + \frac{\kappa_1 c_p \Delta t}{2\eta} \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2 \\ & \leq \frac{1}{2\lambda} \|\psi^0\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \|p^{f,0}\|_{0,\Omega}^2 + \frac{2\alpha^2}{\lambda} \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 + \frac{\eta \Delta t}{2\kappa_1 c_p} \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \\ & \quad - \Delta t \sum_{m=0}^n \int_{\Omega} \delta_t \psi^{m+1} \operatorname{div} \mathbf{u}^{s,m+1}. \end{aligned} \quad (2.21)$$

Finally, for the last term on the right hand side of (2.21), we proceed similar to [4, Section 9], applying summation by parts and the initial conditions (1.10), to obtain that

$$\begin{aligned} & - \Delta t \sum_{m=0}^n \int_{\Omega} \delta_t \psi^{m+1} \operatorname{div} \mathbf{u}^{s,m+1} = - \int_{\Omega} \psi^{n+1} \operatorname{div} \mathbf{u}^{s,n+1} + \Delta t \sum_{m=0}^{n-1} \int_{\Omega} \psi^{m+1} \delta_t \operatorname{div} \mathbf{u}^{s,m+1} \\ & \leq \frac{1}{2\delta_3} \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{\delta_3}{2} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \frac{1}{2\delta_4} \Delta t \sum_{m=0}^{n-1} \|\psi^{m+1}\|_{0,\Omega}^2 + \frac{\delta_4}{2} \Delta t \sum_{m=0}^{n-1} \|\delta_t \mathbf{u}^{s,m+1}\|_{1,\Omega}^2, \end{aligned}$$

and then, taking $\delta_3 := \mu C_{k,1}$ and $\delta_4 := \frac{\mu C_{k,1} \Delta t}{2}$, we arrive at the following estimate

$$\begin{aligned} & \frac{1}{2\lambda} \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\|p^{f,n+1}\|_{0,\Omega}^2 + \Delta t^2 \sum_{m=0}^n \|\delta_t p^{f,m+1}\|_{0,\Omega}^2 \right) + \frac{\kappa_1 c_p \Delta t}{2\eta} \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2 \\ & \leq \frac{1}{2\lambda} \|\psi^0\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \|p^{f,0}\|_{0,\Omega}^2 + \frac{2\alpha^2}{\lambda} \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 + \frac{\eta \Delta t}{2\kappa_1 c_p} \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \end{aligned} \quad (2.22)$$

$$+ \frac{1}{2\mu C_{k,1}} \|\psi^{n+1}\|_{0,\Omega}^2 + \frac{\mu C_{k,1}}{2} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \frac{1}{\mu C_{k,1}} \sum_{m=0}^{n-1} \|\psi^{m+1}\|_{0,\Omega}^2 + \frac{\mu C_{k,1} \Delta t^2}{4} \sum_{m=0}^{n-1} \|\delta_t \mathbf{u}^{s,m+1}\|_{1,\Omega}^2.$$

In this way, by adding (2.19) and (2.22), we get

$$\begin{aligned} & \frac{\mu C_{k,1}}{2} \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \frac{c_0}{2} \|p^{f,n+1}\|_{0,\Omega}^2 + \frac{\kappa_1 c_p \Delta t}{2\eta} \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2 \\ & \leq C_1 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|r^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 \right\} + \left(c_0 + \frac{1}{2\lambda} \right) \|\psi^0\|_{0,\Omega}^2 \\ & \quad + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \|p^{f,0}\|_{0,\Omega}^2 + \left(c_0 + \frac{2\alpha^2}{\lambda} \right) \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 \\ & \quad + \frac{\eta \Delta t}{2\kappa_1 c_p} \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 + \frac{2}{\mu C_{k,1}} \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 \\ & \leq C_2 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \|p^{f,0}\|_{0,\Omega}^2 + \|\psi^0\|_{0,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 \right. \\ & \quad \left. + \sum_{m=0}^n \|\hat{r}^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \right\}, \end{aligned} \quad (2.23)$$

where $C_2 := \max\{C_1, c_0 + \frac{1}{2\lambda}, \frac{1}{2}(c_0 + \frac{\alpha^2}{\lambda}), c_0 + \frac{2\alpha^2}{\lambda}, \frac{\eta}{2\kappa_1 c_p}, \frac{2}{\mu C_{k,1}}\}$ must be understood as a constant independent of λ if λ goes to infinity. On the other hand, with the aim to obtain a bound for $\|\psi^{n+1}\|_{0,\Omega}$ independent of λ , we propose to use the inf-sup condition (2.12). Thus, taking $\phi = \psi^{n+1}$, using the first row of (2.15) and the continuity of a_1 , we easily obtain

$$\begin{aligned} \beta \|\psi^{n+1}\|_{0,\Omega} & \leq \sup_{\mathbf{v}^s \in \mathbf{V}} \frac{b_1(\mathbf{v}^s, \psi^{n+1})}{\|\mathbf{v}^s\|_{1,\Omega}} = \sup_{\mathbf{v}^s \in \mathbf{V}} \frac{-a_1(\mathbf{u}^{s,n+1}, \mathbf{v}^s) + F_{\hat{r}^{n+1}}(\mathbf{v}^s)}{\|\mathbf{v}^s\|_{1,\Omega}} \\ & \leq 2\mu C_{k,2} \|\varepsilon(\mathbf{u}^{s,n+1})\|_{0,\Omega} + \sqrt{C_{k,2}\tau} \|\hat{r}^{n+1}\|_{0,\Omega} + \rho \|\mathbf{b}^{n+1}\|_{0,\Omega}, \end{aligned}$$

or what is the same

$$\|\psi^{n+1}\|_{0,\Omega}^2 \leq C_3 \left\{ \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + \|\hat{r}^{n+1}\|_{0,\Omega}^2 + \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 \right\}, \quad (2.24)$$

where C_3 is a constant depending on $\beta, C_{k,1}, C_{k,2}, \mu, \tau$ and ρ . From (2.23) and (2.24) we finally obtain an estimate concerning the stability of the poroelasticity problem

$$\begin{aligned} & \|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + c_0 \|p^{f,n+1}\|_{0,\Omega}^2 + \|\psi^{n+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2 \\ & \leq C_4 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \|p^{f,0}\|_{0,\Omega}^2 + \|\psi^0\|_{0,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 \right. \\ & \quad \left. + \sum_{m=0}^n \|\hat{r}^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \right\} + C_3 \left\{ \|\hat{r}^{n+1}\|_{0,\Omega}^2 + \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 \right\}. \end{aligned} \quad (2.25)$$

Finally, the stability result (2.16) follows by applying Gronwall's inequality to (2.25). $\square$

Lemma 2.2. For any $\mathbf{u}^{s,n+1} \in \mathbf{V}$, the uncoupled ADR system (2.14) has a unique solution. Moreover there exists $C > 0$, independent of Δt , such that for each n ,

$$\|w_1^{n+1}\|_{0,\Omega} + \|w_2^{n+1}\|_{0,\Omega} + \|\nabla w_1\|_{\ell^2(L^2(\Omega))} + \|\nabla w_2\|_{\ell^2(L^2(\Omega))} \leq C \sqrt{\exp} \left\{ n \Delta t + \|w_1^0\|_{0,\Omega} + \|w_2^0\|_{0,\Omega} \right\}. \quad (2.26)$$

Proof. Note that for each n , the uncoupled ADR equations constitute a semilinear elliptic system; and owing to the uniform boundedness of the matrices $D_i(\mathbf{x})$, $i = 1, 2$ together with the growth condition assumed for f, g ; the problem (2.14) is uniquely solvable (see for instance, [18]). On the other hand, for the continuous dependence, we begin by taking $s_1 = w_1^{n+1}$ in the first equation of (2.14), which yields

$$\int_{\Omega} \delta_t w_1^{n+1} w_1^{n+1} + \int_{\Omega} D_1(\mathbf{x}) \nabla w_1^{n+1} \cdot \nabla w_1^{n+1} + \int_{\Omega} (\delta_t \mathbf{u}^{s,n+1} \cdot \nabla w_1^{n+1}) w_1^{n+1} = \int_{\Omega} f^{n+1} w_1^{n+1},$$

and then, recalling that

$$\int_{\Omega} (\delta_t \mathbf{u}^{s,n+1} \cdot \nabla w_1^{n+1}) w_1^{n+1} = -\frac{1}{2} \int_{\Omega} \operatorname{div} (\delta_t \mathbf{u}^{s,n+1}) (w_1^{n+1})^2, \quad (2.27)$$

we can apply classical Cauchy-Schwarz inequality, to obtain

$$\begin{aligned} \frac{1}{2} \delta_t \|w_1^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t \|\delta_t w_1^{n+1}\|_{0,\Omega}^2 + D_1^{\min} \|\nabla w_1^{n+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|\delta_t \mathbf{u}^{s,n+1}\|_{1,\infty,\Omega} \|w_1^{n+1}\|_{0,\Omega}^2 + \|f^{n+1}\|_{0,\Omega} \|w_1^{n+1}\|_{0,\Omega}. \end{aligned}$$

Under the assumption that $\mathbf{u}^{s,n+1}, \mathbf{u}^{s,n}$ are uniformly bounded in $\mathbf{W}^{1,\infty}(\Omega)$, and after applying Young's inequality, we deduce the following result

$$\begin{aligned} \frac{1}{2} \delta_t \|w_1^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t \|\delta_t w_1^{n+1}\|_{0,\Omega}^2 + D_1^{\min} \|\nabla w_1^{n+1}\|_{0,\Omega}^2 \\ \leq \frac{C_1}{2\Delta t} \|w_1^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \|f^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \|w_1^{n+1}\|_{0,\Omega}^2. \end{aligned}$$

Finally, a preliminary stability result follows by summing over n and multiplying by Δt , which is

$$\begin{aligned} \frac{1}{2} \|w_1^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t^2 \sum_{m=0}^n \|\delta_t w_1^{m+1}\|_{0,\Omega}^2 + D_1^{\min} \Delta t \sum_{m=0}^n \|\nabla w_1^{m+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|w_1^0\|_{0,\Omega}^2 + \frac{1}{2} (C_1 + \Delta t) \sum_{m=0}^n \|w_1^{m+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2} \sum_{m=0}^n \|f^{m+1}\|_{0,\Omega}^2. \end{aligned} \quad (2.28)$$

In much the same way as above, we obtain a stability result for $\|w_2^{n+1}\|_{0,\Omega}$

$$\begin{aligned} \frac{1}{2} \|w_2^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t^2 \sum_{m=0}^n \|\delta_t w_2^{m+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|\nabla w_2^{m+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|w_2^0\|_{0,\Omega}^2 + \frac{1}{2} (C_1 + \Delta t) \sum_{m=0}^n \|w_2^{m+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2} \sum_{m=0}^n \|g^{m+1}\|_{0,\Omega}^2, \end{aligned} \quad (2.29)$$

and then, from (2.28) and (2.29), we get a stability bound for the uncoupled problem (2.14)

$$\begin{aligned} \frac{1}{2} \|w_1^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \|w_2^{n+1}\|_{0,\Omega}^2 + D^{\min} \Delta t \sum_{m=0}^n (\|\nabla w_1^{m+1}\|_{0,\Omega}^2 + \|\nabla w_2^{m+1}\|_{0,\Omega}^2) \\ \leq C_2 \left\{ n\Delta t + \|w_1^0\|_{0,\Omega}^2 + \|w_2^0\|_{0,\Omega}^2 + \sum_{m=0}^n \left(\|w_1^{m+1}\|_{0,\Omega}^2 + \|w_2^{m+1}\|_{0,\Omega}^2 \right) \right\}, \end{aligned} \quad (2.30)$$

where we have used the growth condition on f and g , and $D^{\min} := \min\{D_1^{\min}, D_2^{\min}\}$. Finally, the stability of (2.14) given by (2.26) follows from an application of Gronwall's inequality to (2.30). $\square$

2.4. Existence of a weak solution

The next task consists in showing that (2.2)-(2.6) has a solution. We define auxiliary space-dependent functions m_{w_1}, m_{w_2} in such a way that the solutions of the problem can be expanded as

$$w_1 = e^{-\nu t} m_{w_1}, w_2 = e^{-\nu t} m_{w_2},$$

for some constant $\nu > 0$. Then it is readily seen that w_1, w_2 will satisfy

$$\begin{aligned} \partial_t w_1 - \operatorname{div}(D_1(\mathbf{x}) \nabla w_1) + \partial_t \mathbf{u}^s \cdot \nabla w_1 &= -\nu w_1 + e^{\nu t} f(e^{\nu t} w_1, e^{\nu t} w_2), \\ \partial_t w_2 - \operatorname{div}(D_2(\mathbf{x}) \nabla w_2) + \partial_t \mathbf{u}^s \cdot \nabla w_2 &= -\nu w_2 + e^{\nu t} g(e^{\nu t} w_1, e^{\nu t} w_2), \end{aligned}$$

whose semi-discrete variational counterpart is: Find w_1^{n+1}, w_2^{n+1} such that

$$\begin{aligned} \int_{\Omega} \delta_t w_1^{n+1} s_1 + \int_{\Omega} D_1(\mathbf{x}) \nabla w_1^{n+1} \cdot \nabla s_1 + \int_{\Omega} (\delta_t \mathbf{u}^{s,n+1} \cdot \nabla w_1^{n+1}) s_1 \\ = -\nu \int_{\Omega} w_1^{n+1} s_1 + \int_{\Omega} e^{\nu t_{n+1}} f(e^{\nu t_{n+1}} w_1^{n+1}, e^{\nu t_{n+1}} w_2^{n+1}) s_1 \quad \forall s_1 \in H^1(\Omega), \end{aligned} \quad (2.31)$$

$$\begin{aligned} \int_{\Omega} \delta_t w_2^{n+1} s_2 + \int_{\Omega} D_2(\mathbf{x}) \nabla w_2^{n+1} \cdot \nabla s_2 + \int_{\Omega} (\delta_t \mathbf{u}^{s,n+1} \cdot \nabla w_2^{n+1}) s_2 \\ = -\nu \int_{\Omega} w_2^{n+1} s_2 + \int_{\Omega} e^{\nu t_{n+1}} g(e^{\nu t_{n+1}} w_1^{n+1}, e^{\nu t_{n+1}} w_2^{n+1}) s_2 \quad \forall s_2 \in H^1(\Omega). \end{aligned} \quad (2.32)$$

The system can be equivalently stated in the form

$$\tilde{a}_4(w_1^{n+1}, s_1) + a_4(w_1^{n+1}, s_1) + c(w_1^{n+1}, s_1, \mathbf{u}^{s,n+1}) = \tilde{f}_{f^{n+1}}(s_1) \quad \forall s_1 \in H^1(\Omega), \quad (2.33)$$

$$\tilde{a}_5(w_2^{n+1}, s_2) + a_5(w_2^{n+1}, s_2) + c(w_2^{n+1}, s_2, \mathbf{u}^{s,n+1}) = \tilde{f}_{g^{n+1}}(s_2) \quad \forall s_2 \in H^1(\Omega), \quad (2.34)$$

where

$$\begin{aligned} \tilde{f}_{f^{n+1}}(s_1) &= -\nu \int_{\Omega} w_1^{n+1} s_1 + \int_{\Omega} e^{-\nu t_{n+1}} f(e^{\nu t_{n+1}} w_1^{n+1}, e^{\nu t_{n+1}} w_2^{n+1}) s_1 \\ \tilde{f}_{g^{n+1}}(s_2) &= -\nu \int_{\Omega} w_2^{n+1} s_2 + \int_{\Omega} e^{-\nu t_{n+1}} g(e^{\nu t_{n+1}} w_1^{n+1}, e^{\nu t_{n+1}} w_2^{n+1}) s_2. \end{aligned}$$

Now we define the operator $T : \mathcal{S} \rightarrow \mathcal{S}$ that for each n gives

$$T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) = (w_1^{n+1}, w_2^{n+1}),$$

for a fixed pair $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in \mathcal{S}$, and where $(w_1^{n+1}, w_2^{n+1}) \in [H^1(\Omega)]^2$ is the solution of (2.33)-(2.34) with a given displacement $\mathbf{u}^{s,n+1}$ (that is, the solution of the uncoupled poroelastic problem (2.15)). Our objective is to show that T has a fixed point and in consequence a solution to (2.2)-(2.6) exists. For accomplishment of this purpose, we appeal to the Schauder fixed-point theorem, and all necessary ingredients are collected in what follows.

Lemma 2.3. *The operator T is a self map.*

Proof. For given $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in \mathcal{S}$, we need to show that $0 \leq w_1^{n+1}, w_2^{n+1} \leq e^{(\gamma-\nu)t_{n+1}} M$ for each $n = 0, 1, \dots, N$ where $(w_1^{n+1}, w_2^{n+1}) = T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$. The proof is based on induction and contradiction arguments. Given $w_{1,0} \geq 0$, assume that $w_1^n \geq 0$. We then suppose that $w_1^{n+1} < 0$. Setting $s_1 = -(w_1^{n+1})^- = -\max\{-w_1^{n+1}, 0\}$ in (2.31) gives us

$$-\int_{\Omega} \left(\frac{w_1^{n+1} - w_1^n}{\Delta t} \right) (w_1^{n+1})^- - \int_{\Omega} D_1(\mathbf{x}) \nabla w_1^{n+1} \cdot \nabla (w_1^{n+1})^- - \int_{\Omega} \left(\frac{\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n}}{\Delta t} \cdot \nabla w_1^{n+1} \right) (w_1^{n+1})^-$$

$$\begin{aligned}
&= \nu \int_{\Omega} w_1^{n+1} (w_1^{n+1})^- - \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} (w_1^{n+1})^-, \\
\frac{1}{\Delta t} \int_{\Omega} ((w_1^{n+1})^-)^2 + D_1^{\min} \int_{\Omega} (\nabla (w_1^{n+1})^-)^2 + \int_{\Omega} \left(\frac{\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n}}{2\Delta t} \right) \cdot \nabla ((w_1^{n+1})^-)^2 + \frac{1}{\Delta t} \int_{\Omega} w_1^n (w_1^{n+1})^- \\
&= -\nu \int_{\Omega} ((w_1^{n+1})^-)^2 - \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} (w_1^{n+1})^-,
\end{aligned}$$

and therefore

$$\begin{aligned}
\frac{1}{\Delta t} \int_{\Omega} ((w_1^{n+1})^-)^2 + D_1^{\min} \int_{\Omega} (\nabla (w_1^{n+1})^-)^2 - \int_{\Omega} \left(\frac{\operatorname{div}(\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n})}{2\Delta t} \right) ((w_1^{n+1})^-)^2 + \nu \int_{\Omega} ((w_1^{n+1})^-)^2 \\
= -\frac{1}{\Delta t} \int_{\Omega} w_1^n (w_1^{n+1})^- - \int_{\Omega} e^{-\nu t_{n+1}} (w_1^{n+1})^- f_0.
\end{aligned}$$

Since w_1^n and f_0 are non-negative, the right-hand side of the last equation is non-positive. For $\nu \geq \frac{\|\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n}\|_{1,\infty,\Omega}}{2\Delta t}$ (which is legitimate as can be seen at the end of the proof) along with positive definiteness of $D_1(\mathbf{x})$ throughout Ω implies that $\int_{\Omega} ((w_1^{n+1})^-)^2 \leq 0$; and hence $(w_1^{n+1})^- = 0$. However $(w_1^{n+1})^- > 0$, which contradicts our initial assumption. Proceeding then by induction we obtain that $w_1^{n+1} \geq 0$ for each n . The property for w_2 can be derived in analogous way.

The other part of the inequality (that is, $w_1^n, w_2^n \leq e^{(\gamma-\nu)t_n} M$ for each n) follows the same lines. Given $w_{1,0} \leq M$ we assume that $w_1^n \leq e^{(\gamma-\nu)t_n} M \leq e^{(\gamma-\nu)t_{n+1}} M$, and we further suppose that $w_1^{n+1} > e^{(\gamma-\nu)t_{n+1}} M$. Choosing $s_1 = s_1^{n+1} := (w_1^{n+1} - e^{(\gamma-\nu)t_{n+1}} M)^+$ in (2.33), we can readily obtain

$$\begin{aligned}
\frac{1}{\Delta t} \int_{\Omega} (w_1^{n+1} - w_1^n) s_1^{n+1} + \int_{\Omega} D_1(\mathbf{x}) \nabla w_1^{n+1} \cdot \nabla s_1^{n+1} + \int_{\Omega} \frac{(\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n})}{\Delta t} \cdot \nabla w_1^{n+1} s_1^{n+1} \\
= -\nu \int_{\Omega} w_1^{n+1} s_1^{n+1} + \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} s_1^{n+1},
\end{aligned}$$

which implies that

$$\begin{aligned}
\frac{1}{\Delta t} \int_{\Omega} (s_1^{n+1})^2 + D_1^{\min} \int_{\Omega} |\nabla s_1^{n+1}|^2 - \int_{\Omega} \frac{\operatorname{div}(\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n})}{2\Delta t} (s_1^{n+1})^2 - \int_{\Omega} \frac{(w_1^n - e^{(\gamma-\nu)t_{n+1}} M)}{\Delta t} s_1^{n+1} \\
\leq -\nu \int_{\Omega} (s_1^{n+1})^2 - \nu \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} s_1^{n+1}.
\end{aligned}$$

Using again that $D_1^{\min} > 0$ and the growth condition of f and $w_1^n \leq e^{(\gamma-\nu)t_{n+1}} M$, we can assert that

$$\begin{aligned}
\frac{1}{\Delta t} \int_{\Omega} (s_1^{n+1})^2 + \int_{\Omega} \left(\nu - \frac{\|\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n}\|_{1,\infty,\Omega}}{2\Delta t} \right) (s_1^{n+1})^2 + \nu \int_{\Omega} e^{(\gamma-\nu)t_{n+1}} M s_1^{n+1} \\
\leq -\nu \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} s_1^{n+1} \leq C e^{-\nu t_{n+1}} \int_{\Omega} (1 + |w_1^{n+1}| + |w_2^{n+1}|) s_1^{n+1} \\
\leq C e^{-\nu t_{n+1}} \int_{\Omega} (|s_1^{n+1}| + |s_2^{n+1}| + (1 + 2e^{(\gamma-\nu)t_{n+1}} M)) s_1^{n+1} \\
\leq C_1 \int_{\Omega} (e^{(\gamma-\nu)t_{n+1}} M s_1^{n+1} + (s_1^{n+1})^2 + (s_2^{n+1})^2),
\end{aligned}$$

and hence, after denoting $A(\mathbf{u}, \Delta t) = \frac{\|\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n}\|_{1,\infty,\Omega}}{2\Delta t}$, we can write the bounds

$$\frac{1}{\Delta t} \|s_1^{n+1}\|_{0,\Omega}^2 + (\nu - A(\mathbf{u}, \Delta t) - C_1) \|s_1^{n+1}\|_{0,\Omega}^2 + (\nu - C_1) \int_{\Omega} e^{(\gamma-\nu)t_{n+1}} M s_1^{n+1} - C_1 \|s_2^{n+1}\|_{0,\Omega}^2 \leq 0, \quad (2.35)$$

$$\frac{1}{\Delta t} \|s_2^{n+1}\|_{0,\Omega}^2 + (\nu - A(\mathbf{u}, \Delta t) - C_2) \|s_2^{n+1}\|_{0,\Omega}^2 + (\nu - C_2) \int_{\Omega} e^{(\gamma-\nu)t_{n+1}} M s_2^{n+1} - C_2 \|s_1^{n+1}\|_{0,\Omega}^2 \leq 0. \quad (2.36)$$

We then employ (2.35) and (2.36), which leads to

$$\begin{aligned} & \frac{1}{\Delta t} (\|s_1^{n+1}\|_{0,\Omega}^2 + \|s_2^{n+1}\|_{0,\Omega}^2) + (\nu - A(\mathbf{u}, \Delta t) - \max\{C_1, C_2\}) (\|s_1^{n+1}\|_{0,\Omega}^2 + \|s_2^{n+1}\|_{0,\Omega}^2) \\ & + (\nu - C_1) \int_{\Omega} e^{(\gamma-\nu)t_{n+1}} M s_1^{n+1} + (\nu - C_2) \int_{\Omega} e^{(\gamma-\nu)t_{n+1}} M s_2^{n+1} \leq 0, \end{aligned}$$

and if we choose $\nu \geq A(\mathbf{u}, \Delta t) + \max\{C_1, C_2\}$, then we conclude, from the expression above, that $s_1^{n+1} = s_2^{n+1} = 0$. This leads to a contradiction with $s_1^{n+1}, s_2^{n+1} > 0$, and hence $w_1^{n+1}, w_2^{n+1} \leq e^{(\gamma-\nu)t_{n+1}} M$. An appeal to the induction principle completes the rest of the proof. $\square$

Corollary 1. *The image of the self-map T is contained in $[H^1(\Omega)]^2$.*

Proof. We need to show that $(w_1^{n+1}, w_2^{n+1}) := T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in [H^1(\Omega)]^2$ for any $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) \in \mathcal{S}$. By taking $s_1 = w_1^{n+1}$ in (2.31) and employing (2.27) with the definition of $\mathcal{S}$, we immediately see that

$$\begin{aligned} \frac{1}{\Delta t} \|w_1^{n+1}\|_{0,\Omega}^2 + D_1^{\min} \int_{\Omega} |\nabla w_1^{n+1}|^2 &= \int_{\Omega} \frac{\operatorname{div}(\mathbf{u}^{s,n+1} - \mathbf{u}^{s,n})}{2\Delta t} (w_1^{n+1})^2 + \int_{\Omega} \frac{w_1^n w_1^{n+1}}{\Delta t} \\ &\quad - \nu \|w_1^{n+1}\|_{0,\Omega}^2 + \int_{\Omega} e^{-\nu t_{n+1}} f^{n+1} w_1^{n+1}. \end{aligned} \quad (2.37)$$

Using the boundedness of the terms appearing in the right-hand side of (2.37), we have

$$\|w_1^{n+1}\|_{1,\Omega} \leq \text{Constant},$$

and thus $w_1^{n+1} \in H^1(\Omega)$. Showing that $w_2^{n+1} \in H^1(\Omega)$ is analogous. $\square$

Lemma 2.4. *The map T is continuous.*

Proof. Let $(\hat{w}_{1,k}^{n+1}, \hat{w}_{2,k}^{n+1})_k \in \mathcal{S}$ be a sequence such that $(\hat{w}_{1,k}^{n+1}, \hat{w}_{2,k}^{n+1})_k \rightarrow (\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$ in $[L^2(\Omega)]^2$ as $k \rightarrow \infty$. From the definition of T we have that $(w_{1,k}^{n+1}, w_{2,k}^{n+1}) = T(\hat{w}_{1,k}^{n+1}, \hat{w}_{2,k}^{n+1})$.

Note that $T(\mathcal{S})$ is bounded in $[H^1(\Omega)]^2$ (cf. Corollary 1), and since $H^1(\Omega)$ is compactly embedded into $L^2(\Omega)$, then $T(\mathcal{S})$ is relatively compact in $[L^2(\Omega)]^2$. We then proceed to extract from $(\hat{w}_{1,k}^{n+1}, \hat{w}_{2,k}^{n+1})_k$ a subsequence $(\hat{w}_{1,k_j}^{n+1}, \hat{w}_{2,k_j}^{n+1})_{j \rightarrow \infty}$ which converges to $(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$ a.e. in Ω . Consequently, and owing to the continuity and boundedness of the function, we have that $r(\hat{w}_{1,k_j}^{n+1}, \hat{w}_{1,k_j}^{n+1})$ converges to $r(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$ in $[L^2(\Omega)]^2$. Moreover, since the subsequence $(w_{1,k_j}^{n+1}, w_{2,k_j}^{n+1})_{j \rightarrow \infty}$ is bounded in $[H^1(\Omega)]^2$, there exists a subsequence $(w_{1,(k_j)_q}^{n+1}, w_{2,(k_j)_q}^{n+1})_{q \rightarrow \infty}$ such that

$$(w_{1,(k_j)_q}^{n+1}, w_{2,(k_j)_q}^{n+1})_q \xrightarrow{q \rightarrow \infty} (w_1^{n+1}, w_2^{n+1}),$$

weakly in $[H^1(\Omega)]^2$, strongly in $[L^2(\Omega)]^2$, and a.e. in Ω . And after taking the limit $q \rightarrow \infty$ in (2.31)-(2.32), we can assert that the converging subsequence of $(w_{1,k}^{n+1}, w_{2,k}^{n+1})_k$ in $[L^2(\Omega)]^2$ has as a limit $(w_1^{n+1}, w_2^{n+1}) = T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$. Proceeding in a similar fashion, we can safely say that all convergent subsequences of $(w_{1,k}^{n+1}, w_{2,k}^{n+1})_k$ have a unique limit, which in turn implies that $T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1}) = (w_1^{n+1}, w_2^{n+1})$. Using the fact that the map $T(\mathcal{S})$ is relatively compact in $[L^2(\Omega)]^2$ and has a unique limit for every subsequence of $(w_{1,k}^{n+1}, w_{2,k}^{n+1})_k$, we can conclude that $(w_{1,k}^{n+1}, w_{2,k}^{n+1})_k$ converges to $T(\hat{w}_1^{n+1}, \hat{w}_2^{n+1})$ in $[L^2(\Omega)]^2$. $\square$

Corollary 2. *The map T is compact in $[L^2(\Omega)]^2$.*

Proof. It readily follows from the proof of Lemma 2.4. $\square$

In view of the above results, we are in position to state the following existence theorem.

Theorem 2.1. *The semi-discrete formulation (2.2)-(2.6) for problem (1.9) possesses at least one solution.*

2.5. Uniqueness of weak solutions

Theorem 2.2. *The semi-discrete weak formulation (2.2)-(2.6) of the coupled problem (1.9) has a unique solution.*

Proof. We follow the strategy adopted in [5] and define two solutions $(\mathbf{u}_1^{s,n+1}, p_1^{f,n+1}, \psi_1^{n+1}, w_1^{1,n+1}, w_2^{1,n+1})$ and $(\mathbf{u}_2^{s,n+1}, p_2^{f,n+1}, \psi_2^{n+1}, w_1^{2,n+1}, w_2^{2,n+1})$ associated with initial data $\mathbf{b}_1^{n+1}, \ell_1^{n+1}, \mathbf{u}_1^{s,0}, p_1^{f,0}, \psi_1^0, w_1^1, w_2^1$, and $\mathbf{b}_2^{n+1}, \ell_2^{n+1}, \mathbf{u}_2^{s,0}, p_2^{f,0}, \psi_2^0, w_1^2, w_2^2$, respectively. We then introduce

$$\begin{aligned}\mathcal{U}^{n+1} &= \mathbf{u}_1^{s,n+1} - \mathbf{u}_2^{s,n+1}, \quad \mathcal{P}^{n+1} = p_1^{f,n+1} - p_2^{f,n+1}, \quad \chi^{n+1} = \psi_1^{n+1} - \psi_2^{n+1}, \\ \mathcal{W}_1^{n+1} &= w_1^{1,n+1} - w_1^{2,n+1}, \quad \mathcal{W}_2^{n+1} = w_2^{1,n+1} - w_2^{2,n+1},\end{aligned}$$

and then it follows from (2.2)-(2.4) that

$$\begin{aligned}2\mu \int_{\Omega} \varepsilon(\mathcal{U}^{n+1}) : \varepsilon(\mathbf{v}^s) - \int_{\Omega} \chi^{n+1} \operatorname{div} \mathbf{v}^s - \rho \int_{\Omega} ((\mathbf{b}_1^{n+1} - \mathbf{b}_2^{n+1}) \cdot \mathbf{v}^s \\ - \tau \int_{\Omega} (r_1^{n+1} - r_2^{n+1})(k \otimes k) : \varepsilon(\mathbf{v}^s) = 0, \\ \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \int_{\Omega} \delta_t \mathcal{P}^{n+1} q^f + \int_{\Omega} \frac{\kappa}{\eta} \nabla \mathcal{P}^{n+1} \cdot \nabla q^f - \frac{\alpha}{\lambda} \int_{\Omega} q^f \delta_t \chi^{n+1} - \int_{\Omega} (\ell_1^{n+1} - \ell_2^{n+1}) q^f = 0, \\ - \int_{\Omega} \phi \operatorname{div} \mathcal{U}^{n+1} + \frac{\alpha}{\lambda} \int_{\Omega} \mathcal{P}^{n+1} \phi - \frac{1}{\lambda} \int_{\Omega} \chi^{n+1} \phi = 0,\end{aligned}$$

for all $\mathbf{v}^s \in \mathbf{V}$, all $q^f \in Q$, and all $\phi \in Z$. Similarly as in the proof of Lemma 2.1 we employ $\delta_t \mathcal{U}^{n+1}$, $\mathcal{P}^{n+1}$, $\delta_t \chi^{n+1}$ as test functions, together with (2.9) to arrive at

$$\begin{aligned}\|\mathcal{U}^{n+1}\|_{1,\Omega}^2 + c_0 \|\mathcal{P}^{n+1}\|_{0,\Omega}^2 + \|\chi^{n+1}\|_{0,\Omega}^2 + \|\mathcal{P}\|_{L^2(H^1(\Omega))}^2 \leq C \Big(\|\mathcal{U}^0\|_{1,\Omega}^2 + \|\mathcal{P}^0\|_{0,\Omega}^2 + \|\chi^0\|_{0,\Omega}^2 \\ + \sum_{m=0}^n \|\mathbf{b}_1^{m+1} - \mathbf{b}_2^{m+1}\|_{0,\Omega}^2 + \|\ell_1 - \ell_2\|_{L^2(\Omega)}^2 + \sum_{m=0}^n (\|\mathcal{P}^{m+1}\|_{0,\Omega}^2 + \|\chi^{m+1}\|_{0,\Omega}^2 \\ + \|\mathcal{W}_1^{m+1}\|_{0,\Omega}^2 + \|\mathcal{W}_2^{m+1}\|_{0,\Omega}^2) \Big). \quad (2.38)\end{aligned}$$

In turn, for the ADR problem, we can get from (2.5) and (2.6) with test functions $\mathcal{W}_1^{n+1}$ and $\mathcal{W}_2^{n+1}$, respectively, to get the relations

$$\begin{aligned}\frac{1}{2} (\delta_t \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t \mathcal{W}_1^{n+1}\|_{0,\Omega}^2) + D_1^{\min} |\mathcal{W}_1^{n+1}|_{1,\Omega}^2 \\ \leq \int_{\Omega} (f_1^{n+1} - f_2^{n+1}) \mathcal{W}_1^{n+1} - \int_{\Omega} (\delta_t \mathbf{u}_1^{s,n+1} \cdot \nabla \mathcal{W}_1^{n+1} + \delta_t \mathcal{U}^{n+1} \cdot \nabla w_1^{2,n+1}) \mathcal{W}_1^{n+1}, \\ \frac{1}{2} (\delta_t \|\mathcal{W}_2^{n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t \mathcal{W}_2^{n+1}\|_{0,\Omega}^2) + D_2^{\min} |\mathcal{W}_2^{n+1}|_{1,\Omega}^2 \\ \leq \int_{\Omega} (g_1^{n+1} - g_2^{n+1}) \mathcal{W}_2^{n+1} - \int_{\Omega} (\delta_t \mathbf{u}_1^{s,n+1} \cdot \nabla \mathcal{W}_2^{n+1} + \delta_t \mathcal{U}^{n+1} \cdot \nabla w_2^{2,n+1}) \mathcal{W}_2^{n+1}.\end{aligned}$$

As in Lemma 2.2, we integrate by parts and assume that $w_i^{j,n+1} \in W^{1,\infty}(\Omega)$, $i, j = 1, 2$, which yields

$$\begin{aligned}\frac{1}{2} (\delta_t \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t \mathcal{W}_1^{n+1}\|_{0,\Omega}^2) + D_1^{\min} |\mathcal{W}_1^{n+1}|_{1,\Omega}^2 \leq \|f_1^{n+1} - f_2^{n+1}\|_{0,\Omega} \|\mathcal{W}_1^{n+1}\|_{0,\Omega} \\ + \frac{1}{2} \|\delta_t \mathbf{u}_1^{s,n+1}\|_{1,\infty,\Omega} \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \|w_1^{2,n+1}\|_{1,\infty,\Omega} \|\delta_t \mathcal{U}^{n+1}\|_{0,\Omega} \|\mathcal{W}_1^{n+1}\|_{0,\Omega},\end{aligned}$$

and applying Cauchy Schwarz and Young inequalities together with (2.8), we get the bound

$$\begin{aligned} \frac{1}{2} (\delta_t \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \Delta t \|\delta_t \mathcal{W}_1^{n+1}\|_{0,\Omega}^2) + D_1^{\min} |\mathcal{W}_1^{n+1}|_{1,\Omega}^2 &\leq C \left(\|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \|\mathcal{W}_2^{n+1}\|_{0,\Omega}^2 \right. \\ &\quad \left. + \frac{1}{2\Delta t} \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2} \|\delta_t \mathcal{U}^{n+1}\|_{0,\Omega}^2 + \frac{1}{2\Delta t} \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 \right). \end{aligned}$$

Multiplying by Δt and taking summation over n , we deduce that

$$\begin{aligned} \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \Delta t^2 \sum_{m=0}^n \|\delta_t \mathcal{W}_1^{m+1}\|_{0,\Omega}^2 + D_1^{\min} \Delta t \sum_{m=0}^n |\mathcal{W}_1^{m+1}|_{1,\Omega}^2 &\leq C \left(\|\mathcal{W}_1^0\|_{0,\Omega}^2 + \|\mathcal{U}^0\|_{0,\Omega}^2 \right. \\ &\quad \left. + \sum_{m=0}^n ((1 + \Delta t) \|\mathcal{W}_1^{m+1}\|_{0,\Omega}^2 + \|\mathcal{W}_2^{m+1}\|_{0,\Omega}^2) + \sum_{m=0}^n \|\mathcal{U}^{m+1}\|_{0,\Omega}^2 \right), \end{aligned}$$

and then, proceeding in a similar way as above, we obtain the same bounds for $\mathcal{W}_2^{n+1}$. Therefore, we have

$$\begin{aligned} \|\mathcal{W}_1^{n+1}\|_{0,\Omega}^2 + \|\mathcal{W}_2^{n+1}\|_{0,\Omega}^2 + \Delta t^2 \sum_{m=0}^n (\|\delta_t \mathcal{W}_1^{m+1}\|_{0,\Omega}^2 + \|\delta_t \mathcal{W}_2^{m+1}\|_{0,\Omega}^2) + D^{\min} \Delta t \sum_{m=0}^n (|\mathcal{W}_1^{m+1}|_{1,\Omega}^2 \\ + |\mathcal{W}_2^{m+1}|_{1,\Omega}^2) \leq C \left(\|\mathcal{W}_1^0\|_{0,\Omega}^2 + \|\mathcal{W}_2^0\|_{0,\Omega}^2 + \|\mathcal{U}^0\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathcal{U}^{m+1}\|_{0,\Omega}^2 \right. \\ \left. + (1 + \Delta t) \sum_{m=0}^n (\|\mathcal{W}_1^{m+1}\|_{0,\Omega}^2 + \|\mathcal{W}_2^{m+1}\|_{0,\Omega}^2) \right). \quad (2.39) \end{aligned}$$

The desired estimate is established by combining (2.38) and (2.39), and Gronwall's lemma

$$\begin{aligned} \|\mathcal{U}^{n+1}\|_{1,\Omega} + \|\mathcal{P}^{n+1}\|_{0,\Omega} + \|\chi^{n+1}\|_{0,\Omega} + \|\mathcal{W}_1^{n+1}\|_{0,\Omega} + \|\mathcal{W}_2^{n+1}\|_{0,\Omega} + \|\mathcal{P}\|_{l^2(H^1(\Omega))} + \|\nabla \mathcal{W}_1\|_{l^2(L^2(\Omega))} \\ + \|\nabla \mathcal{W}_2\|_{l^2(L^2(\Omega))} \leq C \left(\|\mathcal{U}^0\|_{1,\Omega} + \|\mathcal{P}^0\|_{0,\Omega} + \|\chi^0\|_{0,\Omega} + \|\mathcal{W}_1^0\|_{0,\Omega} + \|\mathcal{W}_2^0\|_{0,\Omega} \right. \\ \left. + \sum_{m=0}^n \|\mathbf{b}_1^{m+1} - \mathbf{b}_2^{m+1}\|_{0,\Omega} + \|\ell_1 - \ell_2\|_{l^2(L^2(\Omega))} \right), \end{aligned}$$

from which, we can ensure the existence of at most one weak solution to the system (2.2)-(2.6). $\square$

2.6. Continuous dependence on data

Lemma 2.5. *The solution $(\mathbf{u}^{s,n+1}, p^{f,n+1}, \psi^{n+1}, w_1^{n+1}, w_2^{n+1}) \in \mathbf{V} \times Q \times Z \times H^1(\Omega) \times H^1(\Omega)$ of problem (2.2)-(2.6) satisfies*

$$\begin{aligned} \|\mathbf{u}^{s,n+1}\|_{1,\Omega} + \sqrt{c_0} \|p^{f,n+1}\|_{0,\Omega} + \|\psi^{n+1}\|_{0,\Omega} + \|p^f\|_{\ell^2(H^1(\Omega))} + \|w_1^{n+1}\|_{0,\Omega} + \|w_2^{n+1}\|_{0,\Omega} \\ \leq C \sqrt{\exp} \left\{ n\Delta t + \|\mathbf{u}^{s,0}\|_{1,\Omega} + \|p^{f,0}\|_{0,\Omega} + \|\psi^0\|_{0,\Omega} + \|w_1^0\|_{0,\Omega} + \|w_2^0\|_{0,\Omega} + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega} + \|\ell\|_{\ell^2(L^2(\Omega))} \right\}. \end{aligned}$$

where $C > 0$ is a constant independent of Δt and λ .

Proof. We focus first on the Biot system. Proceeding as in the proof of Lemma 2.1, we take $\mathbf{v}^s = \delta_t \mathbf{u}^{s,n+1}$, $q^f = p^{f,n+1}$ and $\phi = \delta_t \psi^{n+1}$ in (2.2), (2.3) and (2.4), respectively, to obtain

$$\|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + c_0 \|p^{f,n+1}\|_{0,\Omega}^2 + \|\psi^{n+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2$$

$$\begin{aligned}
&\leq C_1 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \|p^{f,0}\|_{0,\Omega}^2 + \|\psi^0\|_{0,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 \right. \\
&\quad \left. + \sum_{m=0}^n \|r^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \right\} + C_2 \left\{ \|r^{n+1}\|_{0,\Omega}^2 + \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 \right\}.
\end{aligned} \tag{2.40}$$

In turn, for the ADR problem, we proceed as in the proof of Lemma 2.2, taking $s_1 = w_1^{n+1}$ and $s_2 = w_2^{n+1}$ in (2.5) and (2.6), respectively, to get

$$\begin{aligned}
&\|w_1^{n+1}\|_{0,\Omega}^2 + \|w_2^{n+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n (\|\nabla w_1^{m+1}\|_{0,\Omega}^2 + \|\nabla w_2^{m+1}\|_{0,\Omega}^2) \\
&\leq C_3 \left\{ n + \|w_1^0\|_{0,\Omega}^2 + \|w_2^0\|_{0,\Omega}^2 + \sum_{m=0}^n \left(\|w_1^{m+1}\|_{0,\Omega}^2 + \|w_2^{m+1}\|_{0,\Omega}^2 \right) \right\}.
\end{aligned} \tag{2.41}$$

Combining (2.40) and (2.41), we obtain a preliminar stability bound for the coupled system (2.2)-(2.6)

$$\begin{aligned}
&\|\mathbf{u}^{s,n+1}\|_{1,\Omega}^2 + c_0 \|p^{f,n+1}\|_{0,\Omega}^2 + \|\psi^{n+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|p^{f,m+1}\|_{1,\Omega}^2 + \|w_1^{n+1}\|_{0,\Omega}^2 + \|w_2^{n+1}\|_{0,\Omega}^2 \\
&\leq C_1 \left\{ \|\mathbf{u}^{s,0}\|_{1,\Omega}^2 + \|p^{f,0}\|_{0,\Omega}^2 + \|\psi^0\|_{0,\Omega}^2 + \sum_{m=0}^n \|\psi^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|p^{f,m+1}\|_{0,\Omega}^2 \right. \\
&\quad \left. + \sum_{m=0}^n \|r^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 + \Delta t \sum_{m=0}^n \|\ell^{n+1}\|_{0,\Omega}^2 \right\} + C_2 \left\{ \|r^{n+1}\|_{0,\Omega}^2 + \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 \right\} \\
&\quad + C_3 \left\{ n\Delta t + \|w_1^0\|_{0,\Omega}^2 + \|w_2^0\|_{0,\Omega}^2 + \sum_{m=0}^n \left(\|w_1^{m+1}\|_{0,\Omega}^2 + \|w_2^{m+1}\|_{0,\Omega}^2 \right) \right\},
\end{aligned}$$

and therefore, recalling the bound for r given in Section 2.2, and applying Gronwall's inequality to the resulting estimate, we obtain the desired result. $\square$

3. Mixed-primal Galerkin method

3.1. Fully discrete formulation

Let us consider a family $\{\mathcal{T}_h\}_{h>0}$ of shape-regular, quasi-uniform partitions of the spatial domain $\bar{\Omega}$ into affine elements (triangles in 2D or tetrahedra in 3D) E of diameter h_E , where $h = \max\{h_E : E \in \mathcal{T}_h\}$ denotes the mesh size. Finite-dimensional subspaces of the functional spaces employed in Section 2 will be defined in the following manner

$$\begin{aligned}
\mathbf{V}_h &:= \{\mathbf{v}_h \in \mathbf{C}(\bar{\Omega}) : \mathbf{v}_h|_E \in [\mathbb{P}_1(E) \oplus \text{span}\{b_E\}]^d \forall E \in \mathcal{T}_h, \text{ and } \mathbf{v}_h|_\Gamma = \mathbf{0}\}, \\
Q_h &:= \{q_h \in C(\bar{\Omega}) : q_h|_E \in \mathbb{P}_1(E) \forall E \in \mathcal{T}_h, \text{ and } q_h|_\Sigma = 0\}, \\
Z_h &:= \{\phi_h \in L^2(\Omega) : \phi_h|_E \in \mathbb{P}_1(E) \forall E \in \mathcal{T}_h\}, \quad W_h := \{w_h \in C(\bar{\Omega}) : w_h|_E \in \mathbb{P}_1(E) \forall E \in \mathcal{T}_h\},
\end{aligned} \tag{3.1}$$

where $\mathbb{P}_k(E)$ denotes the space of polynomials of degree less than or equal than k defined locally over $E \in \mathcal{T}_h$, and $b_E := \varphi_1 \varphi_2 \varphi_3$ is a $\mathbb{P}_3$ bubble function in E , and $\varphi_1, \varphi_2, \varphi_3$ are the barycentric coordinates of E . Let us recall that the pair $(\mathbf{V}_h, Z_h)$ (known as the MINI element) is inf-sup stable (see e.g. [8]).

Considering reaction and coupling terms f, g, r discretised implicitly, the fully discrete scheme associated with (2.1) is defined as: From initial data $\mathbf{u}^{s,0}, p^{f,0}, \psi^0, w_1^0, w_2^0$ (which will be projections of the

continuous initial conditions of each field) and for $n = 1, \dots$, find $\mathbf{u}_h^{s,n+1} \in \mathbf{V}_h, p_h^{f,n+1} \in Q_h, \psi_h^{n+1} \in Z_h, w_{1,h}^{n+1} \in W_h, w_{2,h}^{n+1} \in W_h$ such that

$$a_1(\mathbf{u}_h^{s,n+1}, \mathbf{v}_h^s) + b_1(\mathbf{v}_h^s, \psi_h^{n+1}) = F_{r_h^{n+1}}(\mathbf{v}_h^s) \quad \forall \mathbf{v}_h^s \in \mathbf{V}_h, \quad (3.2)$$

$$\tilde{a}_2(p_h^{f,n+1}, q_h^f) + a_2(p_h^{f,n+1}, q_h^f) - \tilde{b}_2(q_h^f, \psi_h^{n+1}) = G_{\ell^{n+1}}(q_h^f) \quad \forall q_h^f \in Q_h, \quad (3.3)$$

$$b_1(\mathbf{u}_h^{s,n+1}, \phi_h) + b_2(p_h^{f,n+1}, \phi_h) - a_3(\psi_h^{n+1}, \phi_h) = 0 \quad \forall \phi_h \in Z_h, \quad (3.4)$$

$$\tilde{a}_4(w_{1,h}^{n+1}, s_{1,h}) + a_4(w_{1,h}^{n+1}, s_{1,h}) + c(w_{1,h}^{n+1}, s_{1,h}, \mathbf{u}_h^{s,n+1}) = J_{f_h^{n+1}}(s_{1,h}) \quad \forall s_{1,h} \in W_h, \quad (3.5)$$

$$\tilde{a}_5(w_{2,h}^{n+1}, s_{2,h}) + a_5(w_{2,h}^{n+1}, s_{2,h}) + c(w_{2,h}^{n+1}, s_{2,h}, \mathbf{u}_h^{s,n+1}) = J_{g_h^{n+1}}(s_{2,h}) \quad \forall s_{2,h} \in W_h. \quad (3.6)$$

3.2. Stability of the discrete solutions

Lemma 3.1. Assume that $(\mathbf{u}_h^{s,n+1}, p_h^{f,n+1}, \psi_h^{n+1}, w_{1,h}^{n+1}, w_{2,h}^{n+1}) \in \mathbf{V}_h \times Q_h \times Z_h \times W_h \times W_h$ is solution of problem (3.2)-(3.6). Then, there exists $C > 0$ independent of λ, h , and Δt , such that

$$\begin{aligned} & \|\mathbf{u}_h^{s,n+1}\|_{1,\Omega} + \sqrt{c_0} \|p_h^{f,n+1}\|_{0,\Omega} + \|\psi_h^{n+1}\|_{0,\Omega} + \|p_h^f\|_{\ell^2(H^1(\Omega))} + \|w_{1,h}^{n+1}\|_{0,\Omega} + \|w_{2,h}^{n+1}\|_{0,\Omega} \\ & \leq C \sqrt{\exp} \left\{ n \Delta t + \|\mathbf{u}_h^{s,0}\|_{1,\Omega} + \|p_h^{f,0}\|_{0,\Omega} + \|\psi_h^0\|_{0,\Omega} + \|w_{1,h}^0\|_{0,\Omega} \right. \\ & \quad \left. + \|w_{2,h}^0\|_{0,\Omega} + \sum_{m=0}^{n+1} \|\mathbf{b}^{m+1}\|_{0,\Omega} + \|\ell\|_{\ell^2(L^2(\Omega))} \right\}. \end{aligned} \quad (3.7)$$

Proof. We proceed similarly to the proof of Lemmas 2.1 and 2.2. We focus first on the stability of (3.2)-(3.4). Taking $\mathbf{v}_h^s = \delta_t \mathbf{u}_h^{s,n+1}$ in (3.2), using Cauchy-Schwarz inequality, applying Young's inequality with constants chosen conveniently, and then, summing over n and multiplying by Δt , we readily get

$$\begin{aligned} & \mu C_{k,1} \|\mathbf{u}_h^{s,n+1}\|_{1,\Omega}^2 + \frac{\mu C_{k,1} \Delta t^2}{4} \sum_{m=0}^n \|\delta_t \mathbf{u}_h^{s,m+1}\|_{1,\Omega}^2 \\ & \leq C_1 \left\{ \|\mathbf{u}_h^{s,0}\|_{1,\Omega}^2 + \sum_{m=0}^n \|\psi_h^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|r_h^{m+1}\|_{0,\Omega}^2 + \sum_{m=0}^n \|\mathbf{b}^{m+1}\|_{0,\Omega}^2 \right\}, \end{aligned} \quad (3.8)$$

where C_1 is a constant depending on $\mu, C_{k,1}, C_{k,2}, \rho$, and τ . Now, in equations (3.3) and (3.4), we take $q_h = p_h^{f,n+1}$ and $\phi_h = \delta_t \psi_h^{n+1}$, respectively, to obtain

$$\begin{aligned} & \frac{1}{2\lambda} \delta_t \|\psi_h^{n+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2\lambda} \|\delta_t \psi_h^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\delta_t \|p_h^{f,n+1}\|_{0,\Omega} + \Delta t \|\delta_t p_h^{f,n+1}\|_{0,\Omega} \right) + \frac{\kappa_1}{\eta} |p_h^{f,n+1}|_{1,\Omega}^2 \\ & \leq \frac{2\alpha}{\lambda} \|p_h^{f,n+1}\|_{0,\Omega} \|\delta_t \psi_h^{n+1}\|_{0,\Omega} + \|\ell^{n+1}\|_{0,\Omega} \|p_h^{f,n+1}\|_{0,\Omega} - \int_{\Omega} \delta_t \psi_h^{n+1} \operatorname{div} \mathbf{u}_h^{s,n+1}, \end{aligned} \quad (3.9)$$

Thus, applying Young's inequality to the first and second term, and summation by parts to the last term, on the right-hand side of (3.9), we obtain

$$\begin{aligned} & \frac{1}{2\lambda} \|\psi_h^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \left(\|p_h^{f,n+1}\|_{0,\Omega}^2 + \Delta t^2 \sum_{m=0}^n \|\delta_t p_h^{f,m+1}\|_{0,\Omega}^2 \right) + \frac{\kappa_1 c_p \Delta t}{2\eta} \sum_{m=0}^n \|p_h^{f,m+1}\|_{1,\Omega}^2 \\ & \leq \frac{1}{2\lambda} \|\psi_h^0\|_{0,\Omega}^2 + \frac{1}{2} \left(c_0 + \frac{\alpha^2}{\lambda} \right) \|p_h^{f,0}\|_{0,\Omega}^2 + \frac{2\alpha^2}{\lambda} \sum_{m=0}^n \|p_h^{f,m+1}\|_{0,\Omega}^2 + \frac{\eta \Delta t}{2\kappa_1 c_p} \sum_{m=0}^n \|\ell^{m+1}\|_{0,\Omega}^2 \end{aligned} \quad (3.10)$$

$$+ \frac{1}{\mu C_{k,1}} \|\psi_h^{n+1}\|_{0,\Omega}^2 + \frac{\mu C_{k,1}}{2} \|\mathbf{u}_h^{s,n+1}\|_{1,\Omega}^2 + \frac{2}{\mu C_{k,1}} \sum_{m=0}^{n-1} \|\psi_h^{m+1}\|_{0,\Omega}^2 + \frac{\mu C_{k,1} \Delta t^2}{4} \sum_{m=0}^{n-1} \|\delta_t \mathbf{u}_h^{m+1}\|_{1,\Omega}^2.$$

On the other hand, as in Lemma 2.1 we target an estimate independent of λ . For that reason we use the discrete version of the inf-sup condition (2.12), which is satisfied by the finite element family (3.1) [14, 8]. Thus, taking $\phi_h = \psi_h^{n+1}$, using (3.2) and the continuity of a_1 , we obtain

$$\begin{aligned} \hat{\beta} \|\psi_h^{n+1}\|_{0,\Omega} &\leq \sup_{\mathbf{v}_h^s \in \mathbf{V}_h} \frac{b_1(\mathbf{v}_h^s, \psi_h^{n+1})}{\|\mathbf{v}_h^s\|_{1,\Omega}} = \sup_{\mathbf{v}_h^s \in \mathbf{V}_h} \frac{-a_1(\mathbf{u}_h^{s,n+1}, \mathbf{v}_h^s) + F_{r_h^{n+1}}(\mathbf{v}_h^s)}{\|\mathbf{v}_h^s\|_{1,\Omega}} \\ &\leq 2\mu C_{k,2} \|\varepsilon(\mathbf{u}_h^{n+1})\|_{0,\Omega} + \sqrt{C_{k,2}\tau} \|r_h^{n+1}\|_{0,\Omega} + \rho \|\mathbf{b}^{n+1}\|_{0,\Omega}, \end{aligned}$$

or what is the same

$$\|\psi_h^{n+1}\|_{0,\Omega}^2 \leq C_2 \left\{ \|\mathbf{u}_h^{s,n+1}\|_{1,\Omega}^2 + \|r_h^{n+1}\|_{0,\Omega}^2 + \|\mathbf{b}^{n+1}\|_{0,\Omega}^2 \right\}, \quad (3.11)$$

where C_2 is a constant depending on $C_{k,1}$, $C_{k,2}$, μ , τ , ρ and the discrete inf-sup constant $\hat{\beta}$. In turn, for the ADR problem (3.5)-(3.6), by taking $s_{1,h} = w_{1,h}^{n+1}$ in (3.5), we get

$$\int_{\Omega} \delta_t w_{1,h}^{n+1} w_{1,h}^{n+1} + \int_{\Omega} D_1(\mathbf{x}) \nabla w_{1,h}^{n+1} \cdot \nabla w_{1,h}^{n+1} + \int_{\Omega} (\delta_t \mathbf{u}_h^{s,n+1} \cdot \nabla w_{1,h}^{n+1}) w_{1,h}^{n+1} = \int_{\Omega} f_h^{n+1} w_{1,h}^{n+1},$$

and then, applying (2.27) and Cauchy-Schwarz inequality, we deduce the estimate

$$\begin{aligned} \frac{1}{2} \delta_t \|w_{1,h}^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t \|\delta_t w_{1,h}^{n+1}\|_{0,\Omega}^2 + D_1^{\min} \|\nabla w_{1,h}^{n+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|\delta_t \mathbf{u}_h^{s,n+1}\|_{1,\infty,\Omega} \|w_{1,h}^{n+1}\|_{0,\Omega}^2 + \|f_h^{n+1}\|_{0,\Omega} \|w_{1,h}^{n+1}\|_{0,\Omega}. \end{aligned}$$

Since Ω is a bounded domain and the elements of $\mathbf{V}_h$ are piecewise polynomials, we know that $\|\mathbf{u}_h^{s,n+1} - \mathbf{u}_h^{s,n}\|_{1,\infty,\Omega} < +\infty$ for each $\mathbf{u}_h^{n+1}, \mathbf{u}_h^n \in \mathbf{V}_h$ (see e.g. [11]), and then, without loss of generality, we may assume that $\|\mathbf{u}_h^{s,n+1} - \mathbf{u}_h^{s,n}\|_{1,\infty,\Omega} \leq M_1$ for some $M_1 \in \mathbb{R}$. Thus, applying Young's inequality, summing over n and multiplying by Δt , we obtain the following result

$$\begin{aligned} \frac{1}{2} \|w_{1,h}^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t^2 \sum_{m=0}^n \|\delta_t w_{1,h}^{m+1}\|_{0,\Omega}^2 + D_1^{\min} \Delta t \sum_{m=0}^n \|\nabla w_{1,h}^{m+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|w_{1,h}^0\|_{0,\Omega}^2 + \frac{1}{2} (M_1 + \Delta t) \sum_{m=0}^n \|w_{1,h}^{m+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2} \sum_{m=0}^n \|f_h^{m+1}\|_{0,\Omega}^2. \end{aligned} \quad (3.12)$$

Moreover, we realise that an estimate for $\|w_{2,h}^{n+1}\|_{0,\Omega}$ stays exactly as above, which is

$$\begin{aligned} \frac{1}{2} \|w_{2,h}^{n+1}\|_{0,\Omega}^2 + \frac{1}{2} \Delta t^2 \sum_{m=0}^n \|\delta_t w_{2,h}^{m+1}\|_{0,\Omega}^2 + D_2^{\min} \Delta t \sum_{m=0}^n \|\nabla w_{2,h}^{m+1}\|_{0,\Omega}^2 \\ \leq \frac{1}{2} \|w_{2,h}^0\|_{0,\Omega}^2 + \frac{1}{2} (M_1 + \Delta t) \sum_{m=0}^n \|w_{2,h}^{m+1}\|_{0,\Omega}^2 + \frac{\Delta t}{2} \sum_{m=0}^n \|g_h^{m+1}\|_{0,\Omega}^2. \end{aligned} \quad (3.13)$$

Finally, the stability stated in (3.7), follows from the growth condition on f_h and g_h , adding (3.8), (3.10), (3.11), (3.12) and (3.13), recalling the bound for r , and applying the discrete Gronwall's inequality. $\square$

Remark 3.1. The solvability analysis of (3.2)-(3.6) can be established similarly to the continuous case. More precisely, as in Section 2.3 we need to define a fixed-point operator, whose well-definiteness will depend upon the




Figure 4.1: Test 1. Convergence of the discretisation for the coupled poro-mechano-chemical problem. Error decay in space (left) and error history in time (right, where errors are computed from (4.2)).

solvability of each uncoupled problem. For the discrete poroelasticity system we can adapt the analysis from [23, Section 3], whereas for the approximate ADR equations we can apply classical techniques for discrete quasi-linear problems [24]. Next, we need to prove the continuity of the operator going from $[W_h]^2$ into itself, which follows as a consequence of the estimate (3.7) in combination with the ideas employed in [5, Section 5.3]. Finally, the result follows from an application of the well-known Brouwer fixed-point theorem.

4. Numerical verification of spatio-temporal convergence

We have not derived theoretically error bounds, but proceed in this Section to examine numerically the rates of convergence of the mixed-primal scheme. Let us consider $\Omega = (0, 1)^2$ with $\Gamma = \{\mathbf{x} : x_1 = 0 \text{ or } x_2 = 0\}$ (the bottom and left edges of the boundary) and $\Sigma = \{\mathbf{x} : x_1 = 1 \text{ or } x_2 = 1\}$ (top and right sides of the square domain). Following [17], we define closed-form solutions to the coupled poro-mechano-chemical system (1.9) as

$$\begin{aligned} \mathbf{u}^s &= u_\infty \frac{t^2}{2} \begin{pmatrix} \sin(\pi x_1) \cos(\pi x_2) + \frac{x_1^2}{\lambda} \\ -\cos(\pi x_1) \sin(\pi x_2) + \frac{x_2^2}{\lambda} \end{pmatrix}, \quad p^f = t(x_1^3 - x_2^4), \quad \psi = p^f - \lambda \operatorname{div} \mathbf{u}^s, \\ w_1 &= t[\exp(x_1) + \cos(\pi x_1) \cos(\pi x_2)], \quad w_2 = t[\exp(-x_2) + \sin(\pi x_1) \sin(\pi x_2)], \end{aligned} \quad (4.1)$$

and we use these smooth functions to construct expressions for the body force $\mathbf{b}(\mathbf{x}, t)$, the fluid source $\ell(\mathbf{x}, t)$, additional mass sources $S_1(\mathbf{x}, t), S_2(\mathbf{x}, t)$ for (1.5)-(1.6); a non-homogeneous displacement and non-homogeneous fluid normal flux on Γ , as well as non-homogeneous Dirichlet boundary pressure and non-homogeneous traction defined on Σ . The model parameters take the values: $u_\infty = \alpha = \gamma = 0.1$, $c_0 = \eta = 10^{-3}$, $\kappa = 10^{-4}$, $D_1 = 0.05$, $D_2 = \rho$, $\beta_1 = 170$, $\beta_2 = 0.1305$, $\beta_3 = 0.7695$, $E = 3 \cdot 10^4$, $\nu = 0.495$ (giving the Lamé constants $\mu = 10033.444$, $\lambda = 993311.037$), and $\tau = 10^5$. For this example we simply take the function that modulates the active stress in (1.8) as $r = w_1 + w_2$ and use $\mathbf{k} = (1, 0)^T$.

To assess the spatial accuracy of the discretisation defined by the finite element spaces specified in (3.1), we construct a sequence of seven uniformly refined meshes and compute individual approximate errors $e(\cdot)$ for each field in their natural spatial norm at the final time $t_{\text{final}} = 0.04$, and the time-stepping scheme (backward Euler and implicit centred differences for first and second order time derivatives, respectively) approximates the polynomial dependence on time in (4.1) exactly. The system is solved by the GMRES Krylov solver with incomplete LU factorisation (ILUT) preconditioning. The stopping criterion on the nonlinear iterations is based on a weighted residual norm dropping below the fixed tolerance of $1 \cdot 10^{-5}$. Moreover, a small fixed time step $\Delta t = 0.01$ is used for all mesh refinements. An

average number of three Newton iterations are needed in all levels to reach convergence. The results are laid out in Figure 4.1 (left) where we observe an optimal error decay of $O(h)$ for all field variables. We also see that the total error is dominated by the total pressure (which is large as these errors are not normalised and since the regime is nearly incompressible), but the convergence rates remain optimal with respect to the expected accuracy given by the interpolation properties of the finite element spaces.

The convergence associated with the time discretisation can be more conveniently assessed considering a different set of closed-form solutions defined on a fixed mesh with 4000 elements

$$\mathbf{u}^s = u_\infty \sin(t) \begin{pmatrix} \frac{x_1^2}{2\lambda} + \frac{x_2^2}{2\lambda} \\ x_1^2 + \frac{x_2^2}{2\lambda} \end{pmatrix}, \quad p^f = \sin(t)(x_1^2 + x_1x_2), \quad w_1 = \sin(t)(x_1^2 - x_2^2), \quad w_2 = \sin(t)(x_1^2 + x_2^2).$$

With the given spatial discretisation, the errors will contain only contributions from the time approximation. We consider now the time interval $(0, 1]$ and choose six time-step uniform refinements $\Delta t \in \{0.5, 0.25, \dots\}$ that we use to compute numerical solutions and cumulative errors up to t_{final} , of a generic individual field s defined as

$$E(s) = \left(\Delta t \sum_{n=1}^N \|s_h^n - s(t^n)\|_{0,\Omega}^2 \right)^{\frac{1}{2}}. \quad (4.2)$$

Figure 4.1 (right) indicates that the errors in time are also of first order, $O(\Delta t)$.

5. Concluding remarks

In this paper we have analysed a model of advection-reaction-diffusion in poroelastic materials. The set of equations assumes the regime of small strains and the coupling mechanisms are primarily dependent on source functions of change of volume, and active stresses. We have derived the well-posedness of the problem stated in mixed-primal form, and we have proposed a suitable mixed finite element scheme. Our work extends the similar-in-spirit contribution [5] in that we are able to derive stability bounds that are robust with respect to the Lamé constants of the solid. Since the proofs do not rely entirely on the specific form of the reaction terms, the present formalism is quite general and could be applied to other systems with similar mathematical and physical structure, such as tumour development dynamics, long bones growth, or embryonic cell poromechanics.

As perspectives of this work, we aim at extending the analysis of Section 2 to the case of finite-strain poroelasticity following the work in [7], and to incorporate viscoelasticity. Further directions include the design of mixed and double-mixed formulations that would improve the accuracy of the method in producing stresses or other variables of applicative interest and also contributing to achieve mass conservation [17, 15], as well as mesh adaptive methods guided by a posteriori error indicators [2, 1].

References

- [1] E. AHMED, J.M. NORDBOTTEN, AND F.A. RADU, *Adaptive asynchronous time-stepping, stopping criteria, and a posteriori error estimates for fixed-stress iterative schemes for coupled poromechanics problems*. J. Comput. Appl. Math., 364 (2020) 112312.
- [2] E. AHMED, F.A. RADU, AND J.M. NORDBOTTEN, *Adaptive poromechanics computations based on a posteriori error estimates for fully mixed formulations of Biot's consolidation mode*. Comput. Methods Appl. Mech. Engrg., 347 (2019) 264–294.
- [3] M. ALVAREZ, G.N. GATICA AND R. RUIZ-BAIER, *An augmented mixed–primal finite element method for a coupled flow–transport problem*. ESAIM: Math. Model. Numer. Anal., 49(5) (2015), 1399–1427.

- [4] I. AMBARTSUMYAN, E. KHATTATOV, I. YOTOV, AND P. ZUNINO, *A Lagrange multiplier method for a Stokes-Biot fluid-poroelastic structure interaction model*. Numer. Math., 140(2) (2018) 513–553.
- [5] V. ANAYA, M. BENDAHMANE, D. MORA, AND R. RUIZ-BAIER, *On a vorticity-based formulation for reaction-diffusion-Brinkman systems*. Netw. Heterog. Media., 13(1) (2018) 69–94.
- [6] V. ANAYA, Z. DE WIJN, B. GOMEZ-VARGAS, D. MORA, AND R. RUIZ-BAIER, *Rotation-based mixed formulations for an elasticity-poroelasticity interface problem*. Submitted preprint (2019).
- [7] L. BERGER, R. BORDAS, D. KAY, AND S. TAVENER, *A stabilized finite element method for finite-strain three-field poroelasticity*. Comput. Mech., 60(1) (2017) 51–68.
- [8] D. BOFFI, F. BREZZI, AND M. FORTIN, *Mixed Finite Element Methods and Applications*, Vol. 44 Springer Series in Computational Mathematics. Springer, Heidelberg (2013).
- [9] M.K. BRUN, E. AHMED, J.M. NORDBOTTEN, AND F.A. RADU, *Well-posedness of the fully coupled quasi-static thermo-poroelastic equations with nonlinear convective transport*. J. Math. Anal. Appl., 471(1-2) (2019) 239–266.
- [10] M.K. BRUN, I. BERRE, J.M. NORDBOTTEN, AND F.A. RADU, *Upscaling of the coupling of hydromechanical and thermal processes in a quasi-static poroelastic medium*. Transp. Porous Media., 124(1) (2018) 137–158.
- [11] G.N. GATICA, B. GOMEZ-VARGAS AND R. RUIZ-BAIER, *Analysis and mixed-primal finite element discretisations for stress-assisted diffusion problems*. Comput. Methods Appl. Mech. Engrg., 337 (2018), 411–438.
- [12] G.N. GATICA, A. MARQUEZ AND S. MEDDAHI, *Analysis of the coupling of primal and dual-mixed finite element methods for a two-dimensional fluid-solid interaction problem*. SIAM J. Numer. Anal., 45(5) (2007) 2072–2097.
- [13] G.N. GATICA, R. OYARZÚA AND F.J. SAYAS, *Analysis of fully-mixed finite element methods for the Stokes-Darcy coupled problem*. Math. Comp., 80 (2011) 1911–1948.
- [14] V. GIRAULT, AND P.-A. RAVIART, *Finite Element Approximation of the Navier-Stokes Equation*, Lecture Notes in Math. 749, Springer-Verlag, Berlin, New York, 1979.
- [15] Q. HONG AND J. KRAUS, *Parameter-robust stability of classical three-field formulation of Biot’s consolidation model*. Electron. Trans. Numer. Anal., 48 (2018) 202–226.
- [16] G.W. JONES AND S.J. CHAPMAN, *Modeling growth in biological materials*. SIAM Rev., 54(1) (2012) 52–118.
- [17] S. KUMAR, R. OYARZÚA, R. RUIZ-BAIER, AND R. SANDILYA, *Conservative discontinuous finite volume and mixed schemes for a new four-field formulation in poroelasticity*. ESAIM: Math. Model. Numer. Anal., (2019) in press.
- [18] O.A. LADYŽENSKAJA, V. A. SOLONNIKOV, AND N.N. URAL’CEVA, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., (1968).
- [19] J.J. LEE, K.-A. MARDAL, AND R. WINTHER, *Parameter-robust discretization and preconditioning of Biot’s consolidation model*. SIAM J. Sci. Comp., 39 (2017) A1–A24.
- [20] J.J. LEE, E. PIERSANTI, K.-A. MARDAL, AND M. ROGNES, *A mixed finite element method for nearly incompressible multiple-network poroelasticity*. SIAM J. Sci. Comp., 41(2) (2019) A722–A747.

- [21] T. MALYSHEVA AND L.E. WHITE, *Sufficient conditions for Hadamard well-posedness of a coupled thermo-chemo-poroelastic system*. *Elect. J. Diff. Eqns.*, 2016(15) (2016) 1–17.
- [22] A.A. NEVILLE, P.C. MATTHEWS, AND H.M. BYRNE, *Interactions between pattern formation and domain growth*. *Bull. Math. Biol.*, 68(8) (2006) 1975–2003.
- [23] R. OYARZÚA AND R. RUIZ-BAIER, *Locking-free finite element methods for poroelasticity*. *SIAM J. Numer. Anal.*, 54(5) (2016) 2951–2973.
- [24] A. QUARTERONI AND A. VALLI, *Numerical Approximation of Partial Differential Equations*. Vol. 23 of Springer Ser. Comput. Math. Springer-Verlag Berlin Heidelberg (1994).
- [25] P. RECHO, A. HALLOU, AND E. HANNEZO, *Theory of mechano-chemical patterning in biphasic biological tissues*. *PNAS.*, 116(12) (2019) 5344–5349.
- [26] J. SCHNAKENBERG, *Simple chemical reaction systems with limit cycle behaviour*. *J. Theoret. Biol.*, 81(3) (1979) 389–400.
- [27] R.E. SHOWALTER, *Diffusion in poro-elastic media*. *J. Math. Anal. Appl.*, 251 (2000) 310–340.
- [28] R.E. SHOWALTER AND B. MOMKEN, *Single-phase flow in composite poroelastic media*. *Math. Methods Appl. Sci.*, 25 (2002) 115–139.
- [29] N. VERMA, B. GÓMEZ-VARGAS, L.M.D.O. VILACA, S. KUMAR, AND R. RUIZ-BAIER, *Advection-diffusion-reaction in poroelastic media. Part II: Linear stability analysis and numerical simulations*, submitted preprint (2019).

Centro de Investigación en Ingeniería Matemática (CI²MA)

PRE-PUBLICACIONES 2019

- 2019-24 JULIO ARACENA, CHRISTOPHER THRAVES: *The weighted sitting closer to friends than enemies problem in the line*
- 2019-25 RAIMUND BÜRGER, STEFAN DIEHL, MARÍA CARMEN MARTÍ, YOLANDA VÁSQUEZ: *A model of flotation with sedimentation: steady states and numerical simulation of transient operation*
- 2019-26 RAIMUND BÜRGER, RAFAEL ORDOÑEZ, MAURICIO SEPÚLVEDA, LUIS M. VILLADA: *Numerical analysis of a three-species chemotaxis model*
- 2019-27 ANÍBAL CORONEL, FERNANDO HUANCAS, MAURICIO SEPÚLVEDA: *Identification of space distributed coefficients in an indirectly transmitted diseases model*
- 2019-28 ANA ALONSO-RODRIGUEZ, JESSIKA CAMAÑO, EDUARDO DE LOS SANTOS, FRANCESCA RAPETTI: *A tree-cotree splitting for the construction of divergence-free finite elements: the high order case*
- 2019-29 RAIMUND BÜRGER, ENRIQUE D. FERNÁNDEZ NIETO, VICTOR OSORES: *A multi-layer shallow water approach for polydisperse sedimentation with sediment compressibility and mixture viscosity*
- 2019-30 LILIANA CAMARGO, BIBIANA LÓPEZ-RODRÍGUEZ, MAURICIO OSORIO, MANUEL SOLANO: *An HDG method for Maxwell equations in heterogeneous media*
- 2019-31 RICARDO OYARZÚA, SANDER RHEBERGEN, MANUEL SOLANO, PAULO ZUÑIGA: *Error analysis of a conforming and locking-free four-field formulation in poroelasticity*
- 2019-32 GABRIEL N. GATICA, MAURICIO MUNAR, FILANDER A. SEQUEIRA: *A mixed virtual element method for the Boussinesq problem on polygonal meshes*
- 2019-33 SERGIO CAUCAO, GABRIEL N. GATICA, RICARDO OYARZÚA, FELIPE SANDOVAL: *Residual-based a posteriori error analysis for the coupling of the Navier-Stokes and Darcy-Forchheimer equations*
- 2019-34 GABRIEL N. GATICA, RICARDO OYARZÚA, NATHALIE VALENZUELA: *A five-field augmented fully-mixed finite element method for the Navier-Stokes/Darcy coupled problem*
- 2019-35 LUIS MIGUEL DE OLIVEIRA VILACA, BRYAN GOMEZ-VARGAS, SARVESH KUMAR, RICARDO RUIZ-BAIER, NITESH VERMA: *Advection-diffusion-reaction in poroelastic media. Part I: Well-posedness and discrete analysis*

Para obtener copias de las Pre-Publicaciones, escribir o llamar a: DIRECTOR, CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE, TEL.: 41-2661324, o bien, visitar la página web del centro: <http://www.ci2ma.udec.cl>



**CENTRO DE INVESTIGACIÓN EN
INGENIERÍA MATEMÁTICA (CI²MA)
Universidad de Concepción**



Casilla 160-C, Concepción, Chile
Tel.: 56-41-2661324/2661554/2661316
<http://www.ci2ma.udec.cl>

