

UNIVERSIDAD DE CONCEPCIÓN



CENTRO DE INVESTIGACIÓN EN  
INGENIERÍA MATEMÁTICA (CI<sup>2</sup>MA)



An adaptive stabilized finite element method for the  
Stokes-Darcy coupled problem

RODOLFO ARAYA, CRISTIAN CÁRCAMO,  
ABNER POZA, EDUARDO VINO

PREPRINT 2022-07

SERIE DE PRE-PUBLICACIONES



# An adaptive stabilized finite element method for the Stokes–Darcy coupled problem

Rodolfo Araya<sup>a</sup>, Cristian Cárcamo<sup>a</sup>, Abner H. Poza<sup>b,\*</sup>, Eduardo Vino<sup>b</sup>

<sup>a</sup>*Departamento de Ingeniería Matemática & CI<sup>2</sup>MA, Universidad de Concepción, Casilla 160-C, Concepción, Chile*

<sup>b</sup>*Departamento de Matemática y Física Aplicadas, Universidad Católica de la Santísima Concepción, Casilla 297, Concepción, Chile*

---

## Abstract

For the Stokes–Darcy coupled problem, which models a fluid that flows in a free medium into a porous medium, we introduce and analyze an adaptive stabilized finite element method using Lagrange equal order element to approximate the velocity and pressure of the fluid. The interface conditions between both domains are given by mass conservation, the balance of normal forces, and the Beavers–Joseph–Saffman conditions. We prove the well-posedness of the discrete problem and present a convergence analysis with optimal error estimates in natural norms. Next, we introduce and analyze a residual-based a posteriori error estimator for the stabilized scheme. Finally, we present some numerical examples to show the quality of our scheme.

*Keywords:* Coupled Stokes–Darcy equation, stabilized finite element method, a priori error analysis, a posteriori error analysis.

---

## 1. Introduction

The coupling between a fluid that flows in a free medium into a porous medium is of particular interest by practitioners, scientists, engineering, and hydrology, for example, to predict how contamination spreads into streams, lakes, and rivers, affecting the water supply (see [35], and references therein). This coupling is also important in the filtration of blood through the arterial wall (for details of medical applications, see for example [6]). In the industrial sector, it is studied in the design of filters and in the hydrocarbon extraction process (for details, see [2, 25]). In this work, we consider the coupling of a fluid modeled by the Stokes equations with the fluid in a porous medium governed by the Darcy equations. At the interface, we have mass conservation, the balance of normal forces, and Beavers–Joseph–Saffman conditions [9, 32].

In the last decades, various ways of obtaining approximations of solutions by finite element methods for both problems separately can be found in the literature (for instance, for the Stokes equation [3, 8, 13, 22, 24, 33, 40], and for Darcy equation [4, 12, 14, 29]). Although it is known that both problems are mathematically very different, a unified study of both problems in a single domain can be found in [7, 11, 15, 17, 39]. In this shortlist of references, stabilized finite element methods are presented, assuming that the physical parameters of the problem allow generating the Darcy or Stokes equations in a limit case.

Regarding the coupled problem, there is a long list of discrete schemes to obtain approximated solutions. For example, in [23], a domain decomposition methodology introduces an iterative method to solve the problem. In [31], a complete theoretical analysis of a primal mixed scheme is presented using a Lagrange multiplier to obtain velocities with continuous normal trace. In [38], stable finite element space for the Stokes equations and a Galerkin least-squares formulation is used for the Darcy equations. In [27], the authors presents a conforming mixed finite element for the corresponding coupled problem using Bernardi–Raugel

---

\*Corresponding author

*Email addresses:* `rodolfo.araya@udec.cl` (Rodolfo Araya), `ccarcamo@udec.cl` (Cristian Cárcamo), `apoza@ucsc.cl` (Abner H. Poza), `evino@magister.ucsc.cl` (Eduardo Vino)

and Raviart–Thomas elements for the velocities and piecewise constants for the pressures. In [41], the mini–element for the Stokes equations and Brezzi–Douglas–Marini (BDM) with linear Lagrange approximations are used to solve the problem. For more schemes, see [18, 42, 19] and the references therein. Also, it is worth to mention the interior penalty method presented in [17], based in a Nitsche-type weak formulation, allowing to obtain approximations of piecewise linear continuous velocities and piecewise constant pressure; the scheme proposed in [37] which add jumps terms over the element edges of the velocities and used nonconforming Crouzeix–Raviart piecewise linear element for the velocities and piecewise constant for the pressures, and, finally, in [5], the authors used the classical mini–element discretization of the Darcy and Stokes problems.

In this work, we introduced and analyzed a new stabilized finite element method for the Stokes–Darcy coupled problem, which allows us to use equal order approximation spaces  $\mathbb{P}_k^d \times \mathbb{P}_k^d$ , for velocities and pressures, respectively, in both domains. Our discrete scheme uses the weak formulation presented in [38], and residual stabilized terms, for the Stokes equations, inspired in [26]. Using a weak inf–sup condition, the new method stability can be ensured; also by using a standard argument of stabilized finite element methods, we prove the convergence of the method with natural norms. Therefore, there are two main contributions in this work: the introduction and numerical analysis of this new scheme and the introduction of a reliable and efficient residual-based a posteriori error estimator (for other references in a posteriori error estimator for this problem, see [10, 18, 28]).

This paper is organized as follows: In Section 2, the model problem, with different boundary conditions on the interface between the free and porosity domains, is presented with its weak formulation. The new stabilized finite element method is introduced in Section 3, including its well-posedness. In Section 4, the a priori error analysis for equal order finite element spaces  $\mathbb{P}_k^d \times \mathbb{P}_k^d$  is developed. The a posteriori error is presented and analyzed in Section 5. Numerical results, showing the theoretical convergence rate and the performance of the residual-based a posteriori error estimator, can be found in Section 6. Finally, a technical result is proved in Appendix A.

## 2. Model problem and preliminary results

Let  $\Omega \subset \mathbb{R}^d$ ,  $d \in \{2, 3\}$ , be an open, bounded domain with Lipschitz continuous boundary  $\partial\Omega$ . This domain is divided into two disjoint open subdomains  $\Omega_S$  and  $\Omega_D$ , both with Lipschitz continuous boundaries, such that  $\bar{\Omega} = \bar{\Omega}_S \cup \bar{\Omega}_D$ . Here  $\Omega_S$  and  $\Omega_D$  represent the domains in the free and porous media, respectively. The interface between both media is given by  $\Gamma := \bar{\Omega}_S \cap \bar{\Omega}_D$  as shown in Figure 1. The remaining parts of the boundaries are  $\Gamma_S := \partial\Omega_S \setminus \Gamma$  and  $\Gamma_D := \partial\Omega_D \setminus \Gamma$ , with  $\Gamma_D$  divided in  $\Gamma_D^{\text{Dir}}$  and  $\Gamma_D^{\text{Neu}}$ , with  $\Gamma_D^{\text{Dir}} \cap \Gamma_D^{\text{Neu}} = \emptyset$  and  $\Gamma_D^{\text{Dir}} \neq \emptyset$ .




Figure 1: Representation of a possible computational domain  $\Omega$ .

The Stokes–Darcy coupled problem consists on finding the velocities  $\mathbf{u} := (\mathbf{u}_S, \mathbf{u}_D)$  and the pressures

$p := (p_S, p_D)$  such that they satisfy the system of equations

$$(P) \quad \begin{cases} -\nabla \cdot \boldsymbol{\sigma}(\mathbf{u}_S, p_S) = \mathbf{f}_S & \text{in } \Omega_S, \\ \nabla \cdot \mathbf{u}_S = 0 & \text{in } \Omega_S, \\ \mathbf{u}_S = \mathbf{0} & \text{on } \Gamma_S, \\ \nu \mathbf{u}_D + \kappa \nabla p_D = \mathbf{0} & \text{in } \Omega_D, \\ \nabla \cdot \mathbf{u}_D = g_D & \text{in } \Omega_D, \\ p_D = 0 & \text{on } \Gamma_D^{\text{Dir}}, \\ \mathbf{u}_D \cdot \mathbf{n}_D = \mathbf{0} & \text{on } \Gamma_D^{\text{Neu}}, \end{cases} \quad (2.1)$$

where  $\boldsymbol{\sigma}(\mathbf{u}_S, p_S) := 2\nu \boldsymbol{\varepsilon}(\mathbf{u}_S) - p_S \mathbf{I}$  is the stress rate tensor,  $\mathbf{I}$  the identity matrix,  $\boldsymbol{\varepsilon}(\mathbf{u}_S) := \frac{1}{2} (\nabla \mathbf{u}_S + \nabla \mathbf{u}_S^t)$  the deformation rate tensor,  $\nu > 0$  the viscosity of the fluid,  $\kappa > 0$  the permeability of the porous media and  $\mathbf{n}_D$  the outward unit normal vector on  $\Gamma_D^{\text{Neu}}$ . Here  $\mathbf{f}_S \in L^2(\Omega_S)^d$  and  $g_D \in L^2(\Omega_D)$ . This problem is completed with mass conservation, equilibrium of normal forces and the Beavers–Joseph–Saffman condition on  $\Gamma$  (for details see [9]):

$$\begin{cases} \mathbf{u}_D \cdot \mathbf{n}_D + \mathbf{u}_S \cdot \mathbf{n}_S = 0, \\ -\mathbf{n}_S \cdot \boldsymbol{\sigma}(\mathbf{u}_S, p_S) \mathbf{n}_S = p_D, \\ -\mathbf{n}_S \cdot \boldsymbol{\sigma}(\mathbf{u}_S, p_S) \boldsymbol{\tau}_i = \frac{\alpha_i}{\kappa^{1/2}} \mathbf{u}_S \cdot \boldsymbol{\tau}_i, \quad i = 1, \dots, d-1, \end{cases} \quad (2.2)$$

where  $\alpha_i$  are positives non-dimensional constants, and  $\boldsymbol{\tau}_i$  are the tangent vectors on  $\Gamma$ .

In the sequel we will use the following Hilbert spaces,

$$\begin{aligned} \mathbf{H}^S &:= \{\mathbf{v} \in H^1(\Omega_S)^d : \mathbf{v} = \mathbf{0} \text{ on } \Gamma_S\} \quad \text{and} \quad \mathbf{H}^D := L^2(\Omega_D)^d, \\ Q^S &:= L^2(\Omega_S) \quad \text{and} \quad Q^D := \{q \in H^1(\Omega_D) : q = 0 \text{ on } \Gamma_D^{\text{Dir}}\}, \\ \mathbf{H} &:= \mathbf{H}^S \times \mathbf{H}^D \quad \text{and} \quad Q := Q^S \times Q^D. \end{aligned}$$

A variational formulation of problem (2.1)–(2.2) can be written as: *Find*  $(\mathbf{u}, p) \in \mathbf{H} \times Q$  *such that*

$$B((\mathbf{u}, p), (\mathbf{v}, q)) = F(\mathbf{v}, q), \quad (2.3)$$

for all  $(\mathbf{v}, q) \in \mathbf{H} \times Q$ , where  $B : (\mathbf{H} \times Q) \times (\mathbf{H} \times Q) \rightarrow \mathbb{R}$  is the bilinear form defined by

$$\begin{aligned} B((\mathbf{u}, p); (\mathbf{v}, q)) &:= 2\nu \kappa (\boldsymbol{\varepsilon}(\mathbf{u}_S), \boldsymbol{\varepsilon}(\mathbf{v}_S))_{\Omega_S} - \kappa (p_S, \nabla \cdot \mathbf{v}_S)_{\Omega_S} + \kappa (q_S, \nabla \cdot \mathbf{u}_S)_{\Omega_S} \\ &\quad + \kappa^{1/2} \sum_{i=1}^{d-1} \alpha_i (\mathbf{u}_S \cdot \boldsymbol{\tau}_i, \mathbf{v}_S \cdot \boldsymbol{\tau}_i)_{\Gamma} + \kappa (p_D, \mathbf{v}_S \cdot \mathbf{n}_S)_{\Gamma} \\ &\quad + \nu (\mathbf{u}_D, \mathbf{v}_D)_{\Omega_D} + \kappa (\nabla p_D, \mathbf{v}_D)_{\Omega_D} - \kappa (\mathbf{u}_D, \nabla q_D)_{\Omega_D} - \kappa (\mathbf{u}_S \cdot \mathbf{n}_S, q_D)_{\Gamma} \\ &\quad + \frac{1}{2\nu} (\nu \mathbf{u}_D + \kappa \nabla p_D, -\nu \mathbf{v}_D + \kappa \nabla q_D)_{\Omega_D}, \end{aligned} \quad (2.4)$$

for all  $(\mathbf{u}, p), (\mathbf{v}, q) \in \mathbf{H} \times Q$ , and  $F : \mathbf{H} \times Q \rightarrow \mathbb{R}$  is the linear functional defined by

$$F(\mathbf{v}, q) := \kappa (\mathbf{f}_S, \mathbf{v}_S)_{\Omega_S} + \kappa (g_D, q_D)_{\Omega_D}, \quad (2.5)$$

for all  $(\mathbf{v}, q) \in \mathbf{H} \times Q$ . Now, it is well-known [38, Theorem 1] that (2.3) has a unique solution  $(\mathbf{u}, p) \in \mathbf{H} \times Q$ . In addition, we will use the following norm on the product space  $\mathbf{H} \times Q$ :

$$\|(\mathbf{w}, r)\| := \left\{ \nu \kappa \|\mathbf{w}_S\|_{1, \Omega_S}^2 + \kappa \|r_S\|_{0, \Omega_S}^2 + \nu \|\mathbf{w}_D\|_{0, \Omega_D}^2 + \frac{\kappa^2}{\nu} \|r_D\|_{1, \Omega_D}^2 \right\}^{1/2},$$

for all  $(\mathbf{w}, r) \in \mathbf{H} \times Q$ .

Throughout this paper  $C$  and  $C_i$ ,  $i > 0$  will denote positive constants independent of the mesh size  $h$ , but who may depend on the physical parameters of the equation.

The following result will be needed throughout the paper.

**Lemma 1.** *There exist positive constants  $C_{\text{Korn}}, C_t$ , such that*

$$C_{\text{Korn}} \|\mathbf{v}_S\|_{1, \Omega_S} \leq \|\varepsilon(\mathbf{v}_S)\|_{0, \Omega_S} \leq C_t \|\mathbf{v}_S\|_{1, \Omega_S},$$

for all  $\mathbf{v}_S \in \mathbf{H}^S$ .

*Proof.* See [20, Theorem 1.2-2]. □

### 3. The stabilized finite element method

From now on, we denote by  $\{\mathcal{T}_h^S\}_{h>0}$  and  $\{\mathcal{T}_h^D\}_{h>0}$  two regular families of triangulations of  $\bar{\Omega}_S$  and  $\bar{\Omega}_D$ , respectively, composed by simplexes that match at the interface  $\Gamma$ . For a  $\mathcal{T}_h^S$  or  $\mathcal{T}_h^D$ , we will denote by  $K$  the elements of the triangulation, and by  $\mathcal{E}_h^S$  the set of all edges (faces) of  $\mathcal{T}_h^S$ , with the splitting  $\mathcal{E}_h^S := \mathcal{E}_{\Omega_S} \cup \mathcal{E}_D^{\text{Dir}} \cup \mathcal{E}_{\Gamma_h}$ , where  $\mathcal{E}_{\Omega_S}$  stands for the edges (faces) lying in the interior of  $\Omega_S$ ,  $\mathcal{E}_D^{\text{Dir}}$  stands for the edges (faces) on the boundary  $\Gamma_S$ , and  $\mathcal{E}_{\Gamma_h}$  stands for the edges (faces) on the boundaries  $\Gamma$ . In the same way, we denote by  $\mathcal{E}_h^D$  the set of all edges (faces) of  $\mathcal{T}_h^D$ , with the splitting  $\mathcal{E}_h^D := \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{Neu}} \cup \mathcal{E}_D^{\text{Dir}} \cup \mathcal{E}_{\Gamma_h}$ , where  $\mathcal{E}_{\Omega_D}$  stands for the edges (faces) lying in the interior of  $\Omega_D$ , and  $\mathcal{E}_D^{\text{Neu}}$  and  $\mathcal{E}_D^{\text{Dir}}$  stand for the edges (faces) on the boundaries  $\Gamma_D^{\text{Neu}}$  and  $\Gamma_D^{\text{Dir}}$ , respectively. As usual  $h_T$  means the diameter of  $T$ ,  $h := \max_{T \in \mathcal{T}_h^S \cup \mathcal{T}_h^D} h_T$ , and

$h_F := |F|$  for  $F \in \mathcal{E}_h^S \cup \mathcal{E}_h^D$ . Finally, for each  $K \in \mathcal{T}_h^S \cup \mathcal{T}_h^D$  and  $F \in \mathcal{E}_h^S \cup \mathcal{E}_h^D$ , we denote by  $\mathcal{N}(K)$  the set of nodes of  $K$ ,  $\mathcal{N}(F)$  the set of nodes of  $F$ ,  $\mathcal{E}(K)$  the set of edges of  $K$ , and define

$$\begin{aligned} \omega_F^S &:= \bigcup_{\substack{F \in \mathcal{E}(K') \\ K' \in \mathcal{T}_h^S}} K', & \tilde{\omega}_K^S &:= \bigcup_{\substack{\mathcal{N}(K) \cap \mathcal{N}(K') \neq \emptyset \\ K' \in \mathcal{T}_h^S}} K', & \tilde{\omega}_F^S &:= \bigcup_{\substack{\mathcal{N}(F) \cap \mathcal{N}(K') \neq \emptyset \\ K' \in \mathcal{T}_h^S}} K', \\ \omega_F^D &:= \bigcup_{\substack{F \in \mathcal{E}(K') \\ K' \in \mathcal{T}_h^D}} K', & \tilde{\omega}_K^D &:= \bigcup_{\substack{\mathcal{N}(K) \cap \mathcal{N}(K') \neq \emptyset \\ K' \in \mathcal{T}_h^D}} K', & \tilde{\omega}_F^D &:= \bigcup_{\substack{\mathcal{N}(F) \cap \mathcal{N}(K') \neq \emptyset \\ K' \in \mathcal{T}_h^D}} K'. \end{aligned}$$

We introduce the following finite element subspaces of  $\mathbf{H}^S$ ,  $\mathbf{H}^D$ ,  $Q^S$  and  $Q^D$ , respectively:

$$\begin{aligned} \mathbf{H}_{h,k}^S &:= \{\mathbf{v} \in C(\bar{\Omega}_S)^d : \mathbf{v}|_K \in \mathbb{P}_k(K)^d, \quad \forall K \in \mathcal{T}_h^S\} \cap \mathbf{H}^S, \\ \mathbf{H}_{h,k}^D &:= \{\mathbf{v} \in C(\bar{\Omega}_D)^d : \mathbf{v}|_K \in \mathbb{P}_k(K)^d, \quad \forall K \in \mathcal{T}_h^D\}, \\ Q_{h,k}^S &:= \{q \in C(\bar{\Omega}_S) : q|_K \in \mathbb{P}_k(K), \quad \forall K \in \mathcal{T}_h^S\}, \\ Q_{h,k}^D &:= \{q \in C(\bar{\Omega}_D) : q|_K \in \mathbb{P}_k(K), \quad \forall K \in \mathcal{T}_h^D\} \cap Q^D, \end{aligned}$$

with  $k \geq 1$ , where  $\mathbb{P}_k$  stands for the space of polynomials of total degree less or equal to  $k$ .

Let  $\mathbf{H}_{h,k} := \mathbf{H}_{h,k}^S \times \mathbf{H}_{h,k}^D$  and  $Q_{h,k} := Q_{h,k}^S \times Q_{h,k}^D$ , and let  $\mathbf{u}_h := (\mathbf{u}_{h,S}, \mathbf{u}_{h,D}) \in \mathbf{H}_{h,k}$  and  $p_h := (p_{h,S}, p_{h,D}) \in Q_{h,k}$ . We consider the following discrete stabilized scheme: Find  $(\mathbf{u}_h, p_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$  such that

$$B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{v}_h, q_h)) = F_{\text{stab}}(\mathbf{v}_h, q_h), \quad (3.6)$$

for all  $(\mathbf{v}_h, q_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$ , where

$$\begin{aligned} B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{v}_h, q_h)) &:= B((\mathbf{u}_h, p_h); (\mathbf{v}_h, q_h)) \\ &+ \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( -2\nu \nabla \cdot \varepsilon(\mathbf{u}_{h,S}) + \nabla p_{h,S}, 2\nu \nabla \cdot \varepsilon(\mathbf{v}_{h,S}) + \nabla q_{h,S} \right)_K \\ &+ \theta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( \nabla \cdot \mathbf{u}_{h,S}, \nabla \cdot \mathbf{v}_{h,S} \right)_K + \nu \lambda \sum_{K \in \mathcal{T}_h^D} h_K^2 \left( \nabla \cdot \mathbf{u}_{h,D}, \nabla \cdot \mathbf{v}_{h,D} \right)_K, \end{aligned} \quad (3.7)$$

and

$$F_{\text{stab}}(\mathbf{v}_h, q_h) := F(\mathbf{v}_h, q_h) + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( \mathbf{f}_S, 2\nu \nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S}) + \nabla q_{h,S} \right)_K + \nu \lambda \sum_{K \in \mathcal{T}_h^D} h_K^2 \left( g_D, \nabla \cdot \mathbf{v}_{h,D} \right)_K.$$

Here  $\theta$  and  $\lambda$  are no negative constants, and  $0 < \beta < \frac{C_I}{2}$ , where  $C_I$  is the constant appearing in the following inverse inequality

$$C_I \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S})\|_{0,K}^2 \leq \|\boldsymbol{\varepsilon}(\mathbf{v}_{h,S})\|_{0,\Omega_S}^2 \quad \forall \mathbf{v}_{h,S} \in \mathbf{H}_{h,k}^S. \quad (3.8)$$

**Remark 1.** *The stabilized finite element method (3.6) is inspired by the scheme proposed in [38], where a standard formulation for the Stokes equation and a Galerkin least-squares formulation for the Darcy equation was introduced. In this new method, we introduce an additional stabilization to the Stokes equation, allowing equal order interpolation spaces for the fluid flow as for the flow in the porous media.*

In the rest of this work, over  $\mathbf{H}_{h,k} \times Q_{h,k}$  we will define the following mesh-dependent norm

$$\|(\mathbf{v}_h, q_h)\|_h := \left\{ \nu \kappa \|\mathbf{v}_{h,S}\|_{1,\Omega_S}^2 + \kappa \|q_{h,S}\|_{0,\Omega_S}^2 + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |q_{h,S}|_{1,K}^2 + \nu \|\mathbf{v}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{\nu} |q_{h,D}|_{1,\Omega_D}^2 \right\}^{1/2},$$

for all  $(\mathbf{v}_h, q_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$ .

As it is well known, the subspaces  $\mathbf{H}_{h,k}^S$  and  $Q_{h,k}^S$  do not satisfy a discrete inf-sup condition, but they satisfy the following weak inf-sup condition

**Lemma 2.** *There exist positive constants  $C_1$  and  $C_2$ , independent of  $\nu$  and  $h$ , such that*

$$\sup_{\mathbf{v}_{h,S} \in \mathbf{H}_{h,k}^S \cap H_0^1(\Omega_S)^d} \frac{(q_{h,S}, \nabla \cdot \mathbf{v}_{h,S})_{\Omega_S}}{\|\mathbf{v}_{h,S}\|_{1,\Omega_S}} \geq C_1 \|q_{h,S}\|_{0,\Omega_S} - C_2 \left\{ \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 |q_{h,S}|_{1,K}^2 \right\}^{1/2}, \quad (3.9)$$

for all  $q_{h,S} \in Q_{h,k}^S$ .

*Proof.* See [26, Lemma 3.3]. □

The following result will be necessary to prove the well-posedness of the stabilized finite element problem (3.6). The proof is based on similar arguments to those used in [4, Lemma 3].

**Lemma 3.** *For all  $(\mathbf{u}_h, p_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$ , there exists a positive constant  $C$ , independent of  $h$ ,  $\nu$ ,  $\kappa$  and  $\alpha_i$ , such that*

$$\sup_{(\mathbf{v}_h, q_h) \in \mathbf{H}_{h,k} \times Q_{h,k}} \frac{B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{v}_h, q_h))}{\|(\mathbf{v}_h, q_h)\|_h} \geq C \|(\mathbf{u}_h, p_h)\|_h.$$

*Proof.* Using Cauchy–Schwarz inequality, (3.8) and Lemma 1, we get

$$\begin{aligned}
& B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{u}_h, p_h)) \\
& \geq 2\nu \kappa \|\varepsilon(\mathbf{u}_{h,S})\|_{0,\Omega_S}^2 + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( -2\nu \nabla \cdot \varepsilon(\mathbf{u}_{h,S}) + \nabla p_{h,S}, 2\nu \nabla \cdot \varepsilon(\mathbf{u}_{h,S}) + \nabla p_{h,S} \right)_K \\
& \quad + \nu \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{1}{2\nu} (\nu \mathbf{u}_{h,D} + \kappa \nabla p_{h,D}, -\nu \mathbf{u}_{h,D} + \kappa \nabla p_{h,D})_{\Omega_D} \\
& \geq 2\nu \kappa \|\varepsilon(\mathbf{u}_{h,S})\|_{0,\Omega_S}^2 - 4\nu \kappa \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \varepsilon(\mathbf{u}_{h,S})\|_{0,K}^2 + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 + \frac{\nu}{2} \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{2\nu} |p_{h,D}|_{1,\Omega_D}^2 \\
& \geq 2\nu \kappa \left(1 - \frac{2\beta}{CI}\right) \|\varepsilon(\mathbf{u}_{h,S})\|_{0,\Omega_S}^2 + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 + \frac{\nu}{2} \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{2\nu} |p_{h,D}|_{1,\Omega_D}^2 \\
& \geq C_3 \nu \kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S}^2 + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 + \frac{\nu}{2} \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{2\nu} |p_{h,D}|_{1,\Omega_D}^2. \tag{3.10}
\end{aligned}$$

Let  $\mathbf{w}_S \in \mathbf{H}_{h,k}^S \cap H_0^1(\Omega_S)^d$  be a function for which the supremum of Lemma 2 is attained. Then

$$\begin{aligned}
& B_{\text{stab}}((\mathbf{u}_h, p_h); ((-\mathbf{w}_S, \mathbf{0}), (0, 0))) = B_{\text{stab}}(((\mathbf{u}_{h,S}, \mathbf{u}_{h,D}), (p_{h,S}, p_{h,D})); ((-\mathbf{w}_S, \mathbf{0}), (0, 0))) \\
& = -2\nu \kappa (\varepsilon(\mathbf{u}_{h,S}), \varepsilon(\mathbf{w}_S))_{\Omega_S} - \kappa^{1/2} \sum_{i=1}^{d-1} \alpha_i (\mathbf{u}_{h,S} \cdot \boldsymbol{\tau}_i, \mathbf{w}_S \cdot \boldsymbol{\tau}_i)_{\Gamma} + \kappa (p_{h,S}, \nabla \cdot \mathbf{w}_S)_{\Omega_S} \\
& \quad + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( 2\nu \nabla \cdot \varepsilon(\mathbf{u}_{h,S}), 2\nu \nabla \cdot \varepsilon(\mathbf{w}_S) \right)_K - \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \left( \nabla p_{h,S}, 2\nu \nabla \cdot \varepsilon(\mathbf{w}_S) \right)_K \\
& \quad - \theta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} (\nabla \cdot \mathbf{u}_{h,S}, \nabla \cdot \mathbf{w}_S)_K.
\end{aligned}$$

Now assume that  $\|\mathbf{w}_S\|_{1,\Omega_S} = \|p_{h,S}\|_{0,\Omega_S}$ , using Cauchy–Schwarz inequality, the inverse inequality (3.8) with Lemma 1, we have

$$\begin{aligned}
& B_{\text{stab}}((\mathbf{u}_h, p_h); ((-\mathbf{w}_S, \mathbf{0}), (0, 0))) \\
& \geq -2\nu \kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S} \|\mathbf{w}_S\|_{1,\Omega_S} + \kappa (p_{h,S}, \nabla \cdot \mathbf{w}_S)_{\Omega_S} \\
& \quad - \kappa \beta 4\nu \left\{ \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \varepsilon(\mathbf{u}_{h,S})\|_{0,K}^2 \right\}^{1/2} \left\{ \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \varepsilon(\mathbf{w}_S)\|_{0,K}^2 \right\}^{1/2} \\
& \quad - 2\kappa \left\{ \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 |p_{h,S}|_{1,K}^2 \right\}^{1/2} \left\{ \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \varepsilon(\mathbf{w}_S)\|_{0,K}^2 \right\}^{1/2} - d\theta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \|\mathbf{u}_{h,S}\|_{1,K} \|\mathbf{w}_S\|_{1,K} \\
& \geq -C_4 \nu \kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S} \|\mathbf{w}_S\|_{1,\Omega_S} + \kappa (p_{h,S}, \nabla \cdot \mathbf{w}_S)_{\Omega_S} - C_5 \kappa \left\{ \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 |p_{h,S}|_{1,K}^2 \right\}^{1/2} \|\mathbf{w}_S\|_{1,\Omega_S} \\
& \quad - \frac{C_6 \theta}{\nu} \|\mathbf{u}_{h,S}\|_{1,\Omega_S} \|\mathbf{w}_S\|_{1,\Omega_S} \\
& \geq -\left(C_4 + \frac{C_6 \theta}{\kappa \nu^2}\right) \nu \kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S} \|\mathbf{w}_S\|_{1,\Omega_S} + C_1 \kappa \|p_{h,S}\|_{0,\Omega_S}^2 - (C_2 + C_5) \kappa \left\{ \beta \sum_{K \in \mathcal{T}_h^S} h_K^2 |p_{h,S}|_{1,K}^2 \right\}^{1/2} \|p_{h,S}\|_{0,\Omega_S}.
\end{aligned}$$



Moreover, using Young inequality with positive constants  $\gamma_1$  and  $\gamma_2$  chosen small enough, we can conclude that

$$\begin{aligned}
& B_{\text{stab}}((\mathbf{u}_h, p_h); ((-\mathbf{w}_S, \mathbf{0}), (0, 0))) \\
& \geq -\frac{1}{2\gamma_1} \left( C_4 + \frac{C_6\theta}{\kappa\nu^2} \right) \nu\kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S}^2 + \kappa \left( C_1 - \frac{\gamma_1}{2} \left( C_4 + \frac{C_6\theta}{\kappa\nu^2} \right) \nu - \frac{(C_2 + C_5)\nu\gamma_2}{2} \right) \|p_{h,S}\|_{0,\Omega_S}^2 \\
& \quad - \frac{C_2 + C_5}{2\gamma_2} \kappa\beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 \\
& \geq -C_7\nu\kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S}^2 + C_8\kappa \|p_{h,S}\|_{0,\Omega_S}^2 - C_9\kappa\beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2.
\end{aligned} \tag{3.11}$$

Defining  $(\mathbf{v}_h, q_h) := (\mathbf{u}_h, p_h) + \delta((-\mathbf{w}_S, \mathbf{0}), (0, 0))$ , and combining (3.10) and (3.11), we get

$$\begin{aligned}
& B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{v}_h, q_h)) = B_{\text{stab}}((\mathbf{u}_h, p_h); (\mathbf{u}_h, q_h)) + \delta B_{\text{stab}}((\mathbf{u}_h, p_h); ((-\mathbf{w}_S, \mathbf{0}), (0, 0))) \\
& \geq \nu\kappa (C_3 - \delta C_7) \|\mathbf{u}_{h,S}\|_{1,\Omega_S}^2 + \kappa (1 - \delta C_9) \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 + \frac{\nu}{2} \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{2\nu} |p_{h,D}|_{1,\Omega_D}^2 \\
& \quad + C_8\delta\kappa \|p_{h,S}\|_{0,\Omega_S}^2 \\
& \geq C \left\{ \nu\kappa \|\mathbf{u}_{h,S}\|_{1,\Omega_S}^2 + \kappa \|p_{h,S}\|_{0,\Omega_S}^2 + \kappa\beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} |p_{h,S}|_{1,K}^2 + \nu \|\mathbf{u}_{h,D}\|_{0,\Omega_D}^2 + \frac{\kappa^2}{\nu} |p_{h,D}|_{1,\Omega_D}^2 \right\} \\
& = C \|(\mathbf{u}_h, p_h)\|_h^2,
\end{aligned} \tag{3.12}$$

choosing  $0 < \delta < \min \left\{ \frac{C_3}{C_7}, \frac{1}{C_9}, \frac{1}{\nu^{1/2}} \right\}$ .

On the other hand, we have

$$\|(\mathbf{v}_h, q_h)\|_h \leq \|(\mathbf{u}_h, p_h)\|_h + \delta\nu^{1/2}\kappa^{1/2} \|p_{h,S}\|_{0,\Omega_S} \leq C \|(\mathbf{u}_h, p_h)\|_h,$$

which combined with (3.12) proves the stability estimate.  $\square$

#### 4. A priori error analysis

We begin this section by recalling a local trace theorem and the definition and properties of the interpolation operators that we will use during the proof of the convergence of the stabilized finite element scheme (3.6).

**Lemma 4.** *There exists a positive constant  $C$ , independent of  $h_K$ , such that*

$$\|\psi\|_{0,\partial K}^2 \leq C \{ h_K^{-1} \|\psi\|_{0,K}^2 + h_K |\psi|_{1,K}^2 \},$$

for all  $K \in \mathcal{T}_h^S \cup \mathcal{T}_h^D$  and all  $\psi \in H^1(K)$ .

*Proof.* See [1, Theorem 3.10] or [16, (10.3.8)].  $\square$

We consider the Lagrange interpolant operator  $\mathbf{\Pi}_h^S : H^{k+1}(\Omega_S)^d \cap \mathbf{H}^S \rightarrow \mathbf{H}_{h,k}^S$ , and the Clément interpolation operator  $\mathcal{C}_h^S : H^k(\Omega) \rightarrow Q_{h,k}^S$ , such that (see [24] for details) for all  $K \in \mathcal{T}_h^S$ , we have

$$|\mathbf{u}_S - \mathbf{\Pi}_h^S \mathbf{u}_S|_{l,K} \leq C h_K^{s-l} |\mathbf{u}_S|_{s,K}, \tag{4.13}$$

$$\|p_S - \mathcal{C}_h^S p_S\|_{0,K} \leq C h_K^s \|p_S\|_{s,\tilde{\omega}_K^S}, \tag{4.14}$$

for all  $\mathbf{u}_S \in H^s(K)^d$  and all  $p_S \in H^s(\tilde{\omega}_K^S)$  with  $0 \leq l \leq 1$ ,  $1 \leq s \leq k+1$ . Here  $C$  are positive constants independent of  $h$ . Also, we consider the Clément interpolation operator  $\mathcal{C}_h^D : H^k(\Omega_D)^d \rightarrow \mathbf{H}_{h,k}^D$ , and the Lagrange interpolant operator  $\Pi_h^D : H^{k+1}(\Omega_D) \cap Q^D \rightarrow Q_{h,k}^D$ , such that for all  $K \in \mathcal{T}_h^D$ , we have

$$\|\mathbf{u}_D - \mathcal{C}_h^D\|_{0,K} \leq Ch_K^s \|\mathbf{u}_D\|_{s,\tilde{\omega}_K^D}, \quad (4.15)$$

$$|p_D - \Pi_h^D p_D|_{l,K} \leq Ch_K^{s-l} |p_D|_{s,K}, \quad (4.16)$$

for all  $\mathbf{u}_D \in H^s(\tilde{\omega}_K^D)^d$  and all  $p_D \in H^s(K)$  with  $0 \leq l \leq 1$ ,  $1 \leq s \leq k+1$ , where  $C$  are positive constants independent of  $h$ .

**Lemma 5.** *Let  $(\mathbf{u}, p) = ((\mathbf{u}_S, \mathbf{u}_D), (p_S, p_D))$  and  $(\mathbf{u}_h, p_h) = ((\mathbf{u}_{h,S}, \mathbf{u}_{h,D}), (p_{h,S}, p_{h,D}))$  be the solutions of (2.3) and (3.6), respectively. If  $\mathbf{u}_S \in H^2(\Omega_S)^d$ ,  $\mathbf{u}_D \in H^1(\Omega_D)^d$  and  $p_S \in H^1(\Omega_S)$ , then it holds*

$$B_{\text{stab}}((\mathbf{u} - \mathbf{u}_h, p - p_h); (\mathbf{v}_h, q_h)) = 0,$$

for all  $(\mathbf{v}_h, q_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$ .

*Proof.* Using the fact that  $\mathbf{u}_S \in H^2(\Omega_S)^d$  and  $p_S \in H^1(\Omega_S)$ , we get that  $\mathbf{f}_S - \nabla \cdot (2\nu\varepsilon(\mathbf{u}_S)) + \nabla p_S = \mathbf{0}$  and  $\nabla \cdot \mathbf{u}_S = 0$ . Using the fact that  $\mathbf{u}_D \in H^1(\Omega_D)^d$  we get  $\nabla \cdot \mathbf{u}_D = 0$ , thus the stabilized terms in the definition of  $B_{\text{stab}}$  vanish.  $\square$

**Theorem 6.** *Let  $(\mathbf{u}, p)$  and  $(\mathbf{u}_h, p_h)$  be the solutions of (2.3) and (3.6), respectively. Assume that  $(\mathbf{u}, p) = ((\mathbf{u}_S, \mathbf{u}_D), (p_S, p_D)) \in H^{k+1}(\Omega_S)^d \cap \mathbf{H}^S \times H^k(\Omega_D)^d \times H^k(\Omega_S) \times H^{k+1}(\Omega_D) \cap Q^D$ , then*

$$\|(\mathbf{u} - \mathbf{u}_h, p - p_h)\| \leq Ch^k \{ \|\mathbf{u}_S\|_{k+1,\Omega_S} + \|p_S\|_{k,\Omega_S} + \|\mathbf{u}_D\|_{k,\Omega_D} + \|p_D\|_{k+1,\Omega_D} \},$$

with  $C > 0$  independent of  $h$ .

*Proof.* We consider the following notations:

$$\begin{aligned} \eta^{\mathbf{u}_S} &:= \mathbf{u}_S - \mathbf{\Pi}_h^S \mathbf{u}_S, & \eta^{p_S} &:= p_S - \mathcal{C}_h^S p_S, \\ \eta^{\mathbf{u}_D} &:= \mathbf{u}_D - \mathcal{C}_h^D \mathbf{u}_D, & \eta^{p_D} &:= p_D - \Pi_h^D p_D, \\ \eta^{\mathbf{u}} &:= (\eta^{\mathbf{u}_S}, \eta^{\mathbf{u}_D}), & \eta^p &:= (\eta^{p_S}, \eta^{p_D}), \\ e_h^{\mathbf{u}} &:= (\mathbf{u}_{h,S} - \mathbf{\Pi}_h^S \mathbf{u}_S, \mathbf{u}_{h,D} - \mathcal{C}_h^D \mathbf{u}_D), & e_h^p &:= (p_{h,S} - \mathcal{C}_h^S p_S, p_{h,D} - \Pi_h^D p_D). \end{aligned}$$

Using the definition of  $B_{\text{stab}}$  given in (3.7), Lemma 5 and Cauchy–Schwarz inequality, we have

$$\begin{aligned}
& B_{\text{stab}}((e_h^{\mathbf{u}}, e_h^p); (\mathbf{v}_h, q_h)) = B_{\text{stab}}((\eta^{\mathbf{u}}, \eta^p); (\mathbf{v}_h, q_h)) - B_{\text{stab}}((\mathbf{u} - \mathbf{u}_h, p - p_h); (\mathbf{v}_h, q_h)) \\
& \leq 2\nu\kappa \|\eta^{\mathbf{u}^S}\|_{1,\Omega_S} \|\mathbf{v}_{h,S}\|_{1,\Omega_S} + \kappa^{1/2} \sum_{i=1}^{d-1} \alpha_i \sum_{F \in \mathcal{E}_{\Gamma_h}} \|\eta^{\mathbf{u}^S} \cdot \boldsymbol{\tau}_i\|_{0,F} \|\mathbf{v}_{h,S} \cdot \boldsymbol{\tau}_i\|_{0,F} + \kappa \sum_{F \in \mathcal{E}_{\Gamma_h}} \|\eta^{p_D}\|_{0,F} \|\mathbf{v}_{h,S} \cdot \mathbf{n}_S\|_{0,F} \\
& \quad + \kappa \|\eta^{p_S}\|_{0,\Omega_S} \|\nabla \cdot \mathbf{v}_{h,S}\|_{0,\Omega_S} + \kappa \|q_{h,S}\|_{0,\Omega_S} \|\nabla \cdot \eta^{\mathbf{u}^S}\|_{0,\Omega_S} \\
& \quad + \kappa \beta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \|-2\nu \nabla \cdot \boldsymbol{\varepsilon}(\eta^{\mathbf{u}^S}) + \nabla \eta^{p_S}\|_{0,K} \|2\nu \nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S}) + \nabla q_{h,S}\|_{0,K} \\
& \quad + \nu \|\eta^{\mathbf{u}^D}\|_{0,\Omega_D} \|\mathbf{v}_{h,D}\|_{0,\Omega_D} + \kappa |\eta^{p_D}|_{1,\Omega_D} \|\mathbf{v}_{h,D}\|_{0,\Omega_D} + \kappa \|\eta^{\mathbf{u}^D}\|_{0,\Omega_D} |q_{h,D}|_{1,\Omega_D} + \kappa \sum_{F \in \mathcal{E}_{\Gamma_h}} \|\eta^{\mathbf{u}^S} \cdot \mathbf{n}_S\|_{0,F} \|q_{h,D}\|_{0,F} \\
& \quad + \frac{1}{\nu} \|\nu \eta^{\mathbf{u}^D} + \kappa \nabla \eta^{p_D}\|_{0,\Omega_D} \|- \nu \mathbf{v}_{h,D} + \kappa \nabla q_{h,D}\|_{0,\Omega_D} \\
& \quad + \theta \sum_{K \in \mathcal{T}_h^S} \frac{h_K^2}{\nu} \|\nabla \cdot \eta^{\mathbf{u}^S}\|_{0,K} \|\nabla \cdot \mathbf{v}_{h,S}\|_{0,K} + \nu \lambda \sum_{K \in \mathcal{T}_h^D} h_K^2 \|\nabla \cdot \eta^{\mathbf{u}^D}\|_{0,K} \|\nabla \cdot \mathbf{v}_{h,D}\|_{0,K} \\
& \leq C \left\{ \|\eta^{\mathbf{u}^S}\|_{1,\Omega_S}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{\mathbf{u}^S}\|_{0,F}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{p_D}\|_{0,F}^2 + \|\eta^{p_S}\|_{0,\Omega_S}^2 \right. \\
& \quad + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|-2\nu \nabla \cdot \boldsymbol{\varepsilon}(\eta^{\mathbf{u}^S}) + \nabla \eta^{p_S}\|_{0,K}^2 + \|\eta^{\mathbf{u}^D}\|_{0,\Omega_D}^2 + |\eta^{p_D}|_{1,\Omega_D}^2 + \|\nu \eta^{\mathbf{u}^D} + \kappa \nabla \eta^{p_D}\|_{0,\Omega_D}^2 \\
& \quad \left. + \sum_{K \in \mathcal{T}_h^S} h_K^2 |\eta^{\mathbf{u}^S}|_{1,K}^2 + \sum_{K \in \mathcal{T}_h^D} h_K^2 |\eta^{\mathbf{u}^D}|_{1,K}^2 \right\}^{1/2} \\
& \quad \left\{ \|\mathbf{v}_{h,S}\|_{1,\Omega_S}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|\mathbf{v}_{h,S}\|_{0,F}^2 + \|q_{h,S}\|_{0,\Omega_S}^2 + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|2\nu \nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S}) + \nabla q_{h,S}\|_{0,K}^2 \right. \\
& \quad + \|\mathbf{v}_{h,D}\|_{0,\Omega_D}^2 + |q_{h,D}|_{1,\Omega_D}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|q_{h,D}\|_{0,F}^2 + \|- \nu \mathbf{v}_{h,D} + \kappa \nabla q_{h,D}\|_{0,\Omega_D}^2 \\
& \quad \left. + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \mathbf{v}_{h,S}\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h^D} h_K^2 \|\nabla \cdot \mathbf{v}_{h,D}\|_{0,K}^2 \right\}^{1/2}. \tag{4.17}
\end{aligned}$$

Now, using mesh regularity, (4.13), and Lemma 4, we get

$$\sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{\mathbf{u}^S}\|_{0,F}^2 \leq C \sum_{K \in \mathcal{T}_h^S} h_K^{-1} \{h_K^{-1} \|\eta^{\mathbf{u}^S}\|_{0,K}^2 + h_K |\eta^{\mathbf{u}^S}|_{1,K}^2\} \leq C h^{2k} \|\mathbf{u}_S\|_{k+1,\Omega_S}. \tag{4.18}$$

Similar arguments allow us to obtain the following estimates:

$$\begin{aligned}
& \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{p_D}\|_{0,F}^2 \leq C h^{2k} \|p_D\|_{k+1,\Omega_D}, \\
& \sum_{K \in \mathcal{T}_h^S} h_K^2 \|-2\nu \nabla \cdot \boldsymbol{\varepsilon}(\eta^{\mathbf{u}^S}) + \nabla \eta^{p_S}\|_{0,K}^2 \leq C h^{2k} \left[ \|\mathbf{u}_S\|_{k+1,\Omega_S}^2 + \|p_S\|_{k,\Omega_S}^2 \right].
\end{aligned}$$

Thus, it follows that

$$\begin{aligned}
& \|\eta^{\mathbf{u}_S}\|_{1,\Omega_S}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{\mathbf{u}_S}\|_{0,F}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F^{-1} \|\eta^{p_D}\|_{0,F}^2 + \|\eta^{p_S}\|_{0,\Omega_S}^2 \\
& + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|-2\nu \nabla \cdot \boldsymbol{\varepsilon}(\eta^{\mathbf{u}_S}) + \nabla \eta^{p_S}\|_{0,K}^2 + \|\eta^{\mathbf{u}_D}\|_{0,\Omega_D}^2 + \|\eta^{p_D}\|_{1,\Omega_D}^2 + \|\nu \eta^{\mathbf{u}_D} + \kappa \nabla \eta^{p_D}\|_{0,\Omega_D}^2 \\
& + \sum_{K \in \mathcal{T}_h^S} h_K^2 |\eta^{\mathbf{u}_S}|_{1,K}^2 + \sum_{K \in \mathcal{T}_h^D} h_K^2 |\eta^{\mathbf{u}_D}|_{1,K}^2 \leq C h^{2k} \left\{ \|\mathbf{u}_S\|_{k+1,\Omega_S}^2 + \|p_S\|_{k,\Omega_S}^2 + \|\mathbf{u}_D\|_{k,\Omega_D}^2 + \|p_D\|_{k+1,\Omega_D}^2 \right\}.
\end{aligned} \tag{4.19}$$

On the other hand, using inverse and Poincaré's inequalities, mesh regularity, and Lemma 4, we obtain

$$\begin{aligned}
& \|\mathbf{v}_{h,S}\|_{1,\Omega_S}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|\mathbf{v}_{h,S}\|_{0,F}^2 + \|q_{h,S}\|_{0,\Omega_S}^2 + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|2\nu \nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S}) + \nabla q_{h,S}\|_{0,K}^2 \\
& + \|\mathbf{v}_{h,D}\|_{0,\Omega_D}^2 + \|q_{h,D}\|_{1,\Omega_D}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|q_{h,D}\|_{0,F}^2 + \|- \nu \mathbf{v}_{h,D} + \kappa \nabla q_{h,D}\|_{0,\Omega_D}^2 \\
& + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|\nabla \cdot \mathbf{v}_{h,S}\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h^D} h_K^2 \|\nabla \cdot \mathbf{v}_{h,D}\|_{0,K}^2 \\
& \leq C \left\{ \|(\mathbf{v}_h, q_h)\|_h^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|\mathbf{v}_{h,S}\|_{0,F}^2 + \sum_{K \in \mathcal{T}_h^S} h_K^2 \|2\nu \nabla \cdot \boldsymbol{\varepsilon}(\mathbf{v}_{h,S}) + \nabla q_{h,S}\|_{0,K}^2 + \frac{\kappa^2}{\nu} \sum_{F \in \mathcal{E}_{\Gamma_h}} h_F \|q_{h,D}\|_{0,F}^2 \right\} \\
& \leq C \|(\mathbf{v}_h, q_h)\|_h^2.
\end{aligned} \tag{4.20}$$

Now, from (4.17), (4.19), (4.20) and Lemma 3, we have

$$\|(\mathbf{e}_h^{\mathbf{u}}, \mathbf{e}_h^p)\|^2 \leq \|(\mathbf{e}_h^{\mathbf{u}}, \mathbf{e}_h^p)\|_h^2 \leq C h^{2k} \left\{ \|\mathbf{u}_S\|_{k+1,\Omega_S}^2 + \|p_S\|_{k,\Omega_S}^2 + \|\mathbf{u}_D\|_{k,\Omega_D}^2 + \|p_D\|_{k+1,\Omega_D}^2 \right\}. \tag{4.21}$$

Using interpolation properties (4.13)-(4.16), we obtain the estimate

$$\|(\eta^{\mathbf{u}}, \eta^p)\|^2 \leq C h^{2k} \left\{ \|\mathbf{u}_S\|_{k+1,\Omega_S}^2 + \|p_S\|_{k,\Omega_S}^2 + \|\mathbf{u}_D\|_{k,\Omega_D}^2 + \|p_D\|_{k+1,\Omega_D}^2 \right\}. \tag{4.22}$$

The result follows using triangle inequality, (4.21) and (4.22).  $\square$

## 5. A posteriori error analysis

This section introduces a residual a posteriori error estimator for the stabilized finite element method (3.6). Throughout this section, we will assume, for simplicity, that  $\mathbf{f}_S$  and  $g_D$  are piecewise polynomials functions in  $\Omega_S$  and  $\Omega_D$ , respectively.

### 5.1. Preliminaries results

For the reliability of the error estimator, we need to define standard bubble functions and some of the results associated with them. We will define these functions considering the case in three dimensions ( $d = 3$ ), but the results are still valid in two dimensions ( $d = 2$ ). For all  $K \in \mathcal{T}_h^S \cup \mathcal{T}_h^D$ , we define the element bubble function  $b_K$  by

$$b_K := (d+1)^{d+1} \prod_{x \in \mathcal{N}(K)} \lambda_x,$$

where  $\lambda_x$  corresponds to the barycentric coordinates associated to node  $x$ . Let  $\hat{K}$  be the standard reference element with vertices  $\hat{n}_1 := (1, 0, 0)$ ,  $\hat{n}_2 := (0, 1, 0)$ ,  $\hat{n}_3 := (0, 0, 1)$  and  $\hat{n}_4 := (0, 0, 0)$ , we define the edge bubble function by

$$b_{\hat{F}} := d^d \hat{\lambda}_1 \hat{\lambda}_2 \hat{\lambda}_4,$$

where  $\hat{F} := \{(\hat{x}, \hat{y}, 0) \in \mathbb{R}^d : 0 \leq \hat{x} + \hat{y} \leq 1, \hat{x} \in [0, 1]\}$ . For  $F \in \mathcal{E}_h^S \cup \mathcal{E}_h^D$  assume that  $\omega_F^S = K_1 \cup K_2$  with  $K_1, K_2 \in \mathcal{T}_h^S$  or  $\omega_F^D = K_1 \cup K_2$  with  $K_1, K_2 \in \mathcal{T}_h^D$ . Let  $G_{F,i}$  be the (orientation preserving) affine transformation such that  $G_{F,i}(\hat{K}) = K_i$  and  $G_{F,i}(\hat{F}) = F$ , with  $i = 1, 2$ . We define the bubble function associated with  $F \in \mathcal{E}_h^S$  by

$$b_F^S := \begin{cases} b_{\hat{F}} \circ G_{F,i}^{-1}, & \text{on } K_i, \quad i = 1, 2, \\ 0 & \text{on } \Omega_S \setminus \omega_F^S, \end{cases} \quad (5.23)$$

and with  $F \in \mathcal{E}_h^D$  by

$$b_F^D := \begin{cases} b_{\hat{F}} \circ G_{F,i}^{-1}, & \text{on } K_i, \quad i = 1, 2, \\ 0 & \text{on } \Omega_D \setminus \omega_F^D. \end{cases} \quad (5.24)$$

Let  $\hat{\Pi} := \{(x, y, 0) : (x, y) \in \mathbb{R}^2\}$  and let  $\hat{Q} : \mathbb{R}^d \rightarrow \hat{\Pi}$  be the orthogonal projection from  $\mathbb{R}^d$  to  $\hat{\Pi}$ . We introduce the lifting operator  $\hat{P}_{\hat{F}} : \mathbb{P}_k(\hat{F}) \rightarrow \mathbb{P}_k(\hat{K})$  given by

$$\hat{s} \mapsto \hat{P}_{\hat{F}}(\hat{s}) = \hat{s} \circ \hat{Q}.$$

Let  $K_i \subseteq \omega_F^S \cup \omega_F^D$ . We define the lifting operator  $P_{F,K_i} : \mathbb{P}_k(F) \rightarrow \mathbb{P}_k(K_i)$  by

$$P_{F,K_i}(s) = \hat{P}_{\hat{F}}(s \circ G_{F,i}) \circ G_{F,i}^{-1}.$$

Using these notations, we can define a lifting operator  $P_F^S : \mathbb{P}_k(F) \rightarrow \mathbb{P}_k(\omega_F^S)$  by

$$s \in \mathbb{P}_k(F) \mapsto P_F^S(s) := \begin{cases} P_{F,K_1}(s) & \text{in } K_1, \\ P_{F,K_2}(s) & \text{in } K_2, \end{cases}$$

for  $\mathbf{s} = (s_1, s_2, s_3) \in \mathbb{P}_k(F)^d$ , we define  $\mathcal{P}_F^S(\mathbf{s})$  by

$$\mathcal{P}_F^S(\mathbf{s}) := (P_F^S(s_1), P_F^S(s_2), P_F^S(s_3)).$$

Similarily, we define

$$\mathcal{P}_F^D(\mathbf{s}) := (P_F^D(s_1), P_F^D(s_2), P_F^D(s_3)),$$

where  $P_F^D : \mathbb{P}_k(F) \rightarrow \mathbb{P}_k(\omega_F^D)$  by

$$s \in \mathbb{P}_k(F) \mapsto P_F^D(s) := \begin{cases} P_{F,K_1}(s) & \text{in } K_1, \\ P_{F,K_2}(s) & \text{in } K_2. \end{cases}$$

The next result can be prove using scaling arguments.

**Theorem 7.** *Let  $b_K$  be the bubble function corresponding to  $K \in \mathcal{T}_h^S \cup \mathcal{T}_h^D$ . Then for all  $\mathbf{v}_h \in \mathbb{P}_n$ ,  $n \geq 0$  there exists a positive constant  $C$ , depending on the regularity of the mesh, such that*

$$\|\mathbf{v}_h\|_{0,K}^2 \leq C (b_K \mathbf{v}_h, \mathbf{v}_h)_K, \quad (5.25)$$

Moreover, let  $b_F^S$  and  $b_F^D$  be the bubbles functions defined in (5.23) and (5.24), respectively. Then there exist positive constants  $C$ , independent of  $h_K$ , such that

$$\|\mathbf{v}_h\|_{0,F}^2 \leq C (b_F^S \mathbf{v}_h, \mathbf{v}_h)_F \quad \forall F \in \mathcal{E}_h^S, \quad (5.26)$$

$$\|\mathbf{v}_h\|_{0,F}^2 \leq C (b_F^D \mathbf{v}_h, \mathbf{v}_h)_F \quad \forall F \in \mathcal{E}_h^D, \quad (5.27)$$

$$\|b_F^S \mathcal{P}_F^S \mathbf{s}_h\|_{0,K} \leq C h_K |b_F^S \mathcal{P}_F^S \mathbf{s}_h|_{1,K} \quad \forall F \in \mathcal{E}_h^S, K \in \mathcal{T}_h^S, \quad (5.28)$$

$$\|b_F^D \mathcal{P}_F^D \mathbf{s}_h\|_{0,K} \leq C h_K |b_F^D \mathcal{P}_F^D \mathbf{s}_h|_{1,K} \quad \forall F \in \mathcal{E}_h^D, K \in \mathcal{T}_h^D, \quad (5.29)$$

$$|b_F^S \mathcal{P}_F^S \mathbf{s}_h|_{1,\omega_F^S} \leq C h_F^{-1/2} \|\mathbf{s}_h\|_{0,F} \quad \forall F \in \mathcal{E}_h^S, \quad (5.30)$$

$$|b_F^D \mathcal{P}_F^D \mathbf{s}_h|_{1,\omega_F^D} \leq C h_F^{-1/2} \|\mathbf{s}_h\|_{0,F} \quad \forall F \in \mathcal{E}_h^D, \quad (5.31)$$

for every polynomial  $\mathbf{v}_h, \mathbf{s}_h$  of degree  $n$  defined in  $K$  and  $F$ , respectively.

*Proof.* See [3, Theorem 19]. □

We denote by  $\mathbf{C}_h^S : \mathbf{H}^S \rightarrow \mathbf{H}_{h,1}^S$ ,  $\mathbf{C}_h^D : \mathbf{H}^D \rightarrow \mathbf{H}_{h,1}^D$  and  $\mathcal{C}_h^D : Q^D \rightarrow Q_{h,1}^D$ , the Clément interpolation operators. Let  $K \in \mathcal{T}_h^S \cup \mathcal{T}_h^D$  and  $F \in \mathcal{E}_h^S$ . These operators satisfy the following estimates (see [21] or [24]): There exist positive constants  $C$ , independent of  $h_K$  and the physical parameters, such that

$$\|\mathbf{v} - \mathbf{C}_h^S \mathbf{v}\|_{0,K} \leq C h_K |\mathbf{v}|_{1,\tilde{\omega}_K^S}, \quad (5.32)$$

$$\|\mathbf{v} - \mathbf{C}_h^S \mathbf{v}\|_{0,F} \leq C h_F^{1/2} |\mathbf{v}|_{1,\tilde{\omega}_F^S}, \quad (5.33)$$

$$\|\mathbf{w} - \mathbf{C}_h^D \mathbf{w}\|_{0,K} \leq C \|\mathbf{w}\|_{0,\tilde{\omega}_K^D}, \quad (5.34)$$

$$\|q - \mathcal{C}_h^D q\|_{0,K} \leq C h_K |q|_{1,\tilde{\omega}_K^D}, \quad (5.35)$$

$$\|q - \mathcal{C}_h^D q\|_{0,F} \leq C h_F^{1/2} |q|_{1,\tilde{\omega}_F^D}, \quad (5.36)$$

for all  $\mathbf{v} \in \mathbf{H}^S$ ,  $\mathbf{w} \in \mathbf{H}^D$  and  $q \in Q^D$ .

### 5.2. A residual error estimator

For each  $K \in \mathcal{T}_h^S$  and each  $F \in \mathcal{E}_{\Omega_S}$ , we define the residuals

$$\mathcal{R}_S^K := \left( \mathbf{f}_S + \nabla \cdot \boldsymbol{\sigma}_h \right) \Big|_K, \quad (5.37)$$

$$\mathcal{R}_{\Omega_S}^F := \llbracket \boldsymbol{\sigma}_h \mathbf{n}_S \rrbracket_F, \quad (5.38)$$

where  $\boldsymbol{\sigma}_h := 2\nu \boldsymbol{\varepsilon}(\mathbf{u}_{h,S}) - p_{h,S} \mathbf{I}$ . For each  $F \in \mathcal{E}_{\Gamma_h}$ , we define

$$\mathcal{R}_{\text{ENF}}^F := \left( -\mathbf{n}_S \cdot \boldsymbol{\sigma}_h \mathbf{n}_S - p_{h,D} \right) \Big|_F, \quad (5.39)$$

$$\mathcal{R}_{\text{BJS},i}^F := \left( -\kappa \mathbf{n}_S \cdot \boldsymbol{\sigma}_h \boldsymbol{\tau}_i - \kappa^{1/2} \alpha_i \mathbf{u}_{h,S} \cdot \boldsymbol{\tau}_i \right) \Big|_F, \quad 1 \leq i \leq d-1, \quad (5.40)$$

$$\mathcal{R}_{S,\Gamma}^F := \mathcal{R}_{\text{ENF}}^F \mathbf{n}_S + \sum_{i=1}^{d-1} \mathcal{R}_{\text{BJS},i}^F \boldsymbol{\tau}_i, \quad (5.41)$$

$$\mathcal{R}_{\text{MC}}^F := \left( \mathbf{u}_{h,S} \cdot \mathbf{n}_S + \mathbf{u}_{h,D} \cdot \mathbf{n}_D \right) \Big|_F. \quad (5.42)$$

Here,  $\llbracket \mathbf{v} \rrbracket_F$  represents the jump of  $\mathbf{v}$  across  $F$ . Similarly, for each  $K \in \mathcal{T}_h^D$  and  $F \in \mathcal{E}_h^D$ , we define

$$\mathcal{R}_{1,D}^K := \left( \nu \mathbf{u}_{h,D} + \kappa \nabla p_{h,D} \right) \Big|_K, \quad (5.43)$$

$$\mathcal{R}_{2,D}^K := \left( g_D - \nabla \cdot \mathbf{u}_{h,D} \right) \Big|_K, \quad (5.44)$$

$$\mathcal{R}_D^F := \begin{cases} \llbracket \mathbf{u}_{h,D} \cdot \mathbf{n}_D \rrbracket_F, & F \in \mathcal{E}_{\Omega_D}, \\ \mathbf{u}_{h,D} \cdot \mathbf{n}_D \Big|_F, & F \in \mathcal{E}_D^{\text{Neu}}, \\ 0, & F \in \mathcal{E}_D^{\text{Dir}} \cup \mathcal{E}_{\Gamma_h}. \end{cases} \quad (5.45)$$

Our residual-based error estimator is given by

$$\eta := \left\{ \sum_{K \in \mathcal{T}_h^S} \eta_{S,K}^2 + \sum_{K \in \mathcal{T}_h^D} \eta_{D,K}^2 \right\}^{1/2}, \quad (5.46)$$

where, for each  $K \in \mathcal{T}_h^S$ , we define

$$\eta_{S,K} := \left\{ \frac{\kappa}{\nu} h_K^2 \|\mathcal{R}_S^K\|_{0,K}^2 + \kappa \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 + \frac{1}{2} \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Omega_S}} \frac{\kappa}{\nu} h_F \|\mathcal{R}_{\Omega_S}^F\|_{0,F}^2 + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Gamma_h}} \frac{\kappa}{\nu} h_F \|\mathcal{R}_{S,\Gamma}^F\|_{0,F}^2 \right\}^{1/2}, \quad (5.47)$$

and for each  $K \in \mathcal{T}_h^D$ ,

$$\begin{aligned} \eta_{D,K} := & \left\{ \frac{1}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K}^2 + \nu h_K^2 \|\mathcal{R}_{2,D}^K\|_{0,K}^2 + \frac{1}{2} \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Omega_D}} \nu h_F \|\mathcal{R}_D^F\|_{0,F}^2 \right. \\ & \left. + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_D^{\text{Neu}}} \nu h_F \|\mathcal{R}_D^F\|_{0,F}^2 + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Gamma_h}} \nu h_F \|\mathcal{R}_{\text{MC}}^F\|_{0,F}^2 \right\}^{1/2}. \end{aligned} \quad (5.48)$$

**Remark 2.** Using integration by parts element-wise, we get for all  $(\mathbf{v}_S, q_S) \in \mathbf{H}^S \times Q^S$  that

$$\begin{aligned} & B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{v}_S, \mathbf{0}), (q_S, 0))) \\ &= \sum_{K \in \mathcal{T}_h^S} \kappa (\mathcal{R}_S^K, \mathbf{v}_S)_K + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa (\mathcal{R}_{S,\Gamma}^F, \mathbf{v}_S)_F + \sum_{F \in \mathcal{E}_{\Omega_S}} \kappa (\mathcal{R}_{\Omega_S}^F, \mathbf{v}_S)_F - \kappa (q_S, \nabla \cdot \mathbf{u}_{h,S})_{\Omega_S}. \end{aligned} \quad (5.49)$$

Similarly, we have that for all  $(\mathbf{v}_D, q_D) \in \mathbf{H}^D \times Q^D$

$$\begin{aligned} B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, \mathbf{v}_D), (0, q_D))) &= \sum_{K \in \mathcal{T}_h^D} \kappa (\mathcal{R}_{2,D}^K, q_D)_K + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa (\mathcal{R}_{\text{MC}}^F, q_D)_F + \sum_{F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{Neu}}} \kappa (\mathcal{R}_D^F, q_D)_F \\ &\quad - \frac{1}{2} \sum_{K \in \mathcal{T}_h^D} \left( \mathcal{R}_{1,D}^K, \mathbf{v}_D + \frac{\kappa}{\nu} \nabla q_D \right)_K. \end{aligned} \quad (5.50)$$

### 5.3. Reliability of the a posteriori error estimator

**Theorem 8.** Let  $(\mathbf{u}, p) \in \mathbf{H} \times Q$  and  $(\mathbf{u}_h, p_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$  solutions of (2.3) and (3.6), respectively. Then, there exists a positive constant  $C$ , independent of  $h$ , such that

$$\|(\mathbf{u} - \mathbf{u}_h, p - p_h)\| \leq C \left\{ \eta + \left[ \sum_{K \in \mathcal{T}_h^S} \frac{h_K^4}{\nu^3 \kappa} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 \right]^{1/2} \right\}.$$

*Proof.* Let  $(\mathbf{v}, q) := ((\mathbf{v}_S, \mathbf{v}_D), (q_S, q_D)) \in \mathbf{H} \times Q$  an arbitrary element, and  $\mathbf{v}_{1,S} := \mathbf{C}_h^S \mathbf{v}_S$ . Applying (5.49)

to  $(\mathbf{v}_S - \mathbf{v}_{1,S}, q_S)$ , using interpolation properties, and Cauchy-Schwarz inequalities, we get

$$\begin{aligned}
& B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{v}_S - \mathbf{v}_{1,S}, \mathbf{0}), (q_S, 0))) \\
& \leq \sum_{K \in \mathcal{T}_h^S} \kappa \|\mathcal{R}_S^K\|_{0,K} \|\mathbf{v}_S - \mathbf{v}_{1,S}\|_{0,K} + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} \|\mathbf{v}_S - \mathbf{v}_{1,S}\|_{0,F} \\
& \quad + \sum_{F \in \mathcal{E}_{\Omega_S}} \kappa \|\mathcal{R}_{\Omega_S}^F\|_{0,F} \|\mathbf{v}_S - \mathbf{v}_{1,S}\|_{0,F} + \sum_{K \in \mathcal{T}_h^S} \kappa \|q_S\|_{0,K} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K} \\
& \leq C \left\{ \sum_{K \in \mathcal{T}_h^S} \kappa h_K \|\mathcal{R}_S^K\|_{0,K} |\mathbf{v}_S|_{1,\tilde{\omega}_K} + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa h_F^{1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} |\mathbf{v}_S|_{1,\tilde{\omega}_F^S} \right. \\
& \quad \left. + \sum_{F \in \mathcal{E}_{\Omega_S}} \kappa h_F^{1/2} \|\mathcal{R}_{\Omega_S}^F\|_{0,F} |\mathbf{v}_S|_{1,\tilde{\omega}_F^S} + \sum_{K \in \mathcal{T}_h^S} \kappa \|q_S\|_{0,K} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K} \right\} \\
& \leq C \left\{ \sum_{K \in \mathcal{T}_h^S} \frac{\kappa}{\nu} h_K^2 \|\mathcal{R}_S^K\|_{0,K}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} \frac{\kappa}{\nu} h_F \|\mathcal{R}_{S,\Gamma}^F\|_{0,F}^2 + \sum_{F \in \mathcal{E}_{\Omega_S}} \frac{\kappa}{\nu} h_F \|\mathcal{R}_{\Omega_S}^F\|_{0,F}^2 + \sum_{K \in \mathcal{T}_h^S} \kappa \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 \right\}^{1/2} \\
& \quad \left\{ \sum_{K \in \mathcal{T}_h^S} \kappa \nu |\mathbf{v}_S|_{1,\tilde{\omega}_K}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa \nu |\mathbf{v}_S|_{1,\tilde{\omega}_F^S}^2 + \sum_{F \in \mathcal{E}_{\Omega_S}} \kappa \nu |\mathbf{v}_S|_{1,\tilde{\omega}_F^S}^2 + \sum_{K \in \mathcal{T}_h^S} \kappa \|q_S\|_{0,K}^2 \right\}^{1/2} \\
& \leq C \left\{ \sum_{K \in \mathcal{T}_h^S} \eta_{S,K}^2 \right\}^{1/2} \|(\mathbf{v}, q)\|. \tag{5.51}
\end{aligned}$$

On the other hand, let  $\mathbf{v}_{1,D} := \mathcal{C}_h^D \mathbf{v}_D$  and  $q_{1,D} := \mathcal{C}_h^D q_D$ . Using a similar procedure, we have

$$\begin{aligned}
& B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, \mathbf{v}_D - \mathbf{v}_{1,D}), (0, q_D - q_{1,D}))) \\
& = \sum_{K \in \mathcal{T}_h^D} \kappa (\mathcal{R}_{2,D}^K, q_D - q_{1,D})_K + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa (\mathcal{R}_{MC}^F, q_D - q_{1,D})_F + \sum_{F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{Neu}}} \kappa (\mathcal{R}_D^F, q_D - q_{1,D})_F \\
& \quad - \frac{1}{2} \sum_{K \in \mathcal{T}_h^D} \left( \mathcal{R}_{1,D}^K, (\mathbf{v}_D - \mathbf{v}_{1,D}) + \frac{\kappa}{\nu} \nabla (q_D - q_{1,D}) \right)_K \\
& \leq \sum_{K \in \mathcal{T}_h^D} \kappa \|\mathcal{R}_{2,D}^K\|_{0,K} \|q_D - q_{1,D}\|_{0,K} + \sum_{F \in \mathcal{E}_{\Gamma_h}} \kappa \|\mathcal{R}_{MC}^F\|_{0,F} \|q_D - q_{1,D}\|_{0,F} + \sum_{F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{Neu}}} \kappa \|\mathcal{R}_D^F\|_{0,F} \|q_D - q_{1,D}\|_{0,F} \\
& \quad + \frac{1}{2} \sum_{K \in \mathcal{T}_h^D} \|\mathcal{R}_{1,D}^K\|_{0,K} \left[ \|\mathbf{v}_D - \mathbf{v}_{1,D}\|_{0,K} + \frac{\kappa}{\nu} |q_D - q_{1,D}|_{1,K} \right] \\
& \leq C \left\{ \sum_{K \in \mathcal{T}_h^D} \nu h_K^2 \|\mathcal{R}_{2,D}^K\|_{0,K}^2 + \sum_{F \in \mathcal{E}_{\Gamma_h}} \nu h_F \|\mathcal{R}_{MC}^F\|_{0,F}^2 + \sum_{F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{Neu}}} \nu h_F \|\mathcal{R}_D^F\|_{0,F}^2 + \sum_{K \in \mathcal{T}_h^D} \frac{1}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K}^2 \right\}^{1/2} \\
& \quad \left\{ \nu \|\mathbf{v}_D\|_{0,\Omega_D} + \frac{\kappa^2}{\nu} |q_D|_{1,\Omega_D} \right\}^{1/2} \\
& \leq C \left\{ \sum_{K \in \mathcal{T}_h^D} \eta_{D,K}^2 \right\}^{1/2} \|(\mathbf{v}, q)\|. \tag{5.52}
\end{aligned}$$

Let  $\tilde{\mathbf{v}}_h := (\mathbf{v}_{1,S}, \mathbf{v}_{1,D})$  and  $\tilde{q}_h := (0, q_{1,D})$ . Combining (5.51) and (5.52), we obtain

$$B((\mathbf{u} - \mathbf{u}_h, p - p_h); (\mathbf{v} - \tilde{\mathbf{v}}_h, q - \tilde{q}_h)) \leq C \eta \|(\mathbf{v}, q)\|. \tag{5.53}$$



Similar arguments proves that

$$B((\mathbf{u} - \mathbf{u}_h, p - p_h); (\tilde{\mathbf{v}}_h, \tilde{q}_h)) \leq C \left\{ \eta + \left[ \sum_{K \in \mathcal{T}_h^S} \frac{h_K^4}{\nu^3 \kappa} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 \right]^{1/2} \right\} \|(\mathbf{v}, q)\|. \quad (5.54)$$

Therefore, from (5.53), (5.54) and Lemma 14, we arrive at

$$\begin{aligned} \|(\mathbf{u} - \mathbf{u}_h, p - p_h)\| &\leq C \sup_{(\mathbf{v}, q) \in \mathbf{H} \times Q} \frac{B((\mathbf{u} - \mathbf{u}_h, p - p_h); (\mathbf{v}, q))}{\|(\mathbf{v}, q)\|} \\ &\leq C \left\{ \eta + \left[ \sum_{K \in \mathcal{T}_h^S} \frac{h_K^4}{\nu^3 \kappa} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 \right]^{1/2} \right\}, \end{aligned}$$

which conclude the proof of the result.  $\square$

**Remark 3.** *The term*

$$T := \left\{ \sum_{K \in \mathcal{T}_h^S} \frac{h_K^4}{\nu^3 \kappa} \|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K}^2 \right\}^{1/2}, \quad (5.55)$$

appearing in Theorem 8, is only present if  $\theta > 0$  in the definition of  $B_{\text{stab}}$  in (3.7). In that case  $T$  is, at least,  $\mathcal{O}(h^2)$  asymptotically, i.e. is a high order term compared to  $\eta$ .

#### 5.4. Efficiency of the a posteriori error estimator

In order to prove the efficiency of error estimate  $\eta$ , we need some preliminary results.

**Lemma 9.** *For all  $K \in \mathcal{T}_h^S$ , there exist two positive constants  $C_1$  and  $C_2$ , independent of  $\nu$ ,  $\kappa$  and  $h$ , such that*

$$\|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K} \leq C_1 \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K},$$

and

$$\sqrt{\frac{\kappa}{\nu}} h_K \|\mathcal{R}_S^K\|_{0,K} \leq C_2 \max \left\{ 1, \frac{1}{\sqrt{\nu}} \right\} \left\{ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K} + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,K} \right\}.$$

*Proof.* Let  $b_K$  be the bubble function associated to  $K \in \mathcal{T}_h^S$ . Since  $\nabla \cdot \mathbf{u}_S \in L^2(\Omega_S)$  and using (2.3), we have that  $\nabla \cdot \mathbf{u}_S = 0$  a.e in  $\Omega_S$ . Then it is clear that

$$\|\nabla \cdot \mathbf{u}_{h,S}\|_{0,K} \leq C_1 \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K},$$

for all  $K \in \mathcal{T}_h^S$ . Now, let us define  $\mathbf{b}_K := b_K \mathcal{R}_S^K$ . From (5.49), (2.4) and an inverse inequality, we obtain

$$\begin{aligned} \kappa (\mathcal{R}_S^K, \mathbf{b}_K)_K &= \sum_{K' \in \mathcal{T}_h^S} \kappa (\mathcal{R}_S^{K'}, \mathbf{b}_K)_{K'} = B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{b}_K, \mathbf{0}), (0, 0))) \\ &= 2\nu \kappa (\varepsilon(\mathbf{u}_S - \mathbf{u}_{h,S}), \varepsilon(\mathbf{b}_K))_K - \kappa (p_S - p_{h,S}, \nabla \cdot \mathbf{b}_K)_K \\ &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K} + \kappa \|p_S - p_{h,S}\|_{0,K} \right\} \|\mathbf{b}_K\|_{1,K} \\ &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K} + \kappa \|p_S - p_{h,S}\|_{0,K} \right\} h_K^{-1} \|\mathbf{b}_K\|_{0,K} \\ &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K} + \kappa \|p_S - p_{h,S}\|_{0,K} \right\} h_K^{-1} \|\mathcal{R}_S^K\|_{0,K}. \end{aligned}$$

Next, using Theorem 7, it follows, for all  $K \in \mathcal{T}_h^S$ , that

$$\kappa h_K \|\mathcal{R}_S^K\|_{0,K} \leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,K} + \kappa \|p_S - p_{h,S}\|_{0,K} \right\}, \quad (5.56)$$

hence, multiplying by  $\sqrt{\frac{1}{\kappa \nu}}$ , we arrive to the desired result.  $\square$

For the next result, we will consider the bubble function  $b_F^S$  defined for all  $F \in \mathcal{E}_{\Omega_S} \cup \mathcal{E}_{\Gamma_h}$ .

**Lemma 10.** *Let  $F \in \mathcal{E}_{\Gamma_h}$ . Then, there exist a positive constant  $C_3$ , independent of  $\nu$ ,  $\kappa$ , and  $h$ , such that:*

$$\begin{aligned} \sqrt{\frac{\kappa}{\nu}} h_F^{1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} \leq C_3 \max \left\{ 1, \frac{1}{\sqrt{\nu}}, \frac{1}{\nu}, \frac{h_F}{\nu\sqrt{\kappa}}, \frac{h_F}{\sqrt{\kappa}} \right\} \left\{ \sqrt{\nu\kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} \right. \\ \left. + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} + \frac{\kappa}{\sqrt{\nu}} \|p_D - p_{h,D}\|_{1,\omega_F^D} \right\}. \end{aligned} \quad (5.57)$$

Furthermore, for  $F \in \mathcal{E}_{\Omega_S}$  there holds

$$\sqrt{\frac{\kappa}{\nu}} h_F^{1/2} \|\mathcal{R}_{\Omega_S}^F\|_{0,F} \leq C_4 \max \left\{ 1, \frac{1}{\sqrt{\nu}} \right\} \left\{ \sqrt{\nu\kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} \right\}, \quad (5.58)$$

where  $C_4$  is a positive constant independent of  $h$ .

*Proof.* Applying Theorem 7, (5.49), the definition of  $B$  and Cauchy–Schwarz inequality, we obtain

$$\begin{aligned} \kappa \|\mathcal{R}_{S,\Gamma}^F\|_{0,F}^2 &\leq C \kappa (\mathcal{R}_{S,\Gamma}^F, b_F^S \mathcal{R}_{S,\Gamma}^F)_F \\ &\leq C \left\{ B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F, \mathbf{0}), (0, 0))) - \sum_{K \in \omega_F^S} \kappa (\mathcal{R}_S^K, b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F)_K \right\} \\ &= C \left\{ \nu \kappa (\varepsilon(\mathbf{u}_S - \mathbf{u}_{h,S}), \varepsilon(b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F))_{\omega_F^S} - \kappa (p_S - p_{h,S}, \nabla \cdot (b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F))_{\omega_F^S} \right. \\ &\quad \left. + \kappa^{1/2} \sum_{i=1}^{d-1} \alpha_i ((\mathbf{u}_S - \mathbf{u}_{h,S}) \cdot \boldsymbol{\tau}_i, b_F^S \mathcal{R}_{S,\Gamma}^F \cdot \boldsymbol{\tau}_i)_F + \kappa (p_D - p_{h,D}, b_F^S \mathcal{R}_{S,\Gamma}^F \cdot \mathbf{n}_S)_F - \sum_{K \in \omega_F^S} \kappa (\mathcal{R}_S^K, b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F)_K \right\} \\ &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} |b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F|_{1,\omega_F^S} + \kappa \|p_S - p_{h,S}\|_{0,\omega_F^S} |b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F|_{1,\omega_F^S} + \kappa^{1/2} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{0,F} \|b_F^S \mathcal{R}_{S,\Gamma}^F\|_{0,F} \right. \\ &\quad \left. + \kappa \|p_D - p_{h,D}\|_{0,F} \|b_F^S \mathcal{R}_{S,\Gamma}^F\|_{0,F} + \sum_{K \in \omega_F^S} \kappa \|\mathcal{R}_S^K\|_{0,K} |b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F|_{0,K} \right\}. \end{aligned}$$

On the other hand, from Lemma 4, (5.28), Theorem 7, mesh regularity, and (5.56), we have

$$\begin{aligned} \kappa \|\mathcal{R}_{S,\Gamma}^F\|_{0,F}^2 &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} h_F^{-1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} + \kappa \|p_S - p_{h,S}\|_{0,\omega_F^S} h_F^{-1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} \right. \\ &\quad \left. + \kappa^{1/2} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{0,F} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} + \kappa \|p_D - p_{h,D}\|_{0,F} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} + \sum_{K \in \omega_F^S} \kappa h_K \|\mathcal{R}_S^K\|_{0,K} |b_F^S \mathcal{P}_F^S \mathcal{R}_{S,\Gamma}^F|_{1,\omega_F^S} \right\} \\ &\leq C \left\{ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \kappa \|p_S - p_{h,S}\|_{0,\omega_F^S} + \kappa^{1/2} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{0,\omega_F^S} \right. \\ &\quad \left. + \kappa^{1/2} h_F |\mathbf{u}_S - \mathbf{u}_{h,S}|_{1,\omega_F^S} + \kappa \|p_D - p_{h,D}\|_{0,\omega_F^D} + \kappa h_F |p_D - p_{h,D}|_{1,\omega_F^D} \right\} h_F^{-1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F}, \end{aligned}$$

thus

$$\begin{aligned} \sqrt{\frac{\kappa}{\nu}} h_F^{1/2} \|\mathcal{R}_{S,\Gamma}^F\|_{0,F} \leq C \max \left\{ 1, \frac{1}{\sqrt{\nu}}, \frac{1}{\nu}, \frac{h_F}{\nu\sqrt{\kappa}}, \frac{h_F}{\sqrt{\kappa}} \right\} & \left\{ \sqrt{\nu\kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} \right. \\ & \left. + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} + \frac{\kappa}{\sqrt{\nu}} \|p_D - p_{h,D}\|_{1,\omega_F^S} \right\}. \end{aligned}$$

Similar arguments, applied to  $F \in \mathcal{E}_{\Omega_S}$ , show that

$$\begin{aligned} \kappa \|\mathcal{R}_{\Omega_S}^F\|_{0,F}^2 & \leq C \kappa (\mathcal{R}_{\Omega_S}^F, b_F^S \mathcal{R}_{\Omega_S}^F)_F \\ & \leq C \left[ B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((b_F^S \mathcal{P}_F^S \mathcal{R}_{\Omega_S}^F, \mathbf{0}), (0, 0))) - \sum_{K \in \omega_F^S} \kappa (\mathcal{R}_S^K, b_F^S \mathcal{P}_F^S \mathcal{R}_{\Omega_S}^F)_K \right] \\ & \leq C \left[ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \kappa \|p_S - p_{h,S}\|_{0,\omega_F^S} \right] |b_F^S \mathcal{P}_F^S \mathcal{R}_{\Omega_S}^F|_{1,\omega_F^S} \\ & \leq C \left[ \nu \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \kappa \|p_S - p_{h,S}\|_{0,\omega_F^S} \right] h_F^{-1/2} \|\mathcal{R}_{\Omega_S}^F\|_{0,F}, \end{aligned}$$

and the result follows.  $\square$

To prove the efficiency of  $\eta$ , we will need the next result

**Lemma 11.** *For all  $K \in \mathcal{T}_h^D$ , there exist two positive constants  $C_5$  and  $C_6$ , independent of  $\nu$ ,  $\kappa$ , and  $h$ , such that*

$$\frac{1}{\sqrt{\nu}} \|\mathcal{R}_{1,D}^K\|_{0,K} \leq C_5 \left\{ \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,K} \right\}, \quad (5.59)$$

and

$$\sqrt{\nu} h_K \|\mathcal{R}_{2,D}^K\|_{0,K} \leq C_6 \left\{ \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,K} \right\}. \quad (5.60)$$

*Proof.* From Theorem 7, (5.50) and Cauchy–Schwarz inequality, it follows that

$$\begin{aligned} \|\mathcal{R}_{1,D}^K\|_{0,K}^2 & \leq C (\mathcal{R}_{1,D}^K, b_K \mathcal{R}_{1,D}^K)_K \leq C \sum_{K' \in \mathcal{T}_h^D} (R_{1,D}^{K'}, b_K \mathcal{R}_{1,D}^K)_{K'} \\ & \leq C B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, -b_K \mathcal{R}_{1,D}^K), (0, 0))) \\ & \leq C \left\{ \nu (\mathbf{u}_D - \mathbf{u}_{h,D}, -b_K \mathcal{R}_{1,D}^K)_K + \kappa (\nabla(p_D - p_{h,D}), -b_K \mathcal{R}_{1,D}^K)_K + \right. \\ & \quad \left. (\nu (\mathbf{u}_D - \mathbf{u}_{h,D}) + \kappa \nabla(p_D - p_{h,D}), b_K \mathcal{R}_{1,D}^K)_K \right\} \\ & \leq C \left\{ \nu \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \kappa |p_D - p_{h,D}|_{1,K} \right\} \|\mathcal{R}_{1,D}^K\|_{0,K}, \end{aligned}$$

which proves (5.59). We can now proceed analogously, using Theorem 7, (5.50), (5.59) and Cauchy–Schwarz

inequality, to conclude that

$$\begin{aligned}
& \kappa \|\mathcal{R}_{2,D}^K\|_{0,K}^2 \leq C \kappa (\mathcal{R}_{2,D}^K, b_K \mathcal{R}_{2,D}^K)_K \leq C \sum_{K' \in \mathcal{T}_h^D} \kappa (R_{2,D}^{K'}, b_{K'} \mathcal{R}_{2,D}^{K'})_{K'} \\
& \leq C \left\{ B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, \mathbf{0}), (0, b_K \mathcal{R}_{2,D}^K))) + \left( \mathcal{R}_{1,D}^K, \frac{\kappa}{\nu} \nabla (b_K \mathcal{R}_{2,D}^K) \right)_K \right\} \\
& \leq C \left\{ -\kappa (\mathbf{u}_D - \mathbf{u}_{h,D}, \nabla (b_K \mathcal{R}_{2,D}^K))_K + \left( \nu (\mathbf{u}_D - \mathbf{u}_{h,D}) + \kappa \nabla (p_D - p_{h,D}), \frac{\kappa}{\nu} \nabla (b_K \mathcal{R}_{2,D}^K) \right)_K + \right. \\
& \quad \left. \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} |b_K \mathcal{R}_{2,D}^K|_{1,K} \right\} \\
& \leq C \left\{ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right\} |b_K \mathcal{R}_{2,D}^K|_{1,K} \\
& \leq C \left\{ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right\} h_K^{-1} \|\mathcal{R}_{2,D}^K\|_{0,K},
\end{aligned}$$

thus, using (5.59) we prove (5.60).  $\square$

**Lemma 12.** For all  $F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{neu}}$ , there exists two positive constants  $C_7$  and  $C_8$ , independent of  $\nu$ ,  $\kappa$ , and  $h$ , such that

$$\sqrt{\nu} h_F^{1/2} \|\mathcal{R}_D^F\|_{0,F} \leq C_7 \left\{ \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,\omega_F^D} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,\omega_F^D} \right\}. \quad (5.61)$$

Furthermore, for all  $F \in \mathcal{E}_{\Gamma_h}$  the following estimate holds:

$$\sqrt{\nu} h_F^{1/2} \|\mathcal{R}_{\text{MC}}^F\|_{0,F} \leq C_8 \max \left\{ 1, \frac{1}{\sqrt{\kappa}}, \frac{h_F}{\sqrt{\kappa}} \right\} \left\{ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,\omega_F^D} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,\omega_F^D} \right\}. \quad (5.62)$$

*Proof.* We use the same ideas that in previous results, but now for  $F \in \mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{neu}}$ . In fact, using Theorem 7, (5.50), and Cauchy-Schwarz inequality, we obtain

$$\begin{aligned}
& \kappa \|\mathcal{R}_D^F\|_{0,F}^2 \leq C \kappa (\mathcal{R}_D^F, b_F^D \mathcal{R}_D^F)_F \\
& \leq C \left\{ B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, \mathbf{0}), (0, b_F^D \mathcal{P}_F^D \mathcal{R}_D^F))) - \sum_{K \in \omega_F^D} \kappa (\mathcal{R}_{2,D}^K, b_F^D \mathcal{P}_F^D \mathcal{R}_D^F)_K + \frac{1}{2} \sum_{K \in \omega_F^D} \left( \mathcal{R}_{1,D}^K, \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_D^F) \right)_K \right\} \\
& \leq C \left\{ -\kappa (\mathbf{u}_D - \mathbf{u}_{h,D}, \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_D^F))_{\omega_F^D} + \left( \nu (\mathbf{u}_D - \mathbf{u}_{h,D}) + \kappa \nabla (p_D - p_{h,D}), \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_D^F) \right)_{\omega_F^D} \right. \\
& \quad \left. - \sum_{K \in \omega_F^D} \kappa (\mathcal{R}_{2,D}^K, b_F^D \mathcal{P}_F^D \mathcal{R}_D^F)_K + \frac{1}{2} \sum_{K \in \omega_F^D} \left( \mathcal{R}_{1,D}^K, \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_D^F) \right)_K \right\} \\
& \leq C \sum_{K \in \omega_F^D} \left[ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \kappa h_K \|\mathcal{R}_{2,D}^K\|_{0,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right] |b_F^D \mathcal{P}_F^D \mathcal{R}_D^F|_{1,\omega_F^D} \\
& \leq C \sum_{K \in \omega_F^D} \left[ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \kappa h_K \|\mathcal{R}_{2,D}^K\|_{0,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right] h_F^{-1/2} \|\mathcal{R}_D^F\|_{0,F},
\end{aligned}$$

which, combined with Lemma 11, leads to (5.61). Again, using Theorem 7, and Cauchy–Schwarz inequality, we get that

$$\begin{aligned}
& \kappa \|\mathcal{R}_{\text{MC}}^F\|_{0,F}^2 \leq C \kappa (\mathcal{R}_{\text{MC}}^F, b_F^D \mathcal{R}_{\text{MC}}^F)_F \\
& \leq C \left\{ B((\mathbf{u} - \mathbf{u}_h, p - p_h); ((\mathbf{0}, \mathbf{0}), (0, b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F))) - \sum_{K \in \omega_F^D} \kappa (\mathcal{R}_{2,D}^K, b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F)_K + \frac{1}{2} \sum_{K \in \omega_F^D} \left( \mathcal{R}_{1,D}^K, \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F) \right)_K \right\} \\
& \leq C \left\{ -\kappa (\mathbf{u}_D - \mathbf{u}_{h,D}, \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F))_{\omega_F^D} - \kappa ((\mathbf{u}_S - \mathbf{u}_{h,S}) \cdot \mathbf{n}_S, b_F^D \mathcal{R}_{\text{MC}}^F)_F \right. \\
& \quad \left. + \left( \nu (\mathbf{u}_D - \mathbf{u}_{h,D}) + \kappa \nabla (p_D - p_{h,D}), \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F) \right)_{\omega_F^D} \right. \\
& \quad \left. - \sum_{K \in \omega_F^D} \kappa (\mathcal{R}_{2,D}^K, b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F)_K + \sum_{K \in \omega_F^D} \left( \mathcal{R}_{1,D}^K, \frac{\kappa}{\nu} \nabla (b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F) \right)_K \right\} \\
& \leq C \left\{ \sum_{K \in \omega_F^D} \left[ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \kappa h_K \|\mathcal{R}_{2,D}^K\|_{0,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right] |b_F^D \mathcal{P}_F^D \mathcal{R}_{\text{MC}}^F|_{1,\omega_F^D} \right. \\
& \quad \left. + \kappa \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{0,F} \|b_F^D \mathcal{R}_{\text{MC}}^F\|_{0,F} \right\} \\
& \leq C \left\{ \sum_{K \in \omega_F^D} \left[ \kappa \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa^2}{\nu} |p_D - p_{h,D}|_{1,K} + \kappa h_K \|\mathcal{R}_{2,D}^K\|_{0,K} + \frac{\kappa}{\nu} \|\mathcal{R}_{1,D}^K\|_{0,K} \right] \right. \\
& \quad \left. + \kappa h_F^{1/2} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{0,F} \right\} h_F^{-1/2} \|\mathcal{R}_{\text{MC}}^F\|_{0,F},
\end{aligned}$$

thus (5.62) is obtained by lemmas 4 and 11.  $\square$

**Theorem 13.** *Let  $(\mathbf{u}, p) \in \mathbf{H} \times Q$  and  $(\mathbf{u}_h, p_h) \in \mathbf{H}_{h,k} \times Q_{h,k}$  solutions of (2.3) and (3.6), respectively. Then, there is two positive constants  $C_9$  and  $C_{10}$ , independent of  $h$ , such that*

$$\begin{aligned}
\eta_{S,K} & \leq C_9 \max \left\{ 1, \frac{1}{\sqrt{\nu}}, \frac{1}{\nu}, \frac{h_F}{\nu \sqrt{\kappa}}, \frac{h_F}{\sqrt{\kappa}} \right\} \left\{ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} \right. \\
& \quad \left. + \frac{\kappa}{\sqrt{\nu}} \|p_D - p_{h,D}\|_{1,\omega_F^D} + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Omega_S}} \left[ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} \right] \right. \\
& \quad \left. + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Gamma_h}} \left[ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\kappa} \|p_S - p_{h,S}\|_{0,\omega_F^S} + \frac{\kappa}{\sqrt{\nu}} \|p_D - p_{h,D}\|_{1,\omega_F^D} \right] \right\},
\end{aligned}$$

for all  $K \in \mathcal{T}_h^S$ , and

$$\begin{aligned}
\eta_{D,K} & \leq C_{10} \max \left\{ 1, \frac{1}{\sqrt{\kappa}}, \frac{h_K}{\sqrt{\kappa}} \right\} \left\{ \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,K} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,K} \right. \\
& \quad \left. + \sum_{F \in \mathcal{E}(K) \cap (\mathcal{E}_{\Omega_D} \cup \mathcal{E}_D^{\text{neu}})} \left[ \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,\omega_F^D} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,\omega_F^D} \right] \right. \\
& \quad \left. + \sum_{F \in \mathcal{E}(K) \cap \mathcal{E}_{\Gamma_h}} \left[ \sqrt{\nu \kappa} \|\mathbf{u}_S - \mathbf{u}_{h,S}\|_{1,\omega_F^S} + \sqrt{\nu} \|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,\omega_F^D} + \frac{\kappa}{\sqrt{\nu}} |p_D - p_{h,D}|_{1,\omega_F^D} \right] \right\},
\end{aligned}$$

for all  $K \in \mathcal{T}_h^D$ .

*Proof.* The proof follows from a direct application of lemmas 9–12.  $\square$

## 6. Numerical experiments

In this section we present three numerical examples to show the quality of our stabilized adaptive scheme. All the examples were performed using the python libraries `Multiphenics` and `FEniCS` [34].

Recall that we use the notation  $\mathbf{H}_{h,k} \times Q_{h,k}$  to mean that the velocity and the pressure, of Stokes and Darcy, are approximated using piecewise continuous polynomials of total degree at most  $k$ .

We define the effectivity index  $E$  as follows

$$E := \frac{\eta}{\|(\mathbf{u} - \mathbf{u}_h, p - p_h)\|}.$$

Thus our adaptive algorithm for the Stokes–Darcy coupled problem is given by

---

### Algorithm 1 Adaptivity procedure

---

**Require:**  $\vartheta \in (0, 1)$  and a coarse mesh  $\mathcal{T}_h = \mathcal{T}_h^S \cup \mathcal{T}_h^D$ .

- 1: Solve the stabilized discrete scheme (3.6) on the current mesh.
  - 2: For each  $K \in \mathcal{T}_h$ , compute the local error indicators  $\eta_{S,K}$  or  $\eta_{D,K}$  given by (5.47) and (5.48), respectively.
  - 3: Given  $K \in \mathcal{T}_h$  such that  $\eta_K \geq \vartheta \max_{K' \in \mathcal{T}_h} \eta_{K'}$ , mark  $K$  and generate a new mesh  $\mathcal{T}_h$  refining the marked elements.
  - 4: If the stop criterion is not satisfied, go to step 1.
- 

#### 6.1. A smooth solution in two dimensions

In this example (see [5]),  $\Omega_S := ]0, 1/2[ \times ]0, 1[$  and  $\Omega_D := ]1/2, 1[ \times ]0, 1[$ , the interface between  $\Omega_S$  and  $\Omega_D$  is given by  $\Gamma := \{(1/2, y) \in \mathbb{R}^2 : 0 < y < 1\}$  and, on the Beavers–Joseph–Saffman interface condition,  $\alpha_1 = 1$ . The boundary  $\Gamma_D$  is divided on  $\Gamma_D^{\text{Dir}} := \{(1, y) \in \mathbb{R}^2 : 0 < y < 1\} \cup \{(x, 1) \in \mathbb{R}^2 : 1/2 < x < 1\}$  and  $\Gamma_D^{\text{Neu}} := \{(x, 0) \in \mathbb{R}^2 : 1/2 < x < 1\}$  as shown in the Figure 6.1.




Figure 2: Configuration of the computational domain  $\Omega$ .

The data  $\mathbf{f}_S$  and  $g_D$  are such that the exact solution of our problem is given by

$$\begin{aligned} \mathbf{u}_S(x, y) &:= (xy(1-y), x^2(1-y)\sin y), & p_S(x, y) &:= 12x^2e^y, \\ \mathbf{u}_D(x, y) &:= (2xy(1-y)(1-x), xy^2(1-y)), & p_D(x, y) &:= 16xy^3 - e - 2. \end{aligned}$$

Finally, we use  $\nu = 1, 10^{-2}$  and  $\kappa = 1$ . The approximation subspaces used for this example are  $\mathbf{H}_{h,k} \times Q_{h,k}$  with  $k = 1$  and  $k = 2$ .

Figures 3 – 6, show the orders of convergence for our proposed scheme in a quasi-uniform refinement. We can observe that the error norms  $\|\mathbf{u}_D - \mathbf{u}_{h,D}\|_{0,\Omega_D}$  and  $\|p_S - p_{h,S}\|_{0,\Omega_S}$  present a better behavior than predicted by Theorem 6; we assume that this is because the solutions of this example are smooth. In Tables 1–2, we present the behavior of the residual error estimator  $\eta$  defined through (5.46). Note that the error estimator has a quite good quality reflected on the fact that effectivity indexes are close to one.




Figure 3: Convergence of the velocities (left) and pressures (right). Here  $\nu = \kappa = 1$  and the interpolation order is  $k = 1$ .




Figure 4: Convergence of the velocities (left) and pressures (right). Here  $\nu = 10^{-2}$ ,  $\kappa = 1$  and the interpolation order is  $k = 1$ .




Figure 5: Convergence of the velocities (left) and pressures (right). Here  $\nu = \kappa = 1$  and the interpolation order is  $k = 2$ .




Figure 6: Convergence of the velocities (left) and pressures (right). Here  $\nu = 10^{-2}$ ,  $\kappa = 1$  and the interpolation order is  $k = 2$ .

$k$	$h$	$\ (\mathbf{u} - \mathbf{u}_h, p - p_h)\ $	order	$\eta$	order	$E$
1	0.0625	0.5223353	–	0.5199705	–	0.9954727
	0.03125	0.2601458	1.0056559	0.2605913	0.9966406	1.0017128
	0.015625	0.1299038	1.0018765	0.1304250	0.9985686	1.0040122
	0.0078125	0.0649220	1.0006647	0.0652396	0.9994017	1.0048916
	0.0039062	0.0324552	1.0002562	0.0326256	0.9997438	1.0052486
2	0.0625	0.0066269	–	0.0077845	–	1.1746854
	0.03125	0.0016570	1.9997191	0.0019468	1.9994783	1.1748814
	0.015625	0.0004144	1.9995227	0.0004869	1.9994128	1.1749709
	0.0078125	0.0001036	1.9996771	0.0001218	1.9996247	1.1750136

Table 1: Orders of convergence and effectivity index. Here  $\nu = \kappa = 1$ .



$k$	$h$	$\ (\mathbf{u} - \mathbf{u}_h, p - p_h)\ $	order	$\eta$	order	$E$
1	0.0625	5.2212942	–	5.1711817	–	0.9904023
	0.03125	2.6006875	1.0055144	2.5912903	0.9968233	0.9963867
	0.015625	1.2986902	1.0018357	1.2968388	0.9986714	0.9985745
	0.0078125	0.6490518	1.0006519	0.6486641	0.9994556	0.9994028
	0.0039062	0.3244693	1.0002517	0.3243835	0.9997714	0.9997356
2	0.0625	0.0662295	–	0.0660592	–	0.9974282
	0.03125	0.0165606	1.9997172	0.0165099	2.0004243	0.9969395
	0.015625	0.0041415	1.9995226	0.0041278	1.9999050	0.9966753
	0.0078125	0.0010356	1.9996773	0.0010320	1.9998723	0.9965406

Table 2: Orders of convergence and effectivity index. Here  $\nu = 10^{-2}$ ,  $\kappa = 1$ .

## 6.2. Simulation of a micro-filtration processes

Inspired by [30], we build a test case based on the simulation of a micro-filtration process that satisfies our interface conditions. Let  $\Omega_S := ]-1/2, 1[ \times ]0, 1/2[$ ,  $\Omega_D := ]0, 1/2[ \times ]-1/4, 0[$ , and the different boundaries conditions are represented in Figure 7. We consider  $\nu = \kappa = 10^{-2}$ ,  $\alpha_1 = 10^{-2}$  on the Beavers–Joseph–Saffman condition, and the data  $\mathbf{f}_S = (0, 0)$  and  $g_D = 0$ .




Figure 7: Boundaries conditions for simulation of a micro-filtration processes.

In this test case we do not have an analytical solution, thus we compare our adapted solution with a numerical solution obtained in a highly uniform refined mesh of 560.000 elements. In Figure 8 we show that our adaptive scheme refine the mesh in the corners of the domain, as well as, at the out flow boundary.

In Figures 9 and 10, we depict the pressure and velocity magnitude isolines, both in the case of our adapted mesh and in the reference solution. Note that we get a good agreement in both solutions.




Figure 8: Initial mesh (left), with 224 elements, and final adapted mesh (right), with 8.674 elements. Here  $k = 1$ .




Figure 9: Isolines of the pressure on the adapted mesh (left) and on the highly uniform refined mesh. Here  $k = 1$ .




Figure 10: Isolines of the velocity magnitude on the adapted mesh (left) and on the highly uniform refined mesh. Here  $k = 1$ .

### 6.3. A analytical solution in three dimensions

The final test illustrates the performance of our adaptive stabilized finite element to simulate a 3D problem. In this example  $\Omega := (0, 1) \times (-1/2, 1/2) \times (0, 1)$ , with  $\Omega_S := (0, 1) \times (-1/2, 0) \times (0, 1)$ ,  $\Omega_D := (0, 1) \times (0, 1/2) \times (0, 1)$ , the interface  $\Gamma := \overline{\Omega_S} \cap \overline{\Omega_D}$ ,  $\nu = \kappa = 1$ , and  $\alpha_1 = \alpha_2 = 1$ . The boundary  $\Gamma_D$  is split as follows

$$\Gamma_D^{\text{Neu}} := \{(x, y, 0) \in \mathbb{R}^3 : 0 < x < 1, 0 < y < 1/2\}, \quad \Gamma_D^{\text{Dir}} := \Gamma_D \setminus \overline{\Gamma_D^{\text{Neu}}},$$

as it is shown in Figure 11. The data  $\mathbf{f}_S$  and  $g_D$  are such that the exact solution is given by

$$\begin{aligned} \mathbf{u}_S(x, y) &:= (e^z \sin y, -e^z \sin y, e^z \cos y - e^z \cos x), \\ p_S(x, y) &:= -\frac{1}{2}e^{2z} + \frac{1}{4}(e^2 - 1), \\ \mathbf{u}_D(x, y) &:= (\cos(\pi z) \sin(\pi x) \sin(\pi y), \sin(\pi z) \cos(\pi x) \sin(\pi y), -2 \sin(\pi z) \sin(\pi x) \cos(\pi y)), \\ p_D(x, y) &:= \sin(\pi z) \sin(\pi x) \cos(\pi y). \end{aligned}$$

All calculations were performed using the approximation spaces  $\mathbf{H}_{h,1} \times Q_{h,1}$ .




Figure 11: Configuration of the computational domain  $\Omega$ .

In Figure 12 we present the convergence orders obtained using our stabilized scheme (3.6).




Figure 12: Convergence of the velocities (left) and pressures (right). Here the interpolation order is  $k = 1$ .

In Table 3 we depict the approximation error as well as the estimator  $\eta$  when  $h$  goes to 0. Note that the effectivity index  $E$  remains bounded close to a constant.

$h$	$\ (\mathbf{u} - \mathbf{u}_h, p - p_h)\ $	order	$\eta$	order	$E$
0.4330127	0.7736884	—	0.6859547	—	0.8866034
0.2165064	0.3876403	0.9970342	0.3053558	1.1676221	0.7877298
0.1082532	0.1912973	1.0189023	0.1472243	1.0524756	0.7696100
0.0541266	0.0949678	1.0103059	0.0728948	1.0141285	0.7675735
0.0270633	0.0473340	1.0045612	0.0362839	1.0064853	0.7665505

Table 3: Effectivity index in a quasi-uniform refinement for  $(\mathbf{u}_h, p_h) \in \mathbf{H}_{h,1} \times \mathcal{Q}_{h,1}$ .

Finally, in Figures 13 and 14 we compare the isovalues of the pressures and the velocities magnitude obtained by our adapted scheme with those of the exact solutions. Note the good agreement in both solutions.




Figure 13: Isovalues of the pressure  $p_{h,D}$  (top left) versus the exact pressure  $p_D$  (bottom left) and isovalues of the velocity magnitude  $|u_{h,D}|$  (top right) versus exact velocity magnitude  $|u_D|$  (bottom right). Here  $k = 1$ .




Figure 14: Isovalues of the pressure  $p_{h,S}$  (top left) versus the exact pressure  $p_S$  (bottom left) and isovalues of the velocity magnitude  $|u_{h,S}|$  (top right) versus exact velocity magnitude  $|u_S|$  (bottom right). Here  $k = 1$ .

## Conclusions

In this work, we have introduced a new stabilized finite element method for the coupled Stokes–Darcy problem. Stability and optimal convergence have been proven even when the polynomial degree used to approximate the pressures is the same as the one used to approximate the velocities. In addition, we have provided a residual-based error estimator composed by two estimators localized on each subdomain. For this error estimator, we proved its efficiency and reliability up to a high order term. These estimates have been verified by numerical experiments in 2D and 3D. The numerical examples presented testify that this approach is effective, for academic illustrations and realistic applications. Firstly, we can look that optimal rates are achieved, thus verifying the theoretical a priori results. Furthermore, it was verified the good performance of the estimator with respect to the viscosity  $\nu$ , adapting on corner singularities and on interfaces. Finally, It was shown that the effectivity indexes are bounded as proven theoretically.

## Acknowledgements

The first author was partially supported by ANID-Chile through the projects Centro de Modelamiento Matemático (FB210005) of the PIA Program: Concurso Apoyo a Centros Científicos y Tecnológicos de Excelencia con Financiamiento Basal, and Fondecyt Regular No 1211649. The third author was partially supported by Dirección de Investigación of the Universidad Católica de la Santísima Concepción through project DIREG 01/2020 and Fondecyt Regular No 1211649. Finally, the fourth author was partially supported by Dirección de Investigación of the Universidad Católica de la Santísima Concepción through project DIREG 01/2020.

## Appendix A. Inf–sup condition for $B$

The next result, used in Section 5, is an adaptation of the arguments presented in [4, Lemma 11].

**Lemma 14.** *There exists a positive constant  $C$ , depending on  $\Omega$ , such that*

$$\sup_{(\mathbf{v}, q) \in \mathbf{H} \times Q} \frac{B((\mathbf{u}, p); (\mathbf{v}, q))}{\|(\mathbf{v}, q)\|} \geq C \|(\mathbf{u}, p)\|,$$

for all  $(\mathbf{u}, p) \in \mathbf{H} \times Q$ .

*Proof.* Using to the definition of  $B$  given in (2.4) and Lemma 1, we have

$$B((\mathbf{u}, p); (\mathbf{u}, p)) \geq C_1 \nu \kappa \|\mathbf{u}_S\|_{1, \Omega_S}^2 + C_2 \nu \|\mathbf{u}_D\|_{0, \Omega_D}^2 + C_3 \frac{\kappa^2}{\nu} |p_D|_{1, \Omega_D}^2. \quad (\text{A.1})$$

On the other hand, by [36, Proposition 5.3.2] we know that for a given  $p_S \in Q^S$  there exists  $\mathbf{w}_S \in \mathbf{H}^S$ ,  $\mathbf{w}_S \neq \mathbf{0}$  and a positive constant  $\zeta$ , such that

$$-(p_S, \nabla \cdot \mathbf{w}_S) \geq \zeta \|\mathbf{w}_S\|_{1, \Omega_S} \|p_S\|_{0, \Omega_S}.$$

Now, if we define  $(\mathbf{w}, 0) := ((\mathbf{w}_S, \mathbf{0}), (0, 0))$ , assuming that  $\|\mathbf{w}_S\|_{1, \Omega_S} = \|p_S\|_{0, \Omega_S}$ , by the Young’s inequality, a trace inequality and Lemma 1, we obtain

$$\begin{aligned} & B((\mathbf{u}, p); (\mathbf{w}, 0)) \\ & \geq -C_4 \nu \kappa \|\mathbf{u}_S\|_{1, \Omega_S} \|\mathbf{w}_S\|_{1, \Omega_S} - C_5 \kappa^{1/2} \|\mathbf{u}_S\|_{1, \Omega_S} \|\mathbf{w}_S\|_{1, \Omega_S} - C_6 \kappa |p_D|_{1, \Omega_D} \|\mathbf{w}_S\|_{1, \Omega_S} + \zeta \kappa \|p_S\|_{0, \Omega_S}^2 \\ & \geq -\frac{\nu \kappa}{2} \left( \frac{C_4}{\gamma_1} + \frac{C_5}{\gamma_2} \right) \|\mathbf{u}_S\|_{1, \Omega_S}^2 + \kappa \left( \zeta - \frac{C_4 \gamma_1 \nu}{2} - \frac{C_5 \gamma_2}{2 \nu \kappa} - \frac{C_6 \gamma_3 \nu}{2 \kappa} \right) \|p_S\|_{0, \Omega_S}^2 - \frac{C_6}{2 \gamma_3} \frac{\kappa^2}{\nu} |p_D|_{1, \Omega_D}^2 \\ & \geq -C_7 \nu \kappa \|\mathbf{u}_S\|_{1, \Omega_S}^2 + C_8 \kappa \|p_S\|_{0, \Omega_S}^2 - C_9 \frac{\kappa^2}{\nu} |p_D|_{1, \Omega_D}^2, \end{aligned} \quad (\text{A.2})$$

where we chose  $\gamma_1, \gamma_2, \gamma_3$  small enough.

Let  $(\mathbf{v}, q) := (\mathbf{u}, p) + \delta(\mathbf{w}, 0)$ . Combining (A.1) with (A.2), we deduce that

$$\begin{aligned}
& B((\mathbf{u}, p); (\mathbf{v}, q)) \\
& \geq C_1 \nu \kappa \|\mathbf{u}_S\|_{1, \Omega_S}^2 + C_2 \nu \|\mathbf{u}_D\|_{0, \Omega_D}^2 + C_3 \frac{\kappa^2}{\nu} |p_D|_{1, \Omega_D}^2 - C_7 \delta \nu \kappa \|\mathbf{u}_S\|_{1, \Omega_S}^2 + C_8 \delta \kappa \|p_S\|_{0, \Omega_S}^2 - C_9 \delta \frac{\kappa^2}{\nu} |p_D|_{1, \Omega_D}^2 \\
& \geq \nu \kappa (C_1 - \delta C_7) \|\mathbf{u}_S\|_{1, \Omega_S}^2 + C_2 \nu \|\mathbf{u}_D\|_{0, \Omega_D}^2 + \frac{\kappa^2}{\nu} (C_3 - \delta C_9) |p_D|_{1, \Omega_D}^2 + C_8 \delta \kappa \|p_S\|_{0, \Omega_S}^2 \\
& \geq C_{10} \|(\mathbf{u}, p)\|^2,
\end{aligned} \tag{A.3}$$

choosing  $0 < \delta < \min \left\{ \frac{C_1}{C_7}, \frac{C_3}{C_9}, \frac{1}{C_8}, \frac{1}{\nu^{1/2}} \right\}$ .

Finally, using triangle inequality yields

$$\|(\mathbf{v}, q)\| \leq \|(\mathbf{u}, p)\| + \delta \|(\mathbf{w}, 0)\| = \|(\mathbf{u}, p)\| + \delta \nu^{1/2} \kappa^{1/2} \|p_S\|_{0, \Omega_S} \leq C_{11} \|(\mathbf{u}, p)\|,$$

thus, using (A.3), we get to the desired result.  $\square$

## References

- [1] Agmon, S., 2010. Lectures on elliptic boundary value problems. AMS Chelsea Publishing, Providence, RI. doi:10.1090/chel/369.
- [2] Amara, M., Capatina, D., Lizaik, L., 2009. Coupling of Darcy-Forchheimer and compressible Navier-Stokes equations with heat transfer. SIAM J. Sci. Comput. 31, 1470–1499. doi:10.1137/070709517.
- [3] Araya, R., Barrenechea, G.R., Poza, A.H., 2008. An adaptive stabilized finite element method for the generalized Stokes problem. J. Comput. Appl. Math. 214, 457–479. doi:10.1016/j.cam.2007.03.011.
- [4] Araya, R., Cárcamo, C., Poza, A.H., 2021. An adaptive stabilized finite element method for the Darcy’s equations with pressure dependent viscosities. Comput. Methods Appl. Mech. Engrg. 387, 114100. doi:10.1016/j.cma.2021.114100.
- [5] Armentano, M., Stockdale, M., 2019. A unified mixed finite element approximations of the Stokes-Darcy coupled problem. Comput. Math. Appl. 77, 2568–2584. doi:10.1016/j.camwa.2018.12.032.
- [6] Badea, L., Discacciati, M., Quarteroni, A., 2010. Numerical analysis of the Navier-Stokes/Darcy coupling. Numer. Math. 115, 195–227. doi:10.1007/s00211-009-0279-6.
- [7] Badia, S., Codina, R., Badia, S., Codina, R., 2009. Unified stabilized finite element formulations for the Stokes and the Darcy problems. SIAM J. Numer. Anal. 47, 1971–2000. doi:10.1137/08072632X.
- [8] Barrenechea, G.R., Valentin, F., 2002. An unusual stabilized finite element method for a generalized Stokes problem. Numer. Math. 92, 653–677. doi:10.1007/s002110100371.
- [9] Beavers, G.S., Joseph, D.D., Beavers, G., Joseph, D., 1967. Boundary conditions at a naturally permeable wall. J. Fluid Mech. 30, 197–207. doi:10.1017/S0022112067001375.
- [10] Bernardi, C., Hecht, F., Nouri, F., 2010. A new finite-element discretization of the Stokes problem coupled with the Darcy equations. IMA J. Numer. Anal. 30, 61–93. doi:10.1093/imanum/drn054.
- [11] Blank, L., Caiazzo, A., Chouly, F., Lozinski, A., Mura, J., Blank, L., Caiazzo, A., Chouly, F., Lozinski, A., Mura, J., 2018. Analysis of a stabilized penalty-free Nitsche method for the Brinkman, Stokes, and Darcy problems. ESAIM Math. Model. Numer. Anal. 52, 2149–2185. doi:10.1051/m2an/2018063.
- [12] Bochev, P.B., Dohrmann, C.R., Bochev, P.B., Dohrmann, C.R., 2006a. A computational study of stabilized, low-order  $C^0$  finite element approximations of Darcy equations. Comput. Mech. 38, 310–333. doi:10.1007/s00466-006-0036-y.
- [13] Bochev, P.B., Dohrmann, C.R., Gunzburger, M.D., Bochev, P.B., Dohrmann, C.R., Gunzburger, M., 2006b. Stabilization of low-order mixed finite elements for the Stokes equations. SIAM J. Numer. Anal. 44, 82–101. doi:10.1137/S0036142905444482.
- [14] Boffi, D., Brezzi, F., Fortin, M., Boffi, D., Brezzi, F., Fortin, M., 2013. Mixed finite element methods and applications. volume 44 of *Springer Series in Computational Mathematics*. Springer, Heidelberg. doi:10.1007/978-3-642-36519-5.
- [15] Braack, M., Schieweck, F., 2011. Equal-order finite elements with local projection stabilization for the Darcy-Brinkman equations. Comput. Methods Appl. Mech. Engrg. 200, 1126–1136. doi:10.1016/j.cma.2010.06.034.
- [16] Brenner, S.C., Scott, L.R., 2008. The mathematical theory of finite element methods. volume 15 of *Texts in Applied Mathematics*. Third ed., Springer, New York. doi:10.1007/978-0-387-75934-0.
- [17] Burman, E., Hansbo, P., 2007. A unified stabilized method for Stokes’ and Darcy’s equations. J. Comput. Appl. Math. 198, 35–51. doi:10.1016/j.cam.2005.11.022.
- [18] Camaño, J., Gatica, G.N., Oyarzúa, Ruiz-Baier, R., Venegas, P., 2015. New fully-mixed finite element methods for the Stokes-Darcy coupling. Comput. Methods Appl. Mech. Engrg. 295, 362–395. doi:10.1016/j.cma.2015.07.007.
- [19] Carvalho, P.G.S., Devloo, P.R.B., Gomes, S.M., Carvalho, P., Devloo, P., Gomes, S., 2020. On the use of divergence balanced  $\mathbf{H}(\text{div})$ - $L^2$  pair of approximation spaces for divergence-free and robust simulations of Stokes, coupled Stokes-Darcy and Brinkman problems. Math. Comput. Simulation 170, 51–78. doi:10.1016/j.matcom.2019.09.002.

- [20] Ciarlet, P.G., 1997. Mathematical elasticity. Vol. II. Theory of plates. volume 27 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam.
- [21] Clément, P., 1975. Approximation by finite element functions using local regularization. R.A.I.R.O. Anal. Numer. 9, 77–84. doi:10.1051/m2an/197509R200771.
- [22] Cockburn, B., Sayas, F., 2014. Divergence-conforming HDG methods for Stokes flows. Math. Comp. 83, 1571–1598. doi:10.1090/S0025-5718-2014-02802-0.
- [23] Discacciati, M., Miglio, E., Quarteroni, A., 2002. Mathematical and numerical models for coupling surface and ground-water flows. Appl. Numer. Math. 43, 57–74. doi:10.1016/S0168-9274(02)00125-3. 19th Dundee Biennial Conference on Numerical Analysis.
- [24] Ern, A., Guermond, J.L., 2004. Theory and practice of finite elements. volume 159 of *Applied Mathematical Sciences*. Springer-Verlag, New York. doi:10.1007/978-1-4757-4355-5.
- [25] Ervin, V.J., Jenkins, E.W., Sun, S., 2009. Coupled generalized nonlinear Stokes flow with flow through a porous medium. SIAM J. Numer. Anal. 47, 929–952. doi:10.1137/070708354.
- [26] Franca, L., Stenberg, R., 1991. Error analysis of Galerkin least squares methods for the elasticity equations. SIAM J. Numer. Anal. 28, 1680–1697. doi:10.1137/0728084.
- [27] Gatica, G.N., Meddahi, S., Oyarzúa, R., 2009. A conforming mixed finite-element method for the coupling of fluid flow with porous media flow. IMA J. Numer. Anal. 29, 86–108. doi:10.1093/imanum/drm049.
- [28] Gatica, G.N., Oyarzúa, R., Sayas, F., 2011. A residual-based a posteriori error estimator for a fully-mixed formulation of the Stokes-Darcy coupled problem. Comput. Methods Appl. Mech. Engrg. 200, 1877–1891. doi:10.1016/j.cma.2011.02.009.
- [29] Hansbo, P., Larson, M.G., Massing, A., Hansbo, P., Larson, M., Massing, A., 2017. A stabilized cut finite element method for the Darcy problem on surfaces. Comput. Methods Appl. Mech. Engrg. 326, 298–318. doi:10.1016/j.cma.2017.08.007.
- [30] Hanspal, N., Waghode, A., Nassehi, V., Wakeman, R., Hanspal, N., Waghode, A., Nassehi, V., Wakeman, R., 2009. Development of a predictive mathematical model for coupled Stokes/Darcy flows in cross-flow membrane filtration. Chem. Eng. J. 149, 132–142. doi:10.1016/j.cej.2008.10.012.
- [31] Layton, W.J., Schieweck, F., Yotov, I., Layton, W., Schieweck, F., Yotov, I., 2002. Coupling fluid flow with porous media flow. SIAM J. Numer. Anal. 40, 2195–2218. doi:10.1137/S0036142901392766.
- [32] Levy, T., Sánchez-Palencia, E., 1975. On boundary conditions for fluid flow in porous media. Internat. J. Engrg. Sci. 13, 923–940. doi:10.1016/0020-7225(75)90054-3.
- [33] Li, M., Shi, D., Dai, Y., 2016. Stabilized low order finite elements for Stokes equations with damping. J. Math. Anal. Appl. 435, 646–660. doi:10.1016/j.jmaa.2015.10.040.
- [34] Logg, A., Wells, G., Mardal, K. (Eds.), 2012. Automated solution of differential equations by the finite element method. The FEniCS book. volume 84 of *Lecture Notes in Computational Science and Engineering*. Springer-Verlag Berlin Heidelberg. doi:10.1007/978-3-642-23099-8.
- [35] Mishra, M., 2019. Hydrodynamic instabilities in porous media flows. Math. Student 88, 49–66.
- [36] Quarteroni, A., Valli, A., 1999. Domain Decomposition Methods for Partial Differential Equations. Oxford University Press.
- [37] Rui, H., Zhang, R., Rui, H., Zhang, R., 2009. A unified stabilized mixed finite element method for coupling Stokes and Darcy flows. Comput. Methods Appl. Mech. Engrg. 198, 2692–2699. doi:10.1016/j.cma.2009.03.011.
- [38] Urquiza, J.M., N’Dri, D., Garon, A., Delfour, M.C., Urquiza, J.M., N’Dri, D., Garon, A., Delfour, M.C., 2008. Coupling Stokes and Darcy equations. Appl. Numer. Math. 58, 525–538. doi:10.1016/j.apnum.2006.12.006.
- [39] Xie, C.M., Luo, Y., Feng, M.F., Xie, C.M., Luo, Y., Feng, M.F., 2011. Analysis of a unified stabilized finite volume method for the Darcy-Stokes problem. Math. Numer. Sin. 33, 133–144.
- [40] Ye, X., Zhang, S., Ye, X., S., Z., 2021. A stabilizer free WG method for the Stokes equations with order two superconvergence on polytopal mesh. Electron. Res. Arch. 29, 3609–3627. doi:10.3934/era.2021053.
- [41] Yu, J., Al Mahbub, M.A., Shi, F., Zheng, H., 2018. Stabilized finite element method for the stationary mixed Stokes-Darcy problem. Adv. Difference Equ. 2018, 346. doi:10.1186/s13662-018-1809-2.
- [42] Zhao, L., Park, E.J., Zhao, L., Park, E., 2020. A lowest-order staggered DG method for the coupled Stokes-Darcy problem. IMA J. Numer. Anal. 40, 2871–2897. doi:10.1093/imanum/drz048.

# Centro de Investigación en Ingeniería Matemática (CI<sup>2</sup>MA)

## PRE-PUBLICACIONES 2021 - 2022

- 2021-26 TOMÁS BARRIOS, EDWIN BEHRENS, ROMMEL BUSTINZA: *Numerical analysis of a stabilized mixed method applied to incompressible elasticity problems with Dirichlet and with mixed boundary conditions*
- 2021-27 LILIANA CAMARGO, MANUEL SOLANO: *A high order unfitted HDG method for the Helmholtz equation with first order absorbing boundary condition*
- 2021-28 ANA ALONSO-RODRIGUEZ, JESSIKA CAMAÑO: *A graph-based algorithm for the approximation of the spectrum of the curl operator*
- 2021-29 VERONICA ANAYA, RUBEN CARABALLO, RICARDO RUIZ-BAIER, HECTOR TORRES: *Augmented finite element formulation for the Navier-Stokes equations with vorticity and variable viscosity*
- 2021-30 CLAUDIO I. CORREA, GABRIEL N. GATICA: *On the continuous and discrete well-posedness of perturbed saddle-point formulations in Banach spaces*
- 2022-01 NESTOR SÁNCHEZ, TONATIUH SANCHEZ-VIZUET, MANUEL SOLANO: *Afternote to Coupling at a distance: convergence analysis and a priori error estimates*
- 2022-02 GABRIEL N. GATICA, CRISTIAN INZUNZA, FILANDER A. SEQUEIRA: *A pseudostress-based mixed-primal finite element method for stress-assisted diffusion problems in Banach spaces*
- 2022-03 ELIGIO COLMENARES, RICARDO OYARZÚA, FRANCISCO PIÑA: *A fully-DG method for the stationary Boussinesq system*
- 2022-04 JUAN MANUEL CÁRDENAS, MANUEL SOLANO: *A high order unfitted hybridizable discontinuous Galerkin method for linear elasticity*
- 2022-05 JAIME MANRÍQUEZ, NGOC-CUONG NGUYEN, MANUEL SOLANO: *A dissimilar non-matching HDG discretization for Stokes flows*
- 2022-06 RAIMUND BÜRGER, STEFAN DIEHL, MARÍA CARMEN MARTÍ, YOLANDA VÁSQUEZ: *A degenerating convection-diffusion system modelling froth flotation with drainage*
- 2022-07 RODOLFO ARAYA, CRISTIAN CÁRCAMO, ABNER POZA, EDUARDO VINO: *An adaptive stabilized finite element method for the Stokes-Darcy coupled problem*

Para obtener copias de las Pre-Publicaciones, escribir o llamar a: DIRECTOR, CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE, TEL.: 41-2661324, o bien, visitar la página web del centro: <http://www.ci2ma.udec.cl>





**CENTRO DE INVESTIGACIÓN EN  
INGENIERÍA MATEMÁTICA (CI<sup>2</sup>MA)  
Universidad de Concepción**



Casilla 160-C, Concepción, Chile  
Tel.: 56-41-2661324/2661554/2661316  
<http://www.ci2ma.udec.cl>

